

## Appendix II-G1

2019 Benthic Assessment Report - Buoy Installation Areas and Sites of Interest

# ATLANTIC SHORES OFFSHORE WIND

## Benthic Assessment Report – Buoy Installation Areas and Sites of Interest



19-P-204565 Atlantic Shores  
Benthic Assessment Report V5  
Buoy Installation Areas  
Terrasond - ASOW

February 14, 2020  
Updated: November 22, 2021

---

Prepared by:

**RPS Ocean Science**

Joseph Zottoli, Alicia Morandi, Paxton Albert, and Alex  
Sousa

55 Village Square Drive  
South Kingstown RI 02879

**T** +1 401 789 6224

**E** [Alicia.Morandi@rpsgroup.com](mailto:Alicia.Morandi@rpsgroup.com)

Prepared for:

**Terrasond – Atlantic Shores Offshore Wind**

## Contents

<b>1</b>	<b>INTRODUCTION</b> .....	<b>1</b>
<b>2</b>	<b>METHODS</b> .....	<b>1</b>
2.1	Field Survey .....	1
2.1.1	Drop-Down Video .....	1
2.1.2	Grab Sampling .....	3
2.2	Lab Analysis .....	4
2.2.1	Grain Size and TOC Analysis .....	4
2.2.2	Benthic Infauna Analysis .....	5
2.3	Video Post-Processing .....	5
2.4	Benthic Community Data Post-Processing .....	7
2.4.1	Taxonomic Composition .....	7
2.4.2	Richness, Diversity, and Evenness .....	7
<b>3</b>	<b>RESULTS</b> .....	<b>8</b>
3.1	Visual Analysis .....	9
3.1.1	IA Sites .....	9
3.1.2	Northern SOI Sites .....	16
3.1.3	Central SOI Sites .....	18
3.1.4	Southern SOI Sites .....	19
3.2	Sediment Analyses .....	21
3.3	Benthic Community Analysis .....	23
3.3.1	Taxonomic Composition .....	23
3.3.2	Richness, Diversity, and Evenness .....	36
<b>4</b>	<b>CMECS CLASSIFICATIONS</b> .....	<b>38</b>
<b>5</b>	<b>REFERENCES</b> .....	<b>43</b>
	<b>APPENDIX A FIELD DATASHEETS</b> .....	<b>44</b>
	<b>APPENDIX B PHOTOMICROSCOPY IMAGES</b> .....	<b>50</b>

## Figures

Figure 1:	Map of sample stations. Note that “B” delineates buoy IAs (e.g., B3-2 is IA site 3 station 2) and “S” delineates SOIs.....	2
Figure 2:	Still image of DDV associated with benthic grab station 1 at IA 1 taken from 0.46 m above the seafloor; field of view: 0 cm <sup>2</sup> .....	10
Figure 3:	Still image of DDV associated with benthic grab station 2 at IA 1 taken from 0.46 m above the seafloor; field of view: 0 cm <sup>2</sup> .....	10
Figure 4:	Still image of DDV associated with benthic grab station 1 at IA 2 taken from 0.46 m above the seafloor; field of view: 2,445 cm <sup>2</sup> .....	12
Figure 5:	Still image of DDV associated with benthic grab station 2 at IA 2 taken from 0.46 m above the seafloor; field of view: 2,223 cm <sup>2</sup> .....	12
Figure 6:	Still image of DDV associated with benthic grab station 1 at IA 3 taken from 0.46 m above the seafloor; field of view: 2,726 cm <sup>2</sup> .....	13
Figure 7:	Still image of DDV associated with benthic grab station 2 at IA 3 taken from 0.46 m above the seafloor; field of view: 2,229 cm <sup>2</sup> .....	14
Figure 8:	Still image of DDV associated with benthic grab station 1 at IA 4 taken from 0.46 m above the seafloor; field of view: 0 cm <sup>2</sup> .....	15
Figure 9:	Still image of DDV associated with benthic grab station 2 at IA 4 taken from 0.46 m above the seafloor; field of view: 2,563 cm <sup>2</sup> .....	16
Figure 10:	Still image of DDV associated with benthic grab station SOI 7 taken from 0.46 m above the seafloor; field of view: 0 cm <sup>2</sup> .....	17
Figure 11:	Still image of DDV associated with benthic grab station SOI 8 taken from 0.46 m above the seafloor; field of view: 1,293 cm <sup>2</sup> .....	17
Figure 12:	Still image of DDV associated with benthic grab station SOI 4 taken from 0.46 m above the seafloor; field of view: 1,393 cm <sup>2</sup> .....	19
Figure 13:	Still image of DDV associated with benthic grab station SOI 1 taken from 0.46 m above the seafloor; field of view: 2,268 cm <sup>2</sup> .....	20
Figure 14:	Still image of DDV associated with benthic grab station SOI 2 taken from 0.46 m above the seafloor; field of view: 2,146 cm <sup>2</sup> .....	20
Figure 15:	Grain size composition at each sample station. Stations located in the WTA are denoted with an asterisk (*). .....	22
Figure 16:	Proportional abundance and proportional number of unique taxa (species or LPTL) for each phylum collected in all benthic grab samples. Results presented as percentage of total. ....	24
Figure 17:	Richness (top) Diversity (middle) and Evenness (bottom) index values for each IA and SOI. Stations located in the WTA are denoted with an asterisk (*). ....	38
Figure 18:	Substrate CMECS classifications by sample site. Note that “B” delineates buoy IAs (e.g., B3-2 is IA site 3 station 2) and “S” delineates SOIs. ....	39

## Tables

Table 1:	Grab-sample station locations and characteristics. Coordinates are North American Datum of 1983 (NAD 83), EPSG 4269. ....	8
Table 2:	Visual characterization of still images associated with two stations at IA 2. ....	11
Table 3:	Visual characterization of still images associated with two stations at IA 3. ....	13
Table 4:	Visual characterization of still images associated with two stations at IA 4. ....	14
Table 5:	Visual characterization of still images associated with northern stations at SOI 7 and SOI 8. ....	16
Table 6:	Visual characterization of still images associated with central stations at SOI 3, 4, 5, and 6. ....	18
Table 7:	Visual characterization of still images associated with southern stations at SOI 1 and SOI 2. ....	19
Table 8:	Grain size composition and total organic carbon content of grab samples. ....	21
Table 9:	Laser diffraction results showing sizes ( $\mu\text{m}$ ) which 10% ( $D_v [10]$ ), 50% ( $D_v [50]$ ), and 90% ( $D_v [90]$ ) of particles in the distribution are smaller than and the volume-weighted mean ( $D[4,3]$ ). ....	22
Table 10:	Phyla present in the fourteen benthic grab samples. ....	23
Table 11:	Density (# of individuals per $0.05 \text{ m}^2$ ) of each Phylum present within each grab sample and the mean density for IA grab sample stations. ....	24
Table 12:	Density of each phyla and taxa (LPTLs) collected at the two grab stations (data combined) in IA 1. ....	26
Table 13:	Density of each phyla and taxa (LPTL) collected at the two grab stations (data combined) in IA 2. ....	28
Table 14:	Density of each phyla and taxa (LPTL) collected at the two grab stations (data combined) in IA 3. ....	29
Table 15:	Density of each phyla and taxa (LPTL) collected at the two grab stations (data combined) in IA 4. ....	30
Table 16:	Density of each phyla and taxa (LPTL) collected at the grab station at SOI 1. ....	31
Table 17:	Density of each phyla and taxa (LPTL) collected at the grab station at SOI 2. ....	32
Table 18:	Density of each phyla and taxa (LPTL) collected at the grab station at SOI 3. ....	33
Table 19:	Density of each phyla and taxa (LPTL) collected at the grab station at SOI 4. ....	34
Table 20:	Density of each phyla and taxa (LPTL) collected at the grab station at SOI 5. ....	35
Table 21:	Density of each phyla and taxa (LPTL) collected at the grab station at SOI 6. ....	36
Table 22:	Community composition parameters calculated for each grab sample station and for each IA. Combined values for individual stations within the four IAs were either averaged ( $\bar{x}$ ) or summed ( $\Sigma$ ) depending on parameter. ....	37
Table 23:	Images of grab samples prior to processing, along with CMECS classifications (Substrate & Biological modifier). ....	40
Table 24:	CMECS hierarchical classification of substrates collected at each station (S) in each IA. Stations located in the WTA are denoted with an asterisk (*). ....	42

## Table of Abbreviations

ASOW	Atlantic Shores Offshore Wind
BOEM	Bureau of Ocean Energy Management
cm	Centimeters
CMECS	Coastal and Marine Ecological Classifications Standards
DDV	Drop Down Video
Dv	Volume Diameter
EPA	United States Environmental Protection Agency
EPSG	European Petroleum Survey Group code used for coordinate systems
EST	Eastern Standard Time
GPS	Global Positioning System
IA	Buoy Installation Area
kg	Kilogram
LPTL	Lowest Practical Taxonomic Level
m	Meters
mg	Milligrams
SOI	Sites of Interest
TOC	Total Organic Carbon

# 1 INTRODUCTION

RPS was contracted by Terrasond to conduct benthic video and grab sampling, post-process the video collected, and compile this benthic assessment report for surveys conducted within four metocean data collection Buoy Installation Areas (IAs) and eight other Sites of Interest (SOIs) in the Atlantic Shores Offshore Wind (ASOW) Lease Area (OCS-A 0499) located offshore of New Jersey. The grab samples and video imagery data conclusions presented here will support interpretation of geophysical data to characterize surficial sediment conditions and classify the benthic habitat according to the Coastal and Marine Ecological Classifications Standards (CMECS; FGDC 2012) in accordance with Bureau of Ocean Energy Management (BOEM) guidelines. This report provides:

- A description of the benthic grab sampling methods, results, and analysis;
- The analysis of benthic grab sampling results using some key statistical analyses such as taxa richness, density per cubic meter, community composition, etc.;
- A description and analysis of the video data collected; and
- CMECS classifications of each sample site based on the video and grain size analyses.

The original version of this report was submitted as Appendix II-G1 of the Atlantic Shores Offshore Wind Construction and Operations Plan (COP) in March of 2021. This updated version contains substrate classification methods that are more consistent with the 2020 Benthic Assessment Report and 2021 Towed Video Report and analyses on the Project. Laser diffraction data with sieve sizes that better aligned to the Wentworth grain size class scale were used for classification instead of the original ASTM particle size distribution method. This has led to a more accurate representation of the percent composition of gravel in each sample. In addition, updated CMECS classifications based on the NMFS recommendations published in March 2021 were applied. The report has also been updated to reflect the Atlantic Shores currently proposed Project 1 and Project 2 boundaries as presented in the COP. The Project 1 and Project 2 boundaries within the Lease Area are collectively referred to herein as the Wind Turbine Area (WTA).

## 2 METHODS

### 2.1 Field Survey

#### 2.1.1 Drop-Down Video

Drop-down video (DDV) was taken in conjunction with grab samples to aid in sample collection and visual habitat classification on October 12-13, 2019. Two samples were taken from each IA with 8 additional grabs at SOIs (Figure 1). The video camera was equipped with an altimeter to record distance above sea floor, temperature probe, parallel-mounted scaling lasers 0.184 meter (m) apart, lights, and a 300-foot (ft)



length of cable that provided real-time viewing of images from the vessel. A silver-colored hook was suspended 0.70 m below the camera with string to standardize image distance and is visible near the center or right side of some images. Due to poor visibility, images were captured just as the grab sampler reached bottom instead of stopping at the standard distance. A YSI EXO sensor with a pH sensor was deployed separately to record pH from a depth of about 5 m at each site and water depth from shipboard sonar was recorded. The video camera was affixed to the grab sampler which was deployed by the Terrasond crew. The equipment was lowered until positioned just above the seafloor (when visibility allowed) and sites were identified as free of sensitive habitat and material prohibitive to sampling (e.g., boulders, large cobbles, other hard bottom, or debris). Samples were then collected at or within 10 m of selected locations.




Figure 1: Map of sample stations. Note that “B” delineates buoy IAs (e.g., B3-2 is IA site 3 station 2) and “S” delineates SOIs.

While viewing the video feed for sampling suitability (i.e., if the site can be sampled with a grab sampler or not due to presence of obstructions and/or sensitive habitat), the sample information (date, time, global positioning satellite [GPS] coordinates, station ID, depth, and video file name) and initial observations of sediment/seafloor characteristics were recorded to aid in post-processing of video data. Grab sample identification numbers were recorded with the video file metadata for sites where they were retrieved.

During video review, attention was given to noting if potentially sensitive benthic habitats (e.g., exposed hard bottom, seagrass/kelp/algal beds, coral species) were present, as per BOEM's Guidelines for Providing Benthic Habitat Survey Information for Renewable Energy Development on the Atlantic Outer Continental Shelf Pursuant to 30 CFR Part 585 (BOEM, 2019).

### 2.1.2 Grab Sampling

Benthic grab samples were acquired using an Ocean Instruments Salish Grab Standard SG-20 sampler. This grab is a modified version of a standard Van Veen sampler with a stainless-steel weighted frame and release system ideal for collection of sediments in soft to hard substrates with a penetration depth up to 20 centimeters (cm) and sampling area of 0.10 square meters (m<sup>2</sup>). A virtual GPS node was placed just off the stern of the vessel where the A-frame was deployed to obtain GPS coordinates for each sample. The actual location on the seafloor at which the sample was taken may have differed from the GPS coordinates by a few meters.

Upon retrieval, the grab sampler was examined for sample acceptability. A sample was deemed acceptable if:

- Sample was more than 50% full;
- Sample was not over penetrated (i.e., not full to the top); and
- Surface structures were undisturbed and even (i.e., not slumped).

If a sample did not fulfill these requirements, the entire contents were returned to the water and another sample attempt was made. If three failed sample attempts occurred at one station, sampling moved on to the next station (no more than two fails occurred in any one sampling station). The results of each attempted grab were recorded in field notes.

Once an acceptable sample was obtained, the following steps were taken:

1. Overlying water was drained using a siphon;
2. A photograph was taken of the sample next to an identification label containing sample identification number and a plastic ruler inserted to record sample depth;
3. Field notes included descriptions of physical features (apparent redox potential discontinuity depth, depth of penetration, sediment color, texture, odor, surface features) and surface macrofauna (e.g. longfin squid), which were then returned to the water.

The grab sample was then divided in two sections using a plexiglass divider. One half of the sample was processed for physical analysis (sediment grain size and total organic carbon [TOC]). For the TOC and grain size analysis, almost the entire top 2 cm of sediment on one half of the sample were collected using a stainless-steel spoon and placed in glass jars; this was approximately 150 milliliters (ml) of sediment for

the grain size sample and 200 ml for the TOC sample. The grain size and TOC samples were stored on ice and sent to Eurofin Test America lab (5575 8<sup>th</sup> St E Tacoma, WA) for initial processing then to Particle Technology Labs (555 Rogers Street, Downers Grove, IL 60515) for completion of sediment analysis.

The other half of the grab sample (i.e., on the other side of the plexiglass divider) was measured for volume and processed for biological community analysis. The sample was then loaded onto a processing table and material washed in a 0.5-mm sieve, using seawater under gentle pressure. The seawater used for sample processing was filtered through a 0.5-mm mesh to prevent planktonic organisms from mixing with the benthic samples.

Organisms, shell fragments, and other material remaining on the sieve were placed into a plastic container using stainless-steel spoon and forceps as needed. The container was filled no more than one-half to two-thirds full with sample and seawater. If the quantity of sample exceeded this volume, it was placed in a second container. The sample was fixed/preserved with 10% buffered formalin solution by filling the remaining space within the bottle with solution. Containers were tightly sealed with tape and stored in a cooler at ambient temperature (not frozen or refrigerated). Prior to sieving the next sample, the sieve was cleaned by scrubbing with a stiff brush and backwashing with pressurized water. The infaunal benthic community samples were sent to EcoAnalysts (1420 S Blaine St ste. 14 Moscow, ID 83843) for processing. Grabs from two reconnaissance sites not in IAs (SOI 7 and SOI 8) were not sampled for infauna.

## 2.2 Lab Analysis

### 2.2.1 Grain Size and TOC Analysis

Grain size and TOC samples were initially analyzed by TestAmerica using ASTM D 422-63 Standard Method for Particle-size Analysis of Soils (ASTM, 2007). The TOC content of sediment samples was analyzed using EPA Method 9060 with results reported in milligrams per kilogram (mg/kg) and percent (EPA, 1986).

Laser diffraction was performed by Particle Technology Labs using the ISO 13320 Standard Method for Particle-size Analysis with a Malvern MasterSizer 3000 Laser diffractor. Data from this method was later found to align better with the Wentworth grain size scale and data were reanalyzed for the November 2021 update of this report using it instead of the TestAmerica results. Laser diffraction is a method of grain size analysis that involves passing a laser beam through sediment samples in order to accurately estimate the grain size distribution of the sediment sample. The principal theory that is used for the Malvern MasterSizer 3000 Laser Diffractor is the Mie Theory, which describes the relationship between scattered light angles and absorption to the relative volume of a particle. Particle Technology Labs then prepared a document that detailed the cumulative volume percentage of each grain size that was analyzed for each sediment sample.

While the majority of the sediment samples were made up of particle sizes that the laser diffraction method could register, larger particulates that exceeded the upper limit of the MasterSizer 3000 Laser Diffractor (3.5 mm) were still present in some samples. To characterize the relative volume percentages of these larger grains, wet sieving was required. The wet sieving was also performed by Particle Technology Labs and the results were then presented as cumulative volume percentages of each grain size.

### 2.2.2 Benthic Infauna Analysis

The benthic infauna analysis was conducted by EcoAnalysts according to the following steps:

1. Benthic infaunal samples were catalogued and verified against the Chain of Custody to ensure samples received match those listed in the shipment.
2. Samples were rinsed with freshwater to remove the formalin and transferred to 70 percent ethanol alcohol for sorting and storage.
3. Organisms were identified to the lowest practical taxonomic level (LPTL) (at least to Family) and counted by taxonomists using the most appropriate taxonomic references for the region (Bousfield, 1973; Cutler, 1994; Winston and Hayward, 2012).
4. Species classification and abundance were recorded in Project data sheets and summarized in both tabular and graphical formats.
5. Prior to performing the infaunal data analyses, the overall dataset was scanned for non-infaunal taxa (i.e., pelagic or planktonic organisms) that were excluded from all analyses; examples include chaetognaths, hyperiid amphipods, and decapod zoea/megalopae.
6. Calculations of abundance included all taxa occurring in each sample whether identified to species level or not.
7. Calculations based on species (diversity, evenness, and number of species) included only those taxa identified to species level.

## 2.3 Video Post-Processing

Post-processing of video data was conducted by RPS to provide:

- General characterization of substrate including bottom type, texture, micro-topography, and presence and approximate thickness (absent, light, moderate, or heavy) of sedimentation (“drape”) covering hard substrates;
- Evidence of benthic activity by organisms (burrows, trails, biogenic reefs);
- Identification of epibenthic macroinvertebrates larger than 4 cm (decapod crustaceans, mollusks [including squid egg mops], echinoderms) and habitat;
- Presence/evidence and general characterization of submerged aquatic vegetation (macroalgae, sea grass);

- Identification of fish and fish habitat (where feasible) as classified by Auster (1998), which can provide back-compatibility with prior sampling depending on what has been previously done in the region and is easily applicable to Essential Fish Habitat determination;
- Identification of organisms to the lowest practical taxonomic level (at least to Family) using standard taxonomic keys for the geographic area;
- Evidence of fishing activity, such as trawl scars, pots, and working nets; and
- Presence of derelict fishing gear, military expended materials, shipwrecks, cultural artifacts, or other marine debris.

All DDV stills were classified according to CMECS (FGDC, 2012). Auster (1998) classification is also included as it is indicative of overall habitat features that can be important to fish, while CMECS focuses more closely on grain size and composition. The BOEM Benthic Habitat guidelines (BOEM, 2019) also require that the developer characterize the benthic community composition which includes documentation of abundance, diversity, percent cover, and community structure. The following were recorded when present and identifiable:

- Characterization and delineation of any submerged aquatic vegetation (seagrass or macroalgae) that occurs within the area of potential adverse effect;
- Characterization and delineation of any hard bottom gradients of low to high relief such as coral (heads/reefs), rock or clay outcroppings, or other shelter-forming features; and
- Identification of communities of sessile and slow-moving marine invertebrates (clams, quahogs, mussels, polychaete worms, anemones, sponges, echinoderms) that may be within the area of potential adverse effect.

The video data were analyzed according to the following steps:

1. A single still image was analyzed from each site by selecting the first clear view of the seafloor at each grab site, with camera positioned approximately 46 cm above the seafloor.
2. The visible area of each still image was defined, measured, and reviewed for evidence of benthic species and activity, submerged aquatic vegetation (macroalgae, sea grass), fishing activity, derelict gear, military expended materials, shipwrecks, and other marine debris and presence/absence of these features were noted. In some instances of poor visibility, video was used in addition to still images to determine presence absence.
3. Selected stills were broadly characterized by texture, microtopography, presence/thickness of sedimentation over hard substrates (i.e., "drape"), and presence of coral heads/reefs, rock outcroppings, other shelter features, or Essential Fish Habitat for NMFS-designated species in the region.
4. The biological component was defined to furthest extent possible for each station when analyzed in conjunction with the grain size and benthic community results according to the CMECS (FGDC, 2012).

## 2.4 Benthic Community Data Post-Processing

The benthic community analysis was based on the benthic macroinfauna laboratory data from Ecoanalysts. Macroinfauna community statistics were calculated using species and abundance estimates in each sample, which were reported as count per 0.05 m<sup>2</sup> grab sample (i.e., half of the grab sample processed for biological analysis). Community composition parameters included: total abundance, number of phyla, number of taxa, Margalef's Richness Index, Shannon Diversity Index, and Pielou's Index of Evenness for each station and within each IA.

### 2.4.1 Taxonomic Composition

Taxa composition was assessed to characterize the high-level trends in taxa data. Taxa composition includes the relative proportions of taxonomic groups by number of identifiable taxa and number of individuals, used to evaluate dominance of common phyla across all samples. Taxa composition was summarized for both individual samples and across the four IAs.

### 2.4.2 Richness, Diversity, and Evenness

Species richness, evenness, and diversity are common ecological parameters used to measure the overall biodiversity of a community or discrete unit. Species richness is the number of unique species or taxonomic groups represented in an area of interest. In this assessment, species richness was calculated using Margalef's Richness Index (Formula 1) for each station and IA to acquire individual and average richness indices.

*Formula 1. Margalef's Richness Index (RI).*

$$RI = \frac{(S - 1)}{\ln(n)}$$

*Where:*

*S= the number of species*

*n= the total number of individuals in the sample*

*Interpretation: The higher the index, the greater the species richness.*

The diversity index for a community considers species richness and the proportion of each unique species. The Shannon Diversity Index (H'; Formula 2) was calculated using the number of each species, the proportion of each species relative to the total number of individuals, and the sum of the proportions. This index was used to assess diversity of each station and IA. The diversity index (H') increases with increasing species richness and evenness.

*Formula 2. H'- Shannon Diversity Index.*

$$H' = - \sum_{i=1}^R p_i \ln(p_i)$$

*Where:*

*p<sub>i</sub> is the proportion of individuals belonging to the with species in the dataset of interest*

*Interpretation: The greater the H', the greater the richness and evenness.*

## ASOW BENTHIC ASSESSMENT REPORT

Evenness of a community refers to the similarity in abundances of different species comprising a population or sample. Pielou's Index of Evenness includes  $H'$  (Shannon-Weiner Diversity Index) in its calculation.

Formula 3.  $J'$ - Pielou's Index of Evenness.

$$J' = \frac{H'}{H_{Max}}$$

Where:

$H'$  is the Shannon- Weiner Diversity Index

$H_{Max}$  is the maximum possible value of  $H'$ , where each species occurs in equal abundances.

$$H_{Max} = \ln(s)$$

Where:  $s$  = Number of species

Interpretation:  $J'$  is constrained between 0 and 1. The greater the value of  $J'$ , the more evenness in the sample.

## 3 RESULTS

All samples were collected on October 12-13, 2019. Sampling stations were located in water between 21 and 35 m deep with bottom temperatures between 18.1°C and 18.9°C (Table 1).

Table 1: Grab-sample station locations and characteristics. Coordinates are North American Datum of 1983 (NAD 83), EPSG 4269.

IA or SOI	Station	Date	Time (EST)	Latitude	Longitude	Sonar-Based Water Depth (m)	Temp (°C)	Penetration Depth (cm)	Infaunal Sample Volume (m <sup>3</sup> )
IA 1*	1	10/12/19	17:25	39° 18.603' N	74° 6.5923' W	28.5	18.87	7.2	0.0021
	2	10/12/19	17:48	39° 18.5498' N	74° 6.4631' W	28.5	18.85	8.2	0.0021
IA 2	1	10/13/19	10:06	39° 33.9308' N	73° 58.1656' W	25.0	18.57	5.1	0.0015
	2	10/13/19	10:35	39° 33.9907' N	73° 58.1989' W	25.0	18.57	9.1	0.0042
IA 3*	1	10/12/19	19:19	39° 17.3562' N	73° 58.5943' W	34.5	18.40	13.9	0.0036
	2	10/12/19	20:06	39° 17.2256' N	73° 58.6107' W	33.5	18.77	7.6	0.0021
IA 4*	1	10/12/19	22:35	39° 12.1396' N	74° 5.0065' W	26.0	18.75	9.0	0.0024
	2	10/12/19	23:03	39° 12.1881' N	74° 4.7826' W	25.5	18.77	8.5	0.0024
SOI 1*	-	10/12/19	21:14	39° 14.1066' N	73° 59.616' W	26.0	18.70	7.3	0.003
SOI 2*	-	10/12/19	00:40	39° 15.4624' N	74° 12.3109' W	26.0	18.57	9.1	0.003
SOI 3*	-	10/13/19	15:19	39° 21.5007' N	74° 11.2525' W	21.0	18.85	9.5	0.002
SOI 4	-	10/13/19	13:43	39° 23.7691' N	74° 2.7639' W	26.0	18.75	9.5	0.003
SOI 5	-	10/13/19	14:43	39° 26.7895' N	74° 5.6972' W	21.0	18.40	11.5	0.003
SOI 6	-	10/13/19	12:39	39° 24.1021' N	73° 57.8008' W	28.0	18.59	9.2	0.0036
SOI 7	-	10/13/19	16:48	39° 33.8481' N	74° 2.1521' W	24.0	18.68	11.8	NA
SOI 8	-	10/13/19	18:07	39° 38.0361' N	73° 57.7192' W	26.0	18.14	8.5	NA

\* denotes benthic grab stations within the WTA

## 3.1 Visual Analysis

The following sections display and describe still images taken from the video camera affixed to the grab sampler (sampler visible in top of each image). Note that data overlaid on the screen may differ slightly from what is reported here due to a time lag in depth reporting (e.g., in Figure 8 the altimeter reads 1.35 m above the seafloor but the grab is resting on the seafloor). Parallel-mounted lasers visible in the center of the images are 0.184 m apart.

Still images were successfully captured and analyzed for 1 station at IA 2 (Figure 3), two stations at IA 3 (Figure 5 and Figure 6), and one station at IA 4 (Figure 8). Still images from other stations were unsuccessful due to extreme turbidity from a recent storm event but video observation during deployment provided limited coverage of the seafloor in some instances in which no notable findings were observed. All IAs appeared to have sandy bottom with varying degrees of shell debris and sand-ripple relief. SOI 8, the northernmost sample site, was the only site with clear presence of gravel. The field of view of each image was measured by calibrating images with the known distance between laser points and measuring the area of a polygon overlaid on the visible seafloor (i.e. unobstructed by grab or darkness) portion of each image. If an image did not include any visible seafloor, it was assigned a field of view of 0 cm<sup>2</sup>

### 3.1.1 IA Sites

#### 3.1.1.1 IA 1

Two sites were sampled in IA 1, located in the WTA, with limited visual coverage due to high turbidity (Figure 2 and Figure 3). Still images were not adequate for identifying habitat types and presence of other features of interest (i.e., aquatic vegetation, evidence of fishing activity, anthropogenic debris). The equipment visible in the still images is the grab sampler to which the camera was attached. The IA 1 sample sites are the most centrally located within the Lease Area, in block 6786H (station 1) and 6787E (station 2).





Figure 2: Still image of DDV associated with benthic grab station 1 at IA 1 taken from 0.46 m above the seafloor; field of view: 0 cm<sup>2</sup>.



Figure 3: Still image of DDV associated with benthic grab station 2 at IA 1 taken from 0.46 m above the seafloor; field of view: 0 cm<sup>2</sup>.

**3.1.1.2 IA 2**

IA 2 is the northernmost IA site in block 6489F, relatively near SOIs 7 and 8 (Table 2, Figure 4, Figure 5).

Table 2: Visual characterization of still images associated with two stations at IA 2.

Site	IA 2 Station 1	IA 2 Station 2
<b>General Characterization</b>	Despite high turbidity, some sparse shell hash was observed on light-colored sand. Video analysis aided in determining presence/absence of features.	Despite high turbidity, some sparse shell hash was observed on light-colored sand. Video analysis aided in determining presence/absence of features.
<b>Field of View</b>	2,445 cm <sup>2</sup>	2,223 cm <sup>2</sup>
<b>Biotic Benthic Activity</b>	None detected	None detected
<b>Epibenthic Macroinvertebrates and Fishes</b>	None detected	None detected
<b>Macroinvertebrate and Fish Habitat</b>	Flat Sand	Shell Aggregate
<b>Aquatic Vegetation</b>	None detected	None detected
<b>Evidence of Fishing Activity</b>	None detected	None detected
<b>Anthropogenic Debris</b>	None detected	None detected



Figure 4: Still image of DDV associated with benthic grab station 1 at IA 2 taken from 0.46 m above the seafloor; field of view: 2,445 cm<sup>2</sup>.



Figure 5: Still image of DDV associated with benthic grab station 2 at IA 2 taken from 0.46 m above the seafloor; field of view: 2,223 cm<sup>2</sup>.

**3.1.1.3 IA 3**

IA 3 is located along the southeast edge of the WTA in block 6789N of the Lease Area, nearest SOI 1 (Table 3, Figure 6, Figure 7).

Table 3: Visual characterization of still images associated with two stations at IA 3.

Site	IA 3 Station 1	IA 3 Station 2
<b>General Characterization</b>	Sandy with trace shell debris and presence of sand dollars, possible small sand waves/ripples	Sandy with sparse shell debris and presence of sand dollars, possible small sand waves/ripples
<b>Field of View</b>	2,726 cm <sup>2</sup>	2,229 cm <sup>2</sup>
<b>Biotic Benthic Activity</b>	None Detected	None Detected
<b>Epibenthic Macroinvertebrates and Fishes</b>	Sand Dollars	Sand Dollars
<b>Macroinvertebrate and Fish Habitat</b>	None Detected	None Detected
<b>Aquatic Vegetation</b>	None Detected	None Detected
<b>Evidence of Fishing Activity</b>	None Detected	None Detected
<b>Anthropogenic Debris</b>	None Detected	None Detected



Figure 6: Still image of DDV associated with benthic grab station 1 at IA 3 taken from 0.46 m above the seafloor; field of view: 2,726 cm<sup>2</sup>.



Figure 7: Still image of DDV associated with benthic grab station 2 at IA 3 taken from 0.46 m above the seafloor; field of view: 2,229 cm<sup>2</sup>.

### 3.1.1.4 IA 4

The first station sampled in IA 4 had limited visual coverage due to high turbidity. Still images were not adequate for identifying habitat types and presence of other objects of interest. IA 4 is the southernmost sample located in the WTA with stations in block 6887N (station 1) and 6887O (station 2) of the Lease Area (Table 4, Figure 8, Figure 9).

Table 4: Visual characterization of still images associated with two stations at IA 4.

Site	IA 4 Station 1	IA 4 Station 2
<b>General Characterization</b>	NA	Sandy with sparse shell debris and presence of sand dollars, possible small sand waves/ripples
<b>Field of View</b>	0 cm <sup>2</sup>	2,563 cm <sup>2</sup>
<b>Biotic Benthic Activity</b>	NA	None Detected
<b>Epibenthic Macroinvertebrates and Fishes</b>	NA	Sand dollars
<b>Macroinvertebrate and Fish Habitat</b>	NA	None Detected
<b>Aquatic Vegetation</b>	NA	None Detected
<b>Evidence of Fishing Activity</b>	NA	None Detected

Site	IA 4 Station 1	IA 4 Station 2
Anthropogenic Debris	NA	None Detected



Figure 8: Still image of DDV associated with benthic grab station 1 at IA 4 taken from 0.46 m above the seafloor; field of view: 0 cm<sup>2</sup>.



Figure 9: Still image of DDV associated with benthic grab station 2 at IA 4 taken from 0.46 m above the seafloor; field of view: 2,563 cm<sup>2</sup>.

### 3.1.2 Northern SOI Sites

Imaging from site SOI 7 had limited visual coverage due to high turbidity. Still images were not adequate for identifying habitat types and presence of other features of interest. SOI 7 is located in the northern portion of the Lease Area, in block 6488J. SOI 8 is the northernmost sample site within the Lease Area, in block 6389O (Table 5, Figure 10, Figure 11).

Table 5: Visual characterization of still images associated with northern stations at SOI 7 and SOI 8.

Site	SOI 7	SOI 8
<b>General Characterization</b>	NA	Gravel with shell debris
<b>Field of View</b>	0 cm <sup>2</sup>	1,293 cm <sup>2</sup>
<b>Biotic Benthic Activity</b>	NA	None Detected
<b>Epibenthic Macroinvertebrates and Fishes</b>	NA	None Detected
<b>Macroinvertebrate and Fish Habitat</b>	NA	None Detected
<b>Aquatic Vegetation</b>	NA	None Detected
<b>Evidence of Fishing Activity</b>	NA	None Detected
<b>Anthropogenic Debris</b>	NA	None Detected




Figure 10: Still image of DDV associated with benthic grab station SOI 7 taken from 0.46 m above the seafloor; field of view: 0 cm<sup>2</sup>.



Figure 11: Still image of DDV associated with benthic grab station SOI 8 taken from 0.46 m above the seafloor; field of view: 1,293 cm<sup>2</sup>.



### 3.1.3 Central SOI Sites

Imaging from site SOI 3 (located in the WTA and in block 6735G of the Lease Area), SOI 4 (block 6688E), SOI 5 (block 6637E), and SOI 6 (block 6689G) had limited visual coverage due to high turbidity. Still images were not adequate for identifying habitat types and presence of other features of interest at SOI 3, 5, or 6 but the seafloor was more visible at SOI 4; thus, only images from SOI 4 are included here (Table 6, Figure 12).

Table 6: Visual characterization of still images associated with central stations at SOI 3, 4, 5, and 6.

Site	SOI 3	SOI 4	SOI 5	SOI 6
<b>General Characterization</b>	NA	Sandy with trace shell debris	NA	NA
<b>Field of View</b>	0 cm <sup>2</sup>	1,393 cm <sup>2</sup>	0 cm <sup>2</sup>	0 cm <sup>2</sup>
<b>Biotic Benthic Activity</b>	NA	None Detected	NA	NA
<b>Epibenthic Macroinvertebrates and Fishes</b>	NA	None Detected	NA	NA
<b>Macroinvertebrate and Fish Habitat</b>	NA	None Detected	NA	NA
<b>Aquatic Vegetation</b>	NA	None Detected	NA	NA
<b>Evidence of Fishing Activity</b>	NA	None Detected	NA	NA
<b>Anthropogenic Debris</b>	NA	None Detected	NA	NA



Figure 12: Still image of DDV associated with benthic grab station SOI 4 taken from 0.46 m above the seafloor; field of view: 1,393 cm<sup>2</sup>.

### 3.1.4 Southern SOI Sites

SOI 1 and SOI 2 are in the southern portion of the Lease Area in the WTA. SOI 1 is located along the eastern edge of the Lease Area in the southern portion of block 6889A and SOI 2 is located along the western edge of the Lease Area in block 6835J (Table 7, Figure 13, Figure 14).

Table 7: Visual characterization of still images associated with southern stations at SOI 1 and SOI 2.

Site	SOI 1	SOI 2
<b>General Characterization</b>	Sandy with sparse shell debris, sand ripples present	Sandy with trace shell debris and presence of sand dollars, possible small sand waves/ripples
<b>Field of View</b>	2,268 cm <sup>2</sup>	2,146 cm <sup>2</sup>
<b>Biotic Benthic Activity</b>	None Detected	None Detected
<b>Epibenthic Macroinvertebrates and Fishes</b>	None Detected	Sand dollars
<b>Macroinvertebrate and Fish Habitat</b>	None Detected	None Detected
<b>Aquatic Vegetation</b>	None Detected	None Detected
<b>Evidence of Fishing Activity</b>	None Detected	None Detected
<b>Anthropogenic Debris</b>	None Detected	None Detected



Figure 13: Still image of DDV associated with benthic grab station SOI 1 taken from 0.46 m above the seafloor; field of view: 2,268 cm<sup>2</sup>.



Figure 14: Still image of DDV associated with benthic grab station SOI 2 taken from 0.46 m above the seafloor; field of view: 2,146 cm<sup>2</sup>.

### 3.2 Sediment Analyses

The following section presents grain size composition and TOC quantification results (Table 8 and Figure 15), laser diffraction (Table 9) and microscopy imaging results from the Particle Tech analyses (Appendix B). Note that the laser diffraction data from the Particle Tech analyses is presented as Dv(10), Dv(50), and Dv(90), which describe the size (volume) median that 10%, 50%, and 90% of the particles within the distribution are smaller than, while the grain size data is presented as composition percentage of each grain size through sieving.

When looking at the grain size analysis from Particle Tech, samples were generally composed of >90% sand, except for those from the northernmost sample sites in the Lease Area, SOI 7 (in block 6488J) and SOI 8 (in block 6389O), and IA 1-1 (in block 6786H), located in the WTA and central Lease Area. The sample from SOI 7 was composed of mostly fine to coarse sand (73%), while the SOI 8 sample was the most coarse, with 58% gravel (Table 8). Based on visual analysis, shell hash from multiple species including Atlantic surf clam (*Spisula solidissima*) and ocean quahog (*Arctica Islandica*) was present in many samples with densities ranging from trace to sparse. This shell hash is likely accounting for some to most of the gravel-sized grain components reported in the grain size results, as the lab analysis does not discern the difference between shell and gravel. The laser diffraction results indicate that nearly all samples were composed of mostly medium to coarse sand, except for SOI 8, where that sample was composed of mostly (>50%) gravel with a Dv (50) of 2.49 mm (Table 9).

Organic carbon content (i.e., TOC) was below the reporting limit of 2,000 mg/kg in all samples. Recent storms may have caused suspension of some organic matter into the water column, reducing TOC measures in the seabed.

Table 8: Grain size composition and total organic carbon content of grab samples.

IA or SOI	Station	Gravel (%)	Very Coarse/ Coarse Sand (%)	Medium Sand (%)	Fine/Very Fine Sand (%)	Mud (%)	NMFS-modified CMECS	Total Organic Carbon (mg/kg)
IA 1*	1	8	36	35	17	4	Gravelly Sand	Not Detected
	2	1	36	56	7	0	Medium Sand	Not Detected
IA 2	1	6	50	41	3	0	Gravelly Sand	Not Detected
	2	4	54	39	3	0	Very Coarse/ Coarse Sand	Not Detected
IA 3*	1	2	34	52	12	0	Medium Sand	Not Detected
	2	3	5	54	36	2	Medium Sand	Not Detected
IA 4*	1	1	33	59	7	0	Medium Sand	Not Detected
	2	9	36	47	8	0	Gravelly Sand	Not Detected
SOI 1*	-	1	41	52	6	0	Medium Sand	Not Detected
SOI 2*	-	0	12	70	18	0	Medium Sand	Not Detected
SOI 3*	-	1	42	51	6	0	Medium Sand	Not Detected
SOI 4	-	0	41	54	5	0	Medium Sand	Not Detected

## ASOW BENTHIC ASSESSMENT REPORT

IA or SOI	Station	Gravel (%)	Very Coarse/Coarse Sand (%)	Medium Sand (%)	Fine/Very Fine Sand (%)	Mud (%)	NMFS-modified CMECS	Total Organic Carbon (mg/kg)
SOI 5	-	1	70	28	1	0	Very Coarse/Coarse Sand	Not Detected
SOI 6	-	1	58	39	2	0	Very Coarse/Coarse Sand	Not Detected
SOI 7	-	27	30	36	7	0	Gravelly Sand	Not Detected
SOI 8	-	58	10	14	17	1	Sandy Gravel	Not Detected

\* denotes benthic grab stations within the WTA




Figure 15: Grain size composition at each sample station. Stations located in the WTA are denoted with an asterisk (\*).

Table 9: Laser diffraction results showing sizes ( $\mu\text{m}$ ) which 10% (Dv [10]), 50% (Dv [50]), and 90% (Dv [90]) of particles in the distribution are smaller than and the volume-weighted mean (D[4,3]).

IA or SOI	Station	Dv (10)	Dv (50)	Dv (90)	D[4,3]
IA 1*	1	156	431	1480	744
	2	267	436	760	501
IA 2	1	306	537	1070	788
	2	313	549	1050	748
IA 3*	1	240	425	774	624
	2	172	277	458	537
IA 4*	1	269	428	703	479
	2	264	472	1580	1160

IA or SOI	Station	Dv (10)	Dv (50)	Dv (90)	D[4,3]
SOI 1*	-	276	462	803	528
SOI 2*	-	223	339	519	362
SOI 3*	-	278	467	840	550
SOI 4	-	284	458	758	500
SOI 5	-	375	636	1040	695
SOI 6	-	325	544	935	608
SOI 7	-	277	571	4120	1380
SOI 8	-	185	2490	7270	2740

\* denotes benthic grab stations within the WTA

### 3.3 Benthic Community Analysis

#### 3.3.1 Taxonomic Composition

Fourteen of the sixteen benthic grab samples collected in this survey were analyzed for infauna and yielded a total of 1322 individual organisms from 8 unique phyla and 62 families (or LPTL; Table 10). The phyla Annelida, Arthropoda, and Mollusca dominated the samples in both abundance and unique number of taxa, representing 90% of all organisms and 91% of all unique taxa (Figure 16).

Table 10: Phyla present in the fourteen benthic grab samples.

Phyla	Abundant Taxonomic Groups (common names)	Abundance <sup>1</sup>	Number of Unique Taxa <sup>1</sup>
Annelida	Polychaete worms, oligochaete worms	535	46
Arthropoda	Amphipods, isopods, tanaids	555	26
Chordata	Lancelet	1	1
Cnidaria	Sea anemone	1	1
Echinodermata	Sand dollar	77	2
Mollusca	Bivalves, sea snails	106	18
Nematoda	Nematodes	40	1
Nemertea	Ribbon worms	7	4
Total		1322	99

<sup>1</sup> Reported as sum of all fourteen 0.05 m<sup>2</sup> grab samples




Figure 16: Proportional abundance and proportional number of unique taxa (species or LPTL) for each phylum collected in all benthic grab samples. Results presented as percentage of total.

Density within the benthic grab sites ranged from 18 organisms in IA 4 grab sample 2 to 280 organisms at SOI 2, with a mean density of 95 organisms per 0.05 m<sup>2</sup>, averaged across all samples (Table 11). Of the IAs, IA 1 and IA 4 contained the highest and lowest densities of infauna with 285 organisms and 49 organisms (total of both grabs), respectively.

Table 11: Density (# of individuals per 0.05 m<sup>2</sup>) of each Phylum present within each grab sample and the mean density for IA grab sample stations.

IA or SOI	Station	Annelida	Arthropoda	Chordata	Cnidaria	Echinodermata	Mollusca	Nematoda	Nemertea	Total Abundance
IA 1*	1	105	66	0	0	0	20	5	1	198
	2	19	55	0	0	4	5	2	0	87
	Average	62	61	0	0	2	13	4	1	141
IA 2	1	46	10	0	1	1	7	0	1	67
	2	65	22	0	0	0	11	6	0	106
	Average	56	16	0	1	1	9	3	1	85
IA 3*	1	39	9	0	0	0	9	1	0	59
	2	32	50	0	0	22	1	1	2	110
	Average	36	30	0	0	11	5	1	1	83
IA 4*	1	6	7	0	0	2	11	4	0	31
	2	7	8	0	0	0	1	0	0	18
	Average	7	8	0	0	1	6	2	0	23
SOI 1*	-	10	12	0	0	5	7	2	1	37

## ASOW BENTHIC ASSESSMENT REPORT

---

SOI 2*	-	59	209	1	0	5	3	2	1	280
SOI 3*	-	82	9	0	0	0	4	4	0	99
SOI 4	-	7	76	0	0	29	3	4	0	119
SOI 5	-	27	4	0	0	5	9	4	0	49
SOI 6	-	31	18	0	0	4	15	5	1	74

\* denotes benthic grab stations within the WTA



3.3.1.1 IA 1

Organisms collected in IA 1 (located in WTA) were classified into 6 phyla and 49 different taxa (LPTLs) (Table 12). The infaunal community recorded from the grabs in IA 1 were dominated by amphipods, polychaete worms, and tanaids. Species of direct economic importance identified in IA 1 included Atlantic rock crab (*Cancer irroratus*) and Atlantic surf clam (*Spisula solidissima*).

Table 12: Density of each phyla and taxa (LPTLs) collected at the two grab stations (data combined) in IA 1.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	<i>Polygordius jouinae</i>	13
	<i>Dipolydora socialis</i>	10
	<i>Marenzelleria viridis</i>	9
	<i>Goniadella gracilis</i>	8
	Cirratulidae	3
	Nephtyidae	3
	<i>Scoletoma fragilis</i>	3
	<i>Ampharete oculata</i>	2
	<i>Exogone hebes</i>	2
	<i>Neanthes acuminata</i> Complex	2
	<i>Protodorvillea kefersteini</i>	2
	<i>Aphroditella hastata</i>	1
	<i>Cabira incerta</i>	1
	<i>Lepidonotus sublevis</i>	1
	<i>Magelona papillicornis</i>	1
	<i>Magelona rosea</i>	1
	<i>Nephtys picta</i>	1
	<i>Oligochaeta</i>	1
	<i>Paradoneis lyra</i>	1
	<i>Phyllodoce mucosa</i>	1
<i>Prionospio</i> sp.	1	
<i>Sphaerodoropsis corrugata</i>	1	
<i>Sthenelais boa</i>	1	
<i>Tharyx</i> sp. A sensu MWRA 2007	1	
	<b>TOTAL</b>	<b>70</b>
Arthropoda	<i>Pseudunciola obliqua</i>	16
	<i>Unciola irrorata</i>	14
	<i>Ampelisca vadorum</i>	10
	<i>Ampelisca verrilli</i>	6
	<i>Tanaissus psammophilus</i>	5
	<i>Monocorophium acherusicum</i>	4

## ASOW BENTHIC ASSESSMENT REPORT

---

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
	Phoxocephalus holbolli	2
	Rhepoxynius hudsoni	2
	Cancer irroratus	1
	Chiridotea tuftsii	1
	Edotea triloba	1
	Eobrolgus spinosus	1
	Pagurus arcuatus	1
	<b>TOTAL</b>	<b>64</b>
Echinodermata	Echinarachnius parma	1
	Echinoidea	1
	<b>TOTAL</b>	<b>2</b>
Mollusca	Tritia trivittata	7
	Angulus versicolor	2
	Astyris lunata	1
	Bivalvia	1
	Cyclocardia borealis	1
	Periploma leanum	1
	Spisula solidissima	1
	Tellinidae	1
	<b>TOTAL</b>	<b>15</b>
Nematoda	Nematoda	4
Nemertea	Palaeonemertea	1
<b>TOTAL DENSITY OF ALL PHyla</b>		<b>156</b>

3.3.1.2 IA 2

Organisms collected in IA 2 were classified into 7 phyla and 28 different taxa (LPTLs) (Table 13). The infaunal community was dominated by polychaete worms, oligochaete worms, tanaisids, and bivalves. One species of direct economic importance, Atlantic surf clam (*Spisula solidissima*), was identified in IA 2.

Table 13: Density of each phyla and taxa (LPTL) collected at the two grab stations (data combined) in IA 2.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	<i>Polygordius jouinae</i>	33
	Oligochaeta	17
	<i>Dispia uncinata</i>	2
	Cirratulidae	1
	<i>Eulalia bilineata</i>	1
	<i>Goniadella gracilis</i>	1
	<i>Marenzelleria viridis</i>	1
	<i>Nephtys picta</i>	1
	<i>Sigalion arenicola</i>	1
	<i>Sphaerodoropsis corrugata</i>	1
<b>TOTAL</b>	<b>59</b>	
Arthropoda	<i>Tanaissus psammophilus</i>	12
	<i>Unciola irrorata</i>	1
	<i>Rhepoxynius hudsoni</i>	1
	<i>Pseudunciola obliqua</i>	1
	<i>Pseudoleptocuma minus</i>	1
	<i>Politolana polita</i>	1
	<i>Chiridotea arenicola</i>	1
	<i>Ampelisca verrilli</i>	1
<b>TOTAL</b>	<b>19</b>	
Cnidaria	<i>Edwardsia elegans</i>	1
Echinodermata	<i>Echinarachnius parma</i>	1
Mollusca	<i>Periploma leanum</i>	5
	<i>Astarte castanea</i>	2
	<i>Spisula solidissima</i>	2
	<i>Angulus tenellus</i>	1
	<i>Angulus versicolor</i>	1
	<i>Crenella sp.</i>	1
<b>TOTAL</b>	<b>14</b>	
Nematoda	Nematoda	3
Nemertea	<i>Tubulanus pellucidus</i>	1
<b>TOTAL DENSITY OF ALL PHyla</b>		<b>96</b>

**3.3.1.3 IA 3**

Organisms collected in IA 3 (located in WTA) were classified into 6 phyla and 44 different taxa (LPTLs) (Table 14). The infaunal community recorded from the grabs in IA 3 were dominated by ostracods, polychaete worms, and common sand dollars (*Echinarachnius parma*). A single species of direct economic importance, Atlantic surf clam (*Spisula solidissima*), was identified in IA 3.

Table 14: Density of each phyla and taxa (LPTL) collected at the two grab stations (data combined) in IA 3.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	Polygordius jouinae	13
	Ampharete oculata	3
	Aricidea (Aricidea) wassi	3
	Scoletoma fragilis	3
	Sthenelais sp.	3
	Onuphis eremita	2
	Ampharete acutifrons	1
	Caulleriella venefica	1
	Cirratulidae	1
	Cirrophorus furcatus	1
	Clymenella mucosa	1
	Dipolydora socialis	1
	Goniadella gracilis	1
	Harmothoe extenuata	1
	Magelona papillicornis	1
	Nephtyidae	1
	Nephtys picta	1
	Oligochaeta	1
	Pherusa affinis	1
	Phyllodoce mucosa	1
Sthenelais limicola	1	
<b>TOTAL</b>	<b>42</b>	
Arthropoda	Ostracoda	14
	Edotea triloba	4
	Ampelisca verrilli	2
	Pagurus arcuatus	2
	Pseudunciola obliquua	2
	Rhepoxynius hudsoni	2
	Unciola irrorata	2

## ASOW BENTHIC ASSESSMENT REPORT

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
	Americhelidium americanum	1
	Jassa falcata	1
	Oxyurostylis smithi	1
	Tanaissus psammophilus	1
	<b>TOTAL</b>	<b>32</b>
Echinodermata	Echinarachnius parma	9
	Echinoidea	3
	<b>TOTAL</b>	<b>12</b>
Mollusca	Angulus versicolor	2
	Crepidula fornicata	1
	Crepidula plana	1
	Pandora inornata	1
	Pleurobranchaea tarda	1
	Spisula solidissima	1
	Tritia trivittata	1
	<b>TOTAL</b>	<b>8</b>
Nematoda	Nematoda	1
Nemertea	Carinoma mutabilis	1
	Tubulanus pellucidus	1
	<b>TOTAL</b>	<b>2</b>
<b>TOTAL DENSITY OF ALL PHYLA</b>		<b>97</b>

### 3.3.1.4 IA 4

Organisms collected in IA 4 (located in WTA) were classified into 5 phyla and 19 different taxa (LPTLs) (Table 15). Overall density of infauna was relatively low compared to other IAs. The infaunal community recorded from the grabs in IA 4 was not dominated by any one species, but polychaete worms, tanaids, and a few species of bivalves were most abundant. Species of direct economic importance identified in IA 4 included Atlantic rock crab (*Cancer irroratus*) and Atlantic surf clam (*Spisula solidissima*).

Table 15: Density of each phyla and taxa (LPTL) collected at the two grab stations (data combined) in IA 4.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	Polygordius jouinae	3
	Nephtyidae	2
	Ampharete oculata	1
	Hemipodia simplex	1
	Sigalion arenicola	1
	<b>TOTAL</b>	<b>8</b>

## ASOW BENTHIC ASSESSMENT REPORT

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Arthropoda	Tanaissus psammophilus	4
	Pagurus sp.	2
	Bathyporeia quoddyensis	1
	Cancer irroratus	1
	Jassa falcata	1
	Parahaustorius attenuatus	1
	Unciola irrorata	1
	<b>TOTAL</b>	<b>11</b>
Echinodermata	Echinoidea	1
Mollusca	Angulus tenellus	3
	Spisula solidissima	2
	Astarte sp.	1
	Crenella sp.	1
	Tritia trivittata	1
	<b>TOTAL</b>	<b>9</b>
Nematoda	Nematoda	2
<b>TOTAL DENSITY OF ALL PHYLA</b>		<b>30</b>

### 3.3.1.5 SOI 1

Organisms collected in SOI 1 (located in WTA) were classified into 6 phyla and 17 different taxa (LPTLs) (Table 16).

Table 16: Density of each phyla and taxa (LPTL) collected at the grab station at SOI 1.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	Exogone hebes	3
	Polygordius jouinae	3
	Eulalia bilineata	1
	Leitoscoloplos robustus	1
	Magelona papillicornis	1
	Onuphis eremita	1
	<b>TOTAL</b>	<b>10</b>
Arthropoda	Tanaissus psammophilus	7
	Rhepoxynius hudsoni	3
	Protohaustorius deichmannae	1
	Unciola irrorata	1
	<b>TOTAL</b>	<b>12</b>
Echinodermata	Echinoidea	3

## ASOW BENTHIC ASSESSMENT REPORT

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
	Echinarachnius parma	2
	<b>TOTAL</b>	<b>5</b>
Mollusca	Angulus tenellus	3
	Astarte castanea	2
	Periploma leanum	2
	<b>TOTAL</b>	<b>7</b>
Nematoda	Nematoda	2
Nemertea	Carinoma mutabilis	1
<b>TOTAL DENSITY OF ALL PHYLA</b>		<b>37</b>

### 3.3.1.6 SOI 2

Organisms collected in SOI 2 (located in WTA) were classified into 7 phyla and 30 different taxa (LPTLs) (Table 17).

Table 17: Density of each phyla and taxa (LPTL) collected at the grab station at SOI 2.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	Magelona papillicornis	22
	Cirrophorus furcatus	5
	Polygordius jouinae	5
	Caulleriella venefica	3
	Cirratulidae	3
	Exogone hebes	3
	Nephtyidae	3
	Oligochaeta	3
	Aricidea (Aricidea) wassi	2
	Aricidea sp.	2
	Tharyx sp. A sensu MWRA 2007	2
	Nephtys picta	1
	Onuphis eremita	1
	Sigalion arenicola	1
	Spiochaetopterus oculatus	1
	Spiophanes bombyx Complex	1
	Streptosyllis arenae	1
	<b>TOTAL</b>	<b>59</b>
Arthropoda	Pseudunciola obliquua	157
	Ampelisca verrilli	32
	Tanaissus psammophilus	15

## ASOW BENTHIC ASSESSMENT REPORT

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
	Ostracoda	3
	Rhepoxynius hudsoni	2
	<b>TOTAL</b>	<b>209</b>
Chordata	Branchiostoma virginiae	1
Echinodermata	Echinarachnius parma	4
	Echinoidea	1
	<b>TOTAL</b>	<b>6</b>
Mollusca	Nucula proxima	1
	Tellinidae	1
	Tritia trivittata	1
	<b>TOTAL</b>	<b>3</b>
Nematoda	Nematoda	2
Nemertea	Lineidae	1
<b>DENSITY OF ALL PHYLA</b>		<b>280</b>

### 3.3.1.7 SOI 3

Organisms collected in SOI 3 (located in WTA) were classified into 4 phyla and 13 different taxa (LPTLs) (Table 18).

Table 18: Density of each phyla and taxa (LPTL) collected at the grab station at SOI 3.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	Polygordius jouinae	65
	Oligochaeta	6
	Hemipodia simplex	4
	Parapionosyllis longicirrata	3
	Exogone hebes	1
	Lumbrinerides acuta	1
	Sigalion arenicola	1
	Streptosyllis arenae	1
	<b>TOTAL</b>	<b>82</b>
Arthropoda	Rhepoxynius hudsoni	4
	Pseudunciola obliqua	3
	Tanaissus psammophilus	2
	<b>TOTAL</b>	<b>9</b>
Mollusca	Angulus tenellus	4
Nematoda	Nematoda	4
<b>DENSITY OF ALL PHYLA</b>		<b>99</b>



3.3.1.8 SOI 4

Organisms collected in SOI 4 were classified into 5 phyla and 19 different taxa (LPTLs) (Table 19).

Table 19: Density of each phyla and taxa (LPTL) collected at the grab station at SOI 4.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	Signalion arenicola	2
	Aricidea (Aricidea) wassi	1
	Caulleriella venefica	1
	Magelona papillicornis	1
	Nephtyidae	1
	Oligochaeta	1
	<b>TOTAL</b>	<b>7</b>
Arthropoda	Pseudunciola obliqua	35
	Tanaissus psammophilus	32
	Rhepoxynius hudsoni	4
	Americhelidium americanum	1
	Bathyporeia quoddyensis	1
	Oxyurostylis smithi	1
	Protohaustorius deichmannae	1
	Ptilanthura tenuis	1
<b>TOTAL</b>	<b>76</b>	
Echinodermata Echinoidea		26
	Echinarachnius parma	3
<b>TOTAL</b>	<b>29</b>	
Mollusca	Tritia trivittata	2
	Periploma sp.	1
<b>TOTAL</b>	<b>3</b>	
Nematoda	Nematoda	4
<b>DENSITY OF ALL PHYLA</b>		<b>119</b>

3.3.1.9 SOI 5

Organisms collected in SOI 5 were classified into 5 phyla and 18 different taxa (LPTLs) (Table 20).

Table 20: Density of each phyla and taxa (LPTL) collected at the grab station at SOI 5.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	Polygordius jouinae	10
	Sigalion arenicola	7
	Travisia forbesii	3
	Dipolydora socialis	2
	Dispio uncinata	1
	Eulalia bilineata	1
	Hemipodia simplex	1
	Nephtys picta	1
	Sphaerodoropsis corrugata	1
	<b>TOTAL</b>	<b>27</b>
Arthropoda	Bathyporeia quoddyensis	2
	Oxyurostylis smithi	1
	Protohaustorius wigleyi	1
<b>TOTAL</b>	<b>4</b>	
Echinodermata	Echinoidea	3
	Echinarachnius parma	2
<b>TOTAL</b>	<b>5</b>	
Mollusca	Periploma sp.	4
	Astarte castanea	3
	Spisula solidissima	2
<b>TOTAL</b>	<b>9</b>	
Nematoda	Nematoda	4
<b>DENSITY OF ALL PHYLA</b>		<b>49</b>

### 3.3.1.10 SOI 6

Organisms collected in SOI 6 were classified into 6 phyla and 19 different taxa (LPTLs) (Table 21).

Table 21: Density of each phyla and taxa (LPTL) collected at the grab station at SOI 6.

Phyla	Taxa (LPTL)	Density (# / 0.05 m <sup>2</sup> )
Annelida	Hemipodia simplex	9
	Goniadella gracilis	8
	Polygordius jouinae	5
	Dipolydora socialis	2
	Nephtys picta	2
	Cauleriella venefica	1
	Dispio uncinata	1
	Lumbrinerides acuta	1
	Sigalion arenicola	1
	Spiophanes bombyx Complex	1
	TOTAL	31
Arthropoda	Tanaissus psammophilus	16
	Americhelidium americanum	1
	Rhepoxynius hudsoni	1
	TOTAL	18
Echinodermata	Echinoidea	4
Mollusca	Periploma sp.	9
	Angulus tenellus	4
	Spisula solidissima	2
	TOTAL	19
Nematoda	Nematoda	5
Nemertea	Tubulanus pellucidus	1
<b>DENSITY OF ALL PHYLA</b>		<b>74</b>

### 3.3.2 Richness, Diversity, and Evenness

Taxonomic richness ranged from 2.18 at SOI 3 to 5.77 at station 2 in IA 3 with a mean richness of 3.96 overall (Table 22). Diversity was consistent between grab stations ranging from 1.39 at SOI 3 to 2.83 at station 2 of IA 3. Evenness ranged from 0.55 at station 1 in IA 2 to 0.94 at station 1 in IA 4. While considering both grabs from each IA, IA 3 had the highest infaunal richness, IA 1 had the most infaunal diversity, and IA 4 had the most even infaunal species composition according to the ecological indices (Figure 17).

## ASOW BENTHIC ASSESSMENT REPORT

Table 22: Community composition parameters calculated for each grab sample station and for each IA. Combined values for individual stations within the four IAs were either averaged ( $\bar{x}$ ) or summed ( $\Sigma$ ) depending on parameter.

IA	Station	Density (# of individuals per 0.05 m <sup>2</sup> )	# of Unique Taxa	Ecological Indices		
				Richness	Diversity	Evenness
IA 1*	1	197	32	5.30	2.70	0.80
	2	85	22	4.05	2.30	0.78
	Combined	$\bar{x} = 141$	$\Sigma = 49$	$\bar{x} = 4.68$	$\bar{x} = 2.50$	$\bar{x} = 0.79$
IA 2	1	66	16	3.10	1.45	0.55
	2	104	17	3.45	2.03	0.72
	Combined	$\bar{x} = 85$	$\Sigma = 28$	$\bar{x} = 3.28$	$\bar{x} = 1.74$	$\bar{x} = 0.63$
IA 3*	1	58	19	4.19	2.08	0.72
	2	108	31	5.77	2.83	0.75
	Combined	$\bar{x} = 83$	$\Sigma = 44$	$\bar{x} = 4.98$	$\bar{x} = 2.45$	$\bar{x} = 0.78$
IA 4*	1	30	16	4.41	2.61	0.94
	2	16	9	2.89	1.92	0.88
	Combined	$\bar{x} = 23$	$\Sigma = 19$	$\bar{x} = 3.65$	$\bar{x} = 2.27$	$\bar{x} = 0.91$
SOI 1*	-	37	17	4.43	2.65	0.93
SOI 2*	-	280	30	4.08	1.76	0.56
SOI 3*	-	99	13	2.18	1.39	0.58
SOI 4	-	119	19	3.77	1.95	0.66
SOI 5	-	49	18	4.11	2.56	0.90
SOI 6	-	74	19	3.72	2.47	0.87

\* denotes benthic grab stations within the WTA




Figure 17: Richness (top) Diversity (middle) and Evenness (bottom) index values for each IA and SOI. Stations located in the WTA are denoted with an asterisk (\*).

## 4 CMECS CLASSIFICATIONS

We applied NMFS-modified CMECS (2021) substrate component classifications to each grab site using a combination of the visual observations of the grab samples after they were brought aboard (Table 23) and the grain size composition from the sediment analysis (Section 3.2). Results are presented as a hierarchy in Table 24. Only one sample was classified in the Gravel Mixes substrate group ( $\geq 30\%$  gravel): SOI 8 was classified as sandy gravel with 58% gravel and 41% sand composition. Samples IA 1: S1 (located in WTA), IA 2: S1, IA 4: S2 (located in WTA), and SOI 7 were classified as gravelly sand ( $\geq 5\%$  gravel) with either trace or sparse clam hash, and some worm hash also in IA 1: S1. These four gravelly sand and

## ASOW BENTHIC ASSESSMENT REPORT

one gravel mix sample are considered complex habitat due to  $\geq 5\%$  gravel content, with gravel defined as particles  $\geq 2$  mm.

Three samples were classified as very coarse/coarse sand, with IA 2: S2 also containing sparse clam hash while SOI 5 and SOI 6 contained trace clam hash. The remaining eight samples were classified as medium sand (Figure 18). Thus over 68% of all samples (11 of 16) and 78% of WTA samples (7 of 9) collected in the Lease Area were classified as soft bottom habitat ( $< 5\%$  gravel cover). No large boulders or sensitive emergent epifauna were sampled.



















Figure 18: Substrate CMECS classifications by sample site. Note that “B” delineates buoy IAs (e.g., B3-2 is IA site 3 station 2) and “S” delineates SOIs.

**ASOW BENTHIC ASSESSMENT REPORT**

Table 23: Images of grab samples prior to processing, along with CMECS classifications (Substrate & Biological modifier).

IA or SOI	Station 1	Station 2
IA 1*	 <p data-bbox="509 562 907 585">Gravelly Sand &amp; Sparse Worm and Clam Hash</p>	 <p data-bbox="1062 562 1190 585">Medium Sand</p>
IA 2	 <p data-bbox="565 858 850 884">Gravelly Sand &amp; Trace Clam Hash</p>	 <p data-bbox="927 858 1325 884">Very Coarse/Coarse Sand &amp; Sparse Clam Hash</p>
IA 3*	 <p data-bbox="565 1157 850 1182">Medium Sand &amp; Trace Clam Hash</p>	 <p data-bbox="984 1157 1269 1182">Medium Sand &amp; Trace Clam Hash</p>
IA 4*	 <p data-bbox="558 1455 862 1480">Medium Sand &amp; Sparse Clam Hash</p>	 <p data-bbox="979 1455 1276 1480">Gravelly Sand &amp; Sparse Clam Hash</p>
SOI 1* / SOI 2*	 <p data-bbox="558 1753 862 1780">Medium Sand &amp; Sparse Clam Hash</p>	 <p data-bbox="984 1753 1269 1780">Medium Sand &amp; Trace Clam Hash</p>

**ASOW BENTHIC ASSESSMENT REPORT**


IA or SOI	Station 1	Station 2
SOI 3* / SOI 4	 <p data-bbox="560 499 857 525">Medium Sand &amp; Sparse Clam Hash</p>	 <p data-bbox="987 499 1284 525">Medium Sand &amp; Trace Clam Hash</p>
SOI 5 / SOI 6	 <p data-bbox="516 798 901 823">Very Coarse/Coarse Sand &amp; Trace Clam Hash</p>	 <p data-bbox="933 798 1318 823">Very Coarse/Coarse Sand &amp; Trace Clam Hash</p>
SOI 7 / SOI 8	 <p data-bbox="560 1096 857 1121">Gravelly Sand &amp; Sparse Clam Hash</p>	 <p data-bbox="987 1096 1284 1121">Sandy Gravel &amp; Sparse Clam Hash</p>

\* denotes benthic grab stations within the WTA



**ASOW BENTHIC ASSESSMENT REPORT**

Table 24: CMECS hierarchical classification of substrates collected at each station (S) in each IA. Stations located in the WTA are denoted with an asterisk (\*).



## 5 REFERENCES

- ASTM D422-63 (2007) e2, Standard Test Method for Particle-Size Analysis of Soils (Withdrawn 2016). ASTM International. West Conshohocken, PA, 2007. [www.astm.org](http://www.astm.org).
- Auster, P.J. 1998. A conceptual model of the impacts of fishing gear on the integrity of fish habitats. *Conservation Biology*. 12(6):1198-1203.
- Bousfield, E.L. 1973. *Shallow-Water Gammaridean Amphipoda of New England*. Cornell University Press, xii + 312pp.
- Cutler, E.B. 1994. *The Sipuncula: Their Systematics, Biology, and Evolution*. Cornell University Press, xvii + 453pp.
- Bureau of Ocean Energy Management (BOEM). 2019. Guidelines for Providing Benthic Habitat Survey Information for Renewable Energy Development on the Atlantic Outer Continental Shelf Pursuant to 30 CFR Part 585. June 2019. BOEM Office of Renewable Energy Programs, US Department of the Interior. 9pp.
- Federal Geographic Data Committee (FGDC). 2012. Coastal and Marine Ecological Classification Standard, June 2012. FGDC-STD-018-2012. 353pp.
- National Marine Fisheries Service (NMFS). 2021. Recommendations for Mapping Fish Habitat. NMFS GARFO Habitat Conservation and Ecosystem Services Division. March 2021. 22 pp.
- U.S. Environmental Protection Agency (EPA). 1986. Method 9060 Total Organic Carbon. SW-846: Hazardous Waste Test Methods, Ch 5.
- Winston, J.E. and P.J Hayward. 2012. *The Marine Bryozoans of the Northeast Coast of the United States: Maine to Virginia*. Virginia Museum of Natural History Memoir 11, xii + 180pp.

## APPENDIX A FIELD DATASHEETS

ASOW BENTHIC ASSESSMENT REPORT

video/grab datasheet

date: 10/12/2019		cruise: Atlantic Shores		samplers: JZ	
station: RL653	lat: 39 21.5007	start time: 15:15:00	file name:		
depth: 21M	long: 74 11.2525	end time: 15:19:30	O <sub>2</sub> : 7.85 mg/L		
YSI depth: 8.7ft	YSI temp: 18.9	YSI pH: 8.9			
rock, <u>unconsolidated mineral</u> , algal, coral, organic, ooze, shell, worm		flat sand and mud, sand waves, biogenic textures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud		pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: Sandy, few small sand fragments					
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N					
Pen. depth: 95mm		TOC/bio/sed ID# RL653		bio volume (inches high) 1.75	
Sediment characterization: Sandy					
station: MBA1651	lat: 39 18.603	start time:	file name:		
depth: 28.5m	long: 74 6.5923	end time: 17:25	7.87 mg/L		
YSI depth: 8.01ft	YSI temp: 18.93	YSI pH: 8.91			
rock, <u>unconsolidated mineral</u> , algal, coral, organic, ooze, shell, worm		flat sand and mud, sand waves, biogenic textures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud		pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: GPS Feed froze for a while					
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N					
Pen. depth: 72mm		TOC/bio/sed ID# MBA1651		bio volume (inches high) 1.75	
Sediment characterization:					
station: MBA1652	lat: 39 18.5498	start time: 17:47:02	file name:		
depth: 28.7	long: 74 06.4631	end time: 17:48:10	7.87 mg/L		
YSI depth: 12.13	YSI temp: 18.992	YSI pH: 8.91			
rock, <u>unconsolidated mineral</u> , algal, coral, organic, ooze, shell, worm		flat sand and mud, sand waves, biogenic textures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud		pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments:					
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N					
Pen. depth: 82mm		TOC/bio/sed ID#		bio volume (inches high) 1.75	
Sediment characterization:					
station: MBA3651	lat: 39 17.3562	start time: 19:23	file name:		
depth: 34.7	long: 73 58.5943	end time: 19:19:40	7.89 mg/L		
YSI depth: 12.87	YSI temp: 18.74	YSI pH: 8.95			
rock, <u>unconsolidated mineral</u> , algal, coral, organic, ooze, shell, worm		flat sand and mud, sand waves, biogenic textures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud		pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: some shell hash, 4 scallop shell					
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N					
Pen. depth: 139		TOC/bio/sed ID#		bio volume (inches high) 3	
Sediment characterization:					

ASOW BENTHIC ASSESSMENT REPORT

video/grab datasheet

2

date: 40 / 12 / 2019		cruise: Atlantic Shores		samplers: J2, Felipe	
station: MBA3658	lat:	start time: 19:46	file name:		
depth: 33.5	long:	end time: 19:49			
YSI depth: 12.47ft	YSI temp: 18.643	YSI pH: 8.94	7.85mg/L		
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm		flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud		pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: didn't close					
<b>Grab</b> Failed <input checked="" type="radio"/> Y / <input type="radio"/> N   >50% Full <input type="radio"/> Y / <input type="radio"/> N   <100% Full <input type="radio"/> Y / <input type="radio"/> N   Undisturbed Surface <input type="radio"/> Y / <input type="radio"/> N					
Pen. depth:		TOC/bio/sed ID#		bio volume (inches high)	
Sediment characterization:					

station: MBA3658		lat: 39 17.2256		start time: 20:05.31		file name:	
depth: 33.5		long: 73 58.6107		end time: 20:06			
YSI depth: 12.47ft		YSI temp: 18.643		YSI pH: 8.94		7.85mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: sand dollars + Longfin Inshore Squid (photo + release)							
<b>Grab</b> Failed <input type="radio"/> Y / <input checked="" type="radio"/> N   >50% Full <input type="radio"/> Y / <input type="radio"/> N   <100% Full <input type="radio"/> Y / <input type="radio"/> N   Undisturbed Surface <input type="radio"/> Y / <input checked="" type="radio"/> N							
Pen. depth: 76		TOC/bio/sed ID#		bio volume (inches high) 1.75			
Sediment characterization:							

station: RL651		lat: 39 14.1066		start time: 21:13		file name:	
depth: 30		long: 73 59.6160		end time: 21:14.26			
YSI depth: 14.06		YSI temp: 18.748		YSI pH: 8.95		7.96mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: Failed 1st attempt, reloaded							
<b>Grab</b> Failed <input type="radio"/> Y / <input checked="" type="radio"/> N   >50% Full <input type="radio"/> Y / <input type="radio"/> N   <100% Full <input type="radio"/> Y / <input type="radio"/> N   Undisturbed Surface <input type="radio"/> Y / <input type="radio"/> N							
Pen. depth: 73		TOC/bio/sed ID#		bio volume (inches high) 2.5"			
Sediment characterization:							

station: MBA4651		lat: 39 12.1396		start time: 22:05		file name:	
depth: 26		long: 74 5.0065		end time: 22:35			
YSI depth: 11.6ft		YSI temp: 18.703		YSI pH: 8.94		8.03mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: some shell hash, some sand dollars							
<b>Grab</b> Failed <input type="radio"/> Y / <input checked="" type="radio"/> N   >50% Full <input type="radio"/> Y / <input type="radio"/> N   <100% Full <input type="radio"/> Y / <input type="radio"/> N   Undisturbed Surface <input type="radio"/> Y / <input type="radio"/> N							
Pen. depth: 90		TOC/bio/sed ID#		bio volume (inches high) 2			
Sediment characterization:							

ASOW BENTHIC ASSESSMENT REPORT

video/grab datasheet

3

date: 10 / 12 / 2019		cruise: Atlantic Shores		samplers: JZ / Edgar	
station: MBA 4652	lat: 39 12.1881	start time: 23:01	file name:		
depth: 25.5	long: 74 4.7806	end time: 23:03			
YSI depth: 14.6	YSI temp: 18.09	YSI pH: 8.96	8.02 %		
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm		flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud		pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: Some shell hash					
<b>Grab</b> Failed Y / <input checked="" type="radio"/> N   >50% Full <input checked="" type="radio"/> N   <100% Full <input checked="" type="radio"/> N   Undisturbed Surface <input checked="" type="radio"/> N					
Pen. depth: 85		TOC/bio/sed ID#		bio volume (inches high) 2"	
Sediment characterization:					

station: RL652		lat: 39 15.4624		start time: 00:38		file name:	
depth: 26		long: 74 12.3109		end time: 00:40			
YSI depth: 9.3 ft		YSI temp: 18.551		YSI pH: 8.91		7.91 mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: Sand dollars							
<b>Grab</b> Failed Y / <input checked="" type="radio"/> N   >50% Full <input checked="" type="radio"/> N   <100% Full <input checked="" type="radio"/> N   Undisturbed Surface <input checked="" type="radio"/> N							
Pen. depth: 91		TOC/bio/sed ID#		bio volume (inches high) 2.15			
Sediment characterization:							

station:		lat:		start time:		file name:	
depth:		long:		end time:			
YSI depth:		YSI temp:		YSI pH:			
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments:							
<b>Grab</b> Failed Y / N   >50% Full Y / N   <100% Full Y / N   Undisturbed Surface Y / N							
Pen. depth:		TOC/bio/sed ID#		bio volume (inches high)			
Sediment characterization:							

station:		lat:		start time:		file name:	
depth:		long:		end time:			
YSI depth:		YSI temp:		YSI pH:			
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments:							
<b>Grab</b> Failed Y / N   >50% Full Y / N   <100% Full Y / N   Undisturbed Surface Y / N							
Pen. depth:		TOC/bio/sed ID#		bio volume (inches high)			
Sediment characterization:							

ASOW BENTHIC ASSESSMENT REPORT

video/grab datasheet

4

date: 10/13/2019		cruise: Atlantic Shores		samplers: JZ, Felipe	
station: MBA2651	lat: 39.33.9308	start time: 10:08	file name:		
depth: 25	long: 73.58.1656	end time: 10:06			
YSI depth: 8.91	YSI temp: 18.462	YSI pH: 8.78	7.86 mg/L		
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm			flat sand and mud, sand waves, biogenic xtures, shell aggreg		
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud			pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs		
comments: Some gravel, 1 sand dollar					
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N					
Pen. depth: 51			TOC/bio/sed ID# bio volume (inches high)		
Sediment characterization:					

station: MBA2652		lat: 39.33.9907		start time:		file name:	
depth: 25		long: 73.58.1989		end time: 10:35			
YSI depth: 5.46		YSI temp: 18.487		YSI pH: 8.90		7.87 mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: Some shells, sandy							
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N							
Pen. depth: 91				TOC/bio/sed ID# bio volume (inches high) 3.5			
Sediment characterization:							

station: RL656		lat: 39.24.021		start time:		file name:	
depth: 28		long: 73.57.6006		end time: 12:39			
YSI depth: 11.45		YSI temp: 18.706		YSI pH: 8.97		7.88 mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: Ocean quahog							
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N							
Pen. depth: 92				TOC/bio/sed ID# bio volume (inches high) 3			
Sediment characterization:							

station: RL659		lat: 39.23.791		start time:		file name:	
depth: 26		long: 74.2.7639		end time: 13:43			
YSI depth: 8.11 ft		YSI temp: 18.825		YSI pH: 9.04		7.69 mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments: Urchin, some shell hash some clay deposits/clumps?							
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N							
Pen. depth: 95				TOC/bio/sed ID# bio volume (inches high) 2.5			
Sediment characterization:							

ASOW BENTHIC ASSESSMENT REPORT

video/grab datasheet

5

date: 0/13/2019		cruise: Atlantic Shores		samplers:	
station: RL655	lat: 39 26.7896	start time:	file name:		
depth: 21	long: 74 5.6972	end time: 14:43			
YSI depth: 14.32 ft	YSI temp: 18.645	YSI pH: 8.92		2.90 mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm			flat sand and mud, sand waves, biogenic xtures, shell aggreg		
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud			pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs		
comments:					
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N Pen. depth: 115 mm TOC/bio/sed ID# bio volume (inches high) 2.5 Sediment characterization:					

station: RL657		lat: 39 33.8481		start time: 16:46		file name:	
depth: 24		long: 74 2.1521		end time: 16:48			
YSI depth: 8.97		YSI temp: 18.697		YSI pH: 8.90		2.79 mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments:							
failed dx <b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N Pen. depth: 118 TOC/bio/sed ID# bio volume (inches high) Sediment characterization: No Bio Sample							

station: RL658		lat: 39 38.0361		start time:		file name:	
depth: 26		long: 73 57.7192		end time: 18:07			
YSI depth: 5.2		YSI temp: 18.454		YSI pH: 8.87		8.15 mg/L	
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments:							
small worm tubes some gravel <b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N Pen. depth: 85 TOC/bio/sed ID# bio volume (inches high) Sediment characterization: No Bio Sample							

station:		lat:		start time:		file name:	
depth:		long:		end time:			
YSI depth:		YSI temp:		YSI pH:			
rock, <b>unconsolidated mineral</b> , algal, coral, organic, ooze, shell, worm				flat sand and mud, sand waves, biogenic xtures, shell aggreg			
gravel, gravel mix, gravelly, slightly gravelly, sand, sandy mud, mud				pebble-cobble, p-c w sponge, buried/disp boulders, piled bldrs			
comments:							
<b>Grab</b> Failed Y/N   >50% Full Y/N   <100% Full Y/N   Undisturbed Surface Y/N Pen. depth: TOC/bio/sed ID# bio volume (inches high) Sediment characterization:							



## APPENDIX B PHOTOMICROSCOPY IMAGES



Figure B- 1. Photomicrograph (1 of 2) of sediment sample from SOI 1 taken at 40x magnification.



Figure B- 2. Photomicrograph (2 of 2) of sediment sample from SOI 1 taken at 40x magnification.




Figure B- 3. Photo of 1180 µm fraction of sediment sample from SOI 1.




Figure B- 4. Photo of 2000 µm fraction of sediment sample from SOI 1.



Figure B- 5. Photomicrograph (1 of 2) of sediment sample from SOI 2 taken at 40x magnification.



Figure B- 6. Photomicrograph (2 of 2) of sediment sample from SOI 2 taken at 40x magnification.




Figure B- 7. Photo of 1180 µm fraction of sediment sample from SOI 2.




Figure B- 8. Photo of 2000 µm fraction of sediment sample from SOI 2.




Figure B- 9. Photomicrograph (1 of 2) of sediment sample from SOI 3 taken at 40x magnification



Figure B- 10. Photomicrograph (2 of 2) of sediment sample from SOI 3 taken at 40x magnification.




Figure B- 11. Photo of 1180  $\mu\text{m}$  fraction of sediment sample from SOI 3.




Figure B- 12. Photo of 2000  $\mu\text{m}$  fraction of sediment sample from SOI 3.




Figure B- 13. Photomicrograph (1 of 2) of sediment sample from SOI 4 taken at 40x magnification.



Figure B- 14. Photomicrograph (2 of 2) of sediment sample from SOI 4 taken at 40x magnification.






Figure B- 15. Photo of 1180 µm fraction of sediment sample from SOI 4.




Figure B- 16. Photo of 2000 µm fraction of sediment sample from SOI 4.




Figure B- 17. Photomicrograph (1 of 2) of sediment sample from SOI 5 taken at 40x magnification.



Figure B- 18. Photomicrograph (2 of 2) of sediment sample from SOI 5 taken at 40x magnification.




Figure B- 19. Photo of 1180 µm fraction of sediment sample from SOI 5.




Figure B- 20. Photo of 2000 µm fraction of sediment sample from SOI 5.



Figure B- 21. Photomicrograph (1 of 2) of sediment sample from SOI 6 taken at 40x magnification.




Figure B- 22. Photomicrograph (2 of 2) of sediment sample from SOI 6 taken at 40x magnification.




Figure B- 23. Photo of 1180  $\mu\text{m}$  fraction of sediment sample from SOI 6.




Figure B- 24. Photo of 2000  $\mu\text{m}$  fraction of sediment sample from SOI 6.




Figure B- 25. Photomicrograph (1 of 2) of sediment sample from SOI 7 taken at 40x magnification.




Figure B- 26. Photomicrograph (2 of 2) of sediment sample from SOI 7 taken at 40x magnification.



Figure B- 27. Photo of 1180  $\mu\text{m}$  fraction of sediment sample from SOI 7.




Figure B- 28. Photo of 2000  $\mu\text{m}$  fraction of sediment sample from SOI 7.




Figure B- 29. Photomicrograph (1 of 2) of sediment sample from SOI 8 taken at 40x magnification.




Figure B- 30. Photomicrograph (2 of 2) of sediment sample from SOI 8 taken at 100x magnification.






Figure B- 31. Photo of 1180 µm fraction of sediment sample from SOI 8.



Figure B- 32. Photo of 2000 µm fraction of sediment sample from SOI 8.




Figure B- 33. Photomicrograph (1 of 2) of sediment sample from grab 1 of 2 from IA 1 taken at 40x magnification.



Figure B- 34. Photomicrograph (2 of 2) of sediment sample from grab 1 of 2 from IA 1 taken at 100x magnification.



Figure B- 35. Photo of 1180 µm fraction of sediment grab 1 of 2 from IA 1.




Figure B- 36. Photo of 2000 µm fraction of sediment grab 1 of 2 from IA 1.




Figure B- 37. Photomicrograph (1 of 2) of sediment grab 2 of 2 from IA 1 taken at 40X magnification.




Figure B- 38. Photomicrograph (2 of 2) of sediment grab 2 of 2 from IA 1 taken at 100X magnification.




Figure B- 39. Photo of 1180 µm fraction of sediment grab 2 of 2 from IA 1.




Figure B- 40. Photo of 2000 µm fraction of sediment grab 2 of 2 from IA 1.




Figure B- 41. Photomicrograph (1 of 2) of sediment grab 1 of 2 from IA 2 taken at 40x magnification.




Figure B- 42. Photomicrograph (2 of 2) of sediment grab 1 of 2 from IA 2 taken at 40x magnification.




Figure B- 43. Photo of 1180  $\mu\text{m}$  fraction of sediment grab 1 of 2 at IA 2.



Figure B- 44. Photo of 2000  $\mu\text{m}$  fraction of sediment grab 1 of 2 at IA 2.




Figure B- 45. Photomicrograph (1 of 2) of sediment from grab 2 of 2 at IA 2 taken at 40x magnification.




Figure B- 46. Photomicrograph (2 of 2) of sediment from grab 2 of 2 at IA 2 taken at 40x magnification.






Figure B- 47. Photo of 1180 µm fraction of sediment from grab 2 of 2 at IA 2.



Figure B- 48. Photo of 2000 µm fraction of sediment from grab 2 of 2 at IA 2.




Figure B- 49. Photomicrograph (1 of 2) of sediment from grab 1 of 2 at IA 3 taken at 40x magnification




Figure B- 50. Photomicrograph (2 of 2) of sediment grab 1 of 2 from IA 3 taken at 100x magnification




Figure B- 51. Photo of 1180  $\mu\text{m}$  fraction of sediment grab 1 of 2 from IA 3.




Figure B- 52. Photo of 2000  $\mu\text{m}$  fraction of sediment grab 1 of 2 at IA 3.




Figure B- 53. Photomicrograph (1 of 2) of sediment grab 2 of 2 from IA 3 taken at 40X magnification.




Figure B- 54. Photomicrograph (2 of 2) of sediment grab 2 of 2 from IA 3 taken at 100X magnification.




Figure B- 55. Photo of 1180  $\mu\text{m}$  fraction of sediment grab 2 of 2 at IA 3.



Figure B- 56. Photo of 2000  $\mu\text{m}$  fraction of sediment grab 2 of 2 at IA 3.




Figure B- 57. Photomicrograph (1 of 2) of sediment sample from grab 1 of 2 from IA 4 taken at 40x magnification.




Figure B- 58. Photomicrograph (2 of 2) of sediment sample from grab 1 of 2 from IA 4 taken at 40x magnification.




Figure B- 59. Photo of 1180 µm fraction of sediment grab 1 of 2 at IA 4.




Figure B- 60. Photo of 2000 µm fraction of sediment grab 1 of 2 at IA 4.




Figure B- 61. Photomicrograph (1 of 2) of sediment sample from grab 2 of 2 from IA 4 taken at 40x magnification.




Figure B- 62. Photomicrograph (2 of 2) of sediment sample from grab 2 of 2 from IA 4 taken at 40x magnification.





Figure B- 63. Photo of 1180 µm fraction of sediment grab 2 of 2 at IA 4.



Figure B- 64. Photo of 2000 µm fraction of sediment grab 2 of 2 at IA 4.