

# GLIDER SURVEILLANCE FOR NEAR-REAL-TIME DETECTION AND SPATIAL MANAGEMENT OF NORTH ATLANTIC RIGHT WHALES

By Katherine L. Indeck, Mark F. Baumgartner, Laurence Lecavalier, Frederick Whoriskey, Delphine Durette-Morin, Neal R. Pettigrew, Jacqueline M. McSweeney, Lesley H. Thorne, Katherine L. Gallagher, Catherine R. Edwards, Erin Meyer-Gutbrod, and Kimberley T.A. Davies

## ABSTRACT

Successful area-based ocean management relies on long-term, persistent biological monitoring using reliable ocean observation assets. Underwater electric gliders fill a unique monitoring niche compared to other platforms because they can autonomously survey across diverse environments—from shallow coastal waters to remote offshore areas—for weeks to months at a time. Gliders equipped with passive acoustic monitoring (PAM) devices are capable of robust, continuous near-real-time monitoring of numerous species of whales. Here, we highlight five case studies to discuss how gliders are being used for area-based monitoring of the internationally migratory and critically endangered North Atlantic right whale to address several different spatial management objectives. Examples include dynamic management of shipping zones and fishery-area closures in Canadian waters, glider-based monitoring in the United States to mitigate vessel strikes and fishing gear entanglements, surveys to assess whale habitat use near offshore wind energy development areas in the northeastern United States, and surveillance of the coastal calving grounds in the southeastern United States. These examples illustrate how PAM-equipped gliders are being used to monitor an endangered cetacean species with complex conservation management needs across its range. These assets are supporting risk reduction measures across diverse regions, and their use is likely to continue to expand in support of species conservation and threat mitigation.

## AUTONOMOUS ACOUSTIC GLIDERS FOR AREA-BASED MANAGEMENT

Area-based ocean management aims to balance human use of the marine environment with biological conservation (Maxwell et al., 2015). There are two primary management frameworks for achieving this: (1) static management areas (e.g., conventional marine protected areas) that are fixed in time (e.g., seasonally) and space based on historical data regarding the occurrence of species needing protection, and (2) dynamic management areas that are triggered in response to recent observations or predictions of species occurrence. Static management is typically applied to known critical habitats or where predictable aggregations of at-risk species frequently overlap with high-threat human

activities (i.e., *high risk areas*). Alternatively, dynamic management is increasingly being used to address short-term, localized, changing, or ephemeral risks. This approach is applied in areas with irregular overlap of at-risk species with human activities, but where the impact of potential interaction is significant (i.e., *high threat areas*). Success of either framework relies on long-term, persistent biological monitoring using reliable ocean observation assets.

Electric gliders are mobile, cost-effective underwater surveillance tools that can be equipped with sensors for measuring oceanographic conditions and recording marine soundscapes (Webb et al., 2001). Glider deployments fill a unique whale surveillance niche compared to other standard platforms. Like visual surveys, gliders survey along transects, but their temporal effort is significantly higher, with deployments lasting up to six months during which monitoring is continuous, including at night and in all types of weather (Baumgartner et al., 2014, 2020). The mobility of gliders allows for regional-scale spatial surveys of habitats or management areas that span hundreds of kilometers and can be remote, a task not easily achievable with individual passive acoustic monitoring (PAM) moorings. Additionally, all profiling electric gliders carry a standard suite of oceanographic sensors for simultaneously monitoring cetacean acoustics and environmental conditions throughout the water column, which is not standard for PAM moorings or visual surveys (e.g., Ruckdeschel et al., 2020). Thus, gliders fill a unique surveillance role that is required to meet whale management objectives that rely on acoustic and environmental monitoring across seasons and variable spatial scales, including in near-real time.

Gliders equipped with PAM devices are capable of robust near-real-time monitoring of numerous whale species (Baumgartner et al., 2013, 2020). One such species is the North Atlantic right whale (*Eubalaena glacialis*, NARW), which is suffering an ongoing unusual mortality event that resulted in 151 documented mortality, serious injury, and morbidity cases from 2017 to 2024: 41 deaths, 39 serious injuries, and 71 sublethal injuries (note that only about one-third of right whale deaths are thought to be documented; NMFS, 2025). The coastal distribution of NARWs spans calving grounds in the southeastern United States to foraging grounds in northern United States and Canadian

waters, resulting in frequent overlap with high density vessel traffic, major shipping lanes, and commercial fisheries operations. As a result, the leading causes of death and injury for NARWs are vessel strikes and fishing gear entanglements (Sharp et al., 2019). Unpredictable shifts have occurred in NARW distributions in recent years, likely linked to the consequences of climate change impacts on habitat suitability and feeding conditions (Meyer-Gutbrod et al., 2018). This resulted in changes to NARW co-occurrence with human activities as well as to existing protection measures. Therefore, glider effort is expanding over larger temporal and spatial scales to better understand and respond to the dynamic behavior of and persistent threats to this critically endangered species.

Glider-derived acoustic detections can provide information on the occurrence and distribution of NARWs in relation to high threat human activities at hourly to daily timescales. Many near-real-time PAM systems deployed to monitor NARWs (i.e., gliders and moored buoys) use a digital acoustic monitoring (DMON) instrument running a low-frequency detection and classification system (LFDCS; Johnson and Hurst, 2007; Baumgartner and Mussoline, 2011; Baumgartner et al., 2013, 2020) that automatically detects and classifies tonal baleen whale sounds in real time. A subset of these detection data is sent to shore periodically (e.g., when a glider surfaces), enabling acoustic analysts to validate detected whale calls in near-real time following a standard protocol (Figure 1; Wilder et al., 2023). Validated detections are then rapidly disseminated to stakeholders via various automated systems. Here, we highlight five case studies to discuss how DMON/LFDCS-equipped gliders are being used internationally for area-based monitoring of NARWs across habitats and in distinct environments, as well as how the acoustic observations are being used to inform management and/or stakeholder actions to mitigate the impacts of anthropogenic threats (Table 1). The goal of these examples is to illustrate how this platform's unique surveillance niche can help address the complex and multifaceted management needs of a migratory endangered species.

## NARW CONSERVATION CASE STUDIES

### CANADIAN DYNAMIC SHIPPING ZONES

The Gulf of St. Lawrence (GoSL), Canada, recently became a foraging hotspot for NARWs (Meyer-Gutbrod et al., 2021). The GoSL is an inland sea bisected by shipping lanes that serve as the sole oceanic connection between North America's Great Lakes (including Canada's largest city, Toronto) and global ports. Regional overlap of whales and high vessel density has contributed to the species' unusual mortality event (Daoust et al., 2018). To reduce the risk of vessel strikes to NARWs in this high threat area, the Canadian government developed a significant new dynamic management plan for shipping in 2018 (Transport




FIGURE 1. Diagram of whale acoustic detection data flow from a digital acoustic monitoring (DMON) instrument running a low-frequency detection and classification system (LFDCS) integrated into a Slocum glider to a shore-side server via the Iridium satellite service and displayed on a publicly accessible website. After analyst review, the presence of North Atlantic right whales is shared with stakeholders via the website and email/text messages. Depending on the area, dynamic management measures may be implemented in response to whale detections.

Canada, 2024). Beginning in 2020, the plan included the implementation of PAM-equipped glider surveys within deep water (>300 m) dynamic shipping zones to trigger mandatory regional speed restrictions of 10 knots in response to NARW acoustic presence (Figure 2). These surveys are conducted annually from April to November and are done in collaboration with the University of New Brunswick and Dalhousie University.

Vessel slowdowns are implemented or can be extended by regulators when a NARW is detected acoustically (via glider) or visually (via aerial surveillance) within or near a dynamic shipping zone. Slowdowns are initially triggered for a period of 15 days and apply to all vessels >13 m transiting within the active slow zone. If a speed limit is already implemented when a new detection is made, the speed limit is reset for an additional 15-day period starting on the day of the new detection, given that it occurs in the last seven days after the start of the previous slowdown (Transport Canada, 2024). When no speed restrictions are in place in the dynamic shipping zones, vessels can transit at a safe operating speed, which may vary depending on the type of vessel. Most commercial vessels normally transit at speeds over 10 knots.

Over the first four years of this dynamic management plan, there were 30 days with near-real-time acoustic detections of NARWs made during 580 glider survey days in the GoSL, triggering 194 days of dynamic shipping zone slowdowns. We found a high degree of interannual and seasonal variation in NARW acoustic occurrence that likely reflected their transitory use of the shipping lanes, as well as within- and between-season shifts in distribution across the region (recent work of author Indeck and colleagues). Gliders triggered more slowdowns than aerial surveillance by a factor of two to five during fall and summer but were less effective during spring, as whales migrating into the GoSL tend to call at lower rates and occur at lower densities than during other behavioral states (Parks et al., 2011; Matthews and Parks, 2021).

#### CANADIAN DYNAMIC FISHING AREAS

In 2018, Fisheries and Oceans Canada (DFO) initiated a new fishery management plan to mitigate entanglement harm to NARWs from fixed-gear fisheries (primarily snow crab, *Chionoecetes opilio*, and lobster, *Homarus americanus*) in Canadian NARW habitats. Measures included mandatory static zones starting in 2018 and 2019, and dynamic

TABLE 1. Summary of area-based monitoring of North Atlantic right whales (NARWs) across Canada and the United States, highlighting how glider-derived acoustic detections are being used to trigger management actions and/or inform stakeholder decisions to mitigate the impacts of various anthropogenic threats.

THREAT	REGION	FRAME- WORK	COMPLIANCE	ACTION	STAKEHOLDERS*
Vessel strike	Atlantic Canada	Dynamic	Mandatory	15-day, 10-knot slowdown of all vessels >13 m transiting the speed-restricted dynamic shipping zone; this is extended an additional 15 days if a second detection occurs during days 8-15.	<ul style="list-style-type: none"> <li>• Transport Canada</li> <li>• Shipping industry</li> </ul>
Fishing gear entanglement	Atlantic Canada	Dynamic	Mandatory	15-day area closure, including a 72-hour gear removal period; if a second detection occurs during days 9-15, the area is put under a seasonal closure.	<ul style="list-style-type: none"> <li>• Fisheries and Oceans Canada</li> <li>• Snow crab and lobster fisheries</li> </ul>
Fishing gear entanglement	Northeast United States	Static	Mandatory	Annual static closure in Lobster Management Area 1 from October 1 to January 31, where traditional fixed-gear fishing with vertical lines is prohibited, based on seasonal presence of NARWs.	<ul style="list-style-type: none"> <li>• National Oceanic and Atmospheric Administration (NOAA)</li> <li>• Lobster fishery</li> </ul>
Noise exposure, habitat degradation	Northeast United States	N/A	N/A	Comparison of pre-and post-construction data in offshore wind energy development areas will allow their potential impacts on NARWs to be assessed and provide information on the environmental drivers of NARW habitat use.	<ul style="list-style-type: none"> <li>• Bureau of Ocean Energy Management (BOEM)</li> <li>• Wind energy developers</li> </ul>
Vessel strike	Northeast United States	Dynamic	Voluntary	15-day, 10-knot slowdown of all vessels >19.8 m as part of NOAA's Slow Zones for Right Whales program.	<ul style="list-style-type: none"> <li>• NOAA</li> <li>• Northeast US mariners</li> </ul>
Vessel strike	Southeast United States	Dynamic	Voluntary	Early Warning System and communication network for vessel strike mitigation, which alerts nearby vessel traffic of NARW presence shortly after a detection.	<ul style="list-style-type: none"> <li>• NOAA</li> <li>• Southeast US mariners</li> </ul>

\*All projects include state/provincial agencies and/or academic institutions that are integral to the success of monitoring and management objectives.

fishery-area closures that began in 2020, supplemented by increased visual and acoustic survey efforts to detect NARW presence, including the use of Slocum acoustic gliders (Fisheries and Oceans Canada, 2023).

Under the current plan, the GoSL is subdivided into 10 minutes latitude × 10 minutes longitude grids. If any NARW is detected within a grid by any monitoring platform

(vessel, airplane, buoy, glider, or validated opportunistic sighting), a temporary closure area is triggered for a period of 15 consecutive days, including a minimum 48-hour gear removal period (Fisheries and Oceans Canada, 2024). Each closure area is a 3 × 3 grid unit that includes the surveyed cell (i.e., the grid containing the NARW detection) and eight surrounding buffer grids, totaling approximately 2,000 km<sup>2</sup>.




FIGURE 2. Map of the eastern United States and Canada, illustrating North Atlantic right whale (NARW) calving grounds off the southeastern United States and a foraging grounds/migratory corridor that extends along the northeastern United States into Canadian waters. Insets highlight the different regions where gliders are being used for area-based monitoring of NARWs. In Atlantic Canada, glider-derived NARW detections are used in the dynamic management of shipping zones (outlined in blue) and contribute to fishery area closures in the southern Gulf of St. Lawrence (GoSL, yellow circle); the green shading indicates all of Transport Canada’s vessel traffic management areas, which include a restricted area in the southern GoSL (gray shading), a voluntary seasonal slowdown zone in the Cabot Strait to the southeast of the dynamic shipping zones, and static 10-knot speed zones to the north and south of the dynamic shipping zones. In the Gulf of Maine, nine years of glider deployments provided insight on seasonal patterns of NARW presence, which informed the establishment of the region’s seasonal restricted area (gray shading) within Lobster Management Area 1 (LMA-1, outlined in black), where ongoing missions continue to monitor NARW occurrence. Glider missions in the New York Bight play an important role in assessing NARW habitat use relative to offshore wind energy development in the northeastern United States, with deployments in wind planning areas (shaded green polygons), wind lease areas (shaded red polygons), and busy shipping lanes (green outlines), as acoustic detections supplement visual observations in triggering dynamic voluntary slow zones (shaded orange squares). Lastly, in US southeast waters, glider deployments off the coast of Georgia contribute to an early warning system in and around heavily trafficked shipping lanes (green outlines) to mitigate the threat of vessel strike for female NARWs and their newborn calves. Yellow arrows in each inset panel indicate the general geographic span of glider missions conducted in that region.

Buffer grids are included in the trigger to account for NARW movement after the detection is made, because NARWs can travel 80 km d<sup>-1</sup> on average (Baumgartner and Mate, 2005). DFO is then responsible for surveying the closure area with an aerial platform during the 15-day closure. If an NARW is not detected again, visually or acoustically, within the closure area during days 9-15 and after two clearance flights (on separate days) with two trained Marine Mammal Observers on board have been completed, then the area is reopened to fishing on day 16. However, if an NARW is detected within the closure area during days 9-15, the area is put under a seasonal closure, effectively ending fishing in that area for the rest of the monitoring season on November 15 (Fisheries and Oceans Canada, 2024).

Gliders have been used to trigger fishery-area closures in the GoSL each year since 2020. During the first four years (2020-2023), the gliders triggered 13, 46, 21, and 48 grid closures, respectively, comprising a total closed area of approximately 8,700 nm<sup>2</sup> (30,000 km<sup>2</sup>) across years. Both the DFO fisheries and the Transport Canada shipping management plans have been reviewed and adapted every year, as more has been learned about NARW presence and distribution in the GoSL.

#### US MONITORING TO MITIGATE FISHING GEAR ENTANGLEMENTS AND VESSEL STRIKES

Glider-based monitoring of NARWs in US waters serves several purposes, including informing mitigation efforts for fishing gear entanglements and vessel strikes. Near-real-time acoustic detections of NARWs from gliders began in the Gulf of Maine in 2012 (Baumgartner et al., 2013), in a region that in 2021 was designated a seasonal restricted area within Lobster Management Area 1, where traditional fixed-gear fishing with vertical lines is now prohibited annually from October 1 to January 31 because of the seasonal presence of NARWs (Figure 2). Glider-based detections from regular surveys of NARWs conducted by the Woods Hole Oceanographic Institution and The University of Maine were used, in part, to justify this restricted area, as well as to defend its existence in US federal court (Bowling, 2022).

Vessel strike mitigation in the United States currently consists of mandatory vessel speed restrictions in relatively small static management areas for vessels with lengths over 19.8 m, and voluntary vessel speed restrictions dynamically triggered by visual or acoustic detections of NARWs outside of the static management areas. Speed in both areas is limited to 10 knots, and dynamic management areas persist for 15 days. The program to encourage cooperation with voluntary vessel speed restrictions based on near-real-time acoustic detections was established in late 2020 and is called the National Oceanic and Atmospheric

Administration's Slow Zones for Right Whales. In the four years since its inception, 154 Slow Zones have been triggered or extended by acoustic detections of NARWs, and 51 (33%) of those Slow Zones were triggered or extended by gliders operated by the Woods Hole Oceanographic Institution, Rutgers University, Stony Brook University, and The University of Maine during 57 separate glider missions. The remaining Slow Zones were triggered by moored buoys operated by the Woods Hole Oceanographic Institution and carrying the same DMON/LFDCS system as the gliders (Baumgartner et al., 2019).

#### US OFFSHORE WIND DEVELOPMENT AREAS

In pursuit of ambitious renewable energy targets, the United States plans to develop its eastern seaboard with offshore wind energy farms over the upcoming decade. Lease areas in northeastern US waters are in various stages of turbine installation, and there is a coordinated effort between the Bureau of Ocean Energy Management (BOEM), state agencies, wind energy developers, and the scientific community to address the ecological impacts of offshore wind energy development (OWD; Van Parijs et al., 2021). These impacts are anticipated to span the marine food chain through nuanced linkages between the hydrodynamics and food web ecology at turbine, wind energy area, and regional scales (NASEM, 2024).

Within the US Northeast, increases in NARW occurrence have been observed south of traditional foraging grounds in the Gulf of Maine since approximately 2010, in regions where considerable OWD is ongoing or upcoming (Davis et al., 2017; Meyer-Gutbrod et al., 2022). Further, OWD is occurring in regions such as the New York Bight, which has historically received limited survey effort and has lacked density estimates and detailed distributional data for large whales until recently (Zoidis et al., 2021). PAM-equipped gliders operated by Rutgers and Stony Brook Universities are playing a key role in assessing the habitat use of NARWs and other large whales relative to OWD in the northeastern United States.

Since 2020, gliders have surveyed for over 700 days and have transited more than 14,000 km in and adjacent to wind lease areas in New York and New Jersey (Figures 2 and 3). These surveys have documented detections of NARWs on 10%-20% of survey days from November to March, and <5% of survey days from March to October. Continued monitoring, and the comparison of pre- and post-development occurrence data in OWD areas, will allow potential impacts of OWD on NARWs to be assessed. Further, by providing NARW detections, along with concurrently sampled subsurface oceanographic data, glider surveys will help to improve our understanding of the environmental drivers

of NARW habitat use in these previously understudied regions. Given the rapid environmental change occurring in the northeastern United States, this information will be critical to distinguishing impacts of OWD on habitat use from effects of environmental variability.

#### US COASTAL CALVING GROUND SURVEILLANCE

Pregnant NARWs migrate to nearshore southeast US calving grounds, spanning the states of Florida, Georgia, and South Carolina, to give birth and nurse their newborn calves between the months of November and April (Gowan and Ortega-Ortiz, 2014). Non-reproductive individuals also migrate to this calving ground during the winter months (Gowan et al., 2019). Although gliders have been used for near-real-time detections of NARWs in northern foraging grounds off the coast of the United States and Canada for more than a decade, gliders were not used until 2023 for PAM in the southeast US calving ground, where glider-based PAM faces two major challenges. First, vocalization rates of lactating females compared to other demographic groups are lower in the calving ground (Parks et al., 2019b), and these calls tend to be low amplitude (Parks et al., 2019a). This reduces the likelihood of acoustically detecting a mother-calf pair in this region. Second, the calving ground is situated close to the coast over a shallow portion of the inner continental shelf. Typical depth occupancy in the calving ground is between 10 m and 25 m, which provides limited vertical space for a glider to operate its dive-climb flight pattern. The frequent shift between ascending and descending status requires more frequent engagement of the buoyancy pump, which is both energetically costly and produces self-noise that may mask whale vocalizations. Further, strong gradients in temperature, salinity, and thus sound speed, may further limit detection range.

Despite these challenges, a pilot program using gliders for NARW PAM was recently implemented through a collaboration between the University of South Carolina and the Skidaway Institute of Oceanography. So far, these efforts consisted of a two-week mission in January 2023, and two four-week missions from January through March 2024 (Figures 2 and 3). During these three missions, which operated in water as shallow as 11 m, three definite NARW detections were made. The NARW detection on January 20, 2024, was the first definite glider-based acoustic detection in southeast US waters, and it triggered an alert from the Southeast Early Warning System notifications program for vessel strike mitigation. These pilot missions indicate that glider-based PAM may be a useful tool for supplementing aerial-based detections of NARWs in the calving ground, especially providing coverage when aerial surveillance is not feasible due to weather or other logistical constraints.




FIGURE 3. Photos of gliders in the field show them, from the top, ready for deployment in the Gulf of St. Lawrence, Canada, being deployed by field personnel on the coastal calving grounds of Georgia, USA, and in the water during deployment in the New York Bight, USA.

#### CONSERVATION IMPLICATIONS

North Atlantic right whales are at risk of extinction before the end of the century, as climate change continues to initiate distributional and behavioral changes that inadvertently increase mortality due to vessel strikes and entanglements (Meyer-Gutbrod et al., 2021). As a result, the Canadian and US governments are investing millions of dollars in technologies to support species monitoring, research to better predict future whale distributions, and mitigation efforts to address complex threats to vulnerable species. In the last five years, glider efforts have rapidly expanded, with cumulative deployments totaling thousands of days across the NARW migratory range for conservation applications (Figure 3). Here, we highlighted several examples of how PAM-equipped underwater gliders are being used for

vessel strike and entanglement mitigation by enhancing risk reduction for dynamic management areas, regional fisheries, and offshore wind energy projects (Table 1). Going forward, this technology has the capacity to contribute to more spatial conservation strategies, such as marine protected areas. These areas tend to encompass vast and/or remote areas that are logistically difficult for personnel to survey, which can hamper authorities' ability to enforce protective measures. Therefore, the use of gliders is likely to continue expanding into the future and across the marine domain in support of species conservation and threat mitigation.

Glider-based PAM offers key spatial monitoring capabilities for NARW threat management but is typically used in conjunction with other, complementary monitoring platforms, such as aerial surveillance and PAM moorings. Because NARWs are an internationally mobile cetacean species that spans diverse habitats and protection measures, surveillance assets must fulfill different requirements (e.g., temporal effort, spatial scale, deployment location, data type) across the NARW range, depending on the goal(s) of each individual monitoring program. Because the space-time needs of successful range-wide management are so complex, no one tool could possibly achieve all monitoring imperatives. As one example, gliders were deployed in the Cabot Strait voluntary seasonal slowdown zone in the GoSL for two years. Despite being a known high-use migratory corridor, we did not acoustically detect any NARWs in near-real time. This may have been because of behaviorally influenced calling rates, missed whales traveling close to shore (i.e., deployment location vs. whale movements), or platform type (e.g., mid-endurance mobile glider vs. long-endurance stationary array). Thus, glider-based PAM is being used alongside a suite of other tools, including moored PAM, visual monitoring, and distribution modeling, to aid conservation goals.

We have presented several different area-based threats, management goals, and mitigation plans across glider survey regions. These highlight the need for continued research/support for additional and/or new monitoring platforms (including gliders) to be incorporated into existing and future management plans for the conservation of NARWs. However, the efficacy of glider detections at achieving conservation goals depends largely on the regional regulatory measures in place being informed by these detections. For example, in a recent study, the average percentage of mariners found to be cooperating with 10-knot speed requests in US dynamic management areas was less than 50%, compared to higher compliance in some, but not all, mandatory seasonal management areas (>85% in most areas, but <25% for the largest commercial vessels outside four ports in the southeastern United States; NMFS,

2020). In contrast, Canada has made mitigation measures in the dynamic shipping zones of the GoSL mandatory and achieved 99% compliance during the 2023 monitoring season (Bilodeau, 2023). Furthermore, slowdowns in eastern Canadian waters affect vessels down to 13 m, whereas US speed restrictions currently only apply to vessels that are greater than or equal to 19.8 m. This difference is significant, as four of 13 vessel-related NARW deaths in US waters since 2008 were attributable to boats less than 20 m and, therefore, not subject to slowdown measures (Redfern, 2023). No matter how capable a technology or monitoring system, its overall conservation impact is intertwined with the prevailing management policies.

## REFERENCES

- Baumgartner, M.F., and B.R. Mate. 2005. Summer and fall habitat of North Atlantic right whales (*Eubalaena glacialis*) inferred from satellite telemetry. *Canadian Journal of Fisheries and Aquatic Sciences* 62(3):527-543, <https://doi.org/10.1139/f04-238>.
- Baumgartner, M.F., and S.E. Mussoline. 2011. A generalized baleen whale call detection and classification system. *The Journal of the Acoustical Society of America* 129(5):2,889-2,902, <https://doi.org/10.1121/1.3562166>.
- Baumgartner, M.F., D.M. Fratantoni, T.P. Hurst, M.W. Brown, T.V. Cole, S.M. Van Parijs, and M. Johnson. 2013. Real-time reporting of baleen whale passive acoustic detections from ocean gliders. *The Journal of the Acoustical Society of America* 134(3):1,814-1,823, <https://doi.org/10.1121/1.4816406>.
- Baumgartner, M.F., K.M. Stafford, P. Winsor, H. Statscewich, and D.M. Fratantoni. 2014. Glider-based passive acoustic monitoring in the Arctic. *Marine Technology Society Journal* 48(5):40-51, <https://doi.org/10.4031/MTSJ.48.5.2>.
- Baumgartner, M.F., J. Bonnell, S.M. Van Parijs, P.J. Corkeron, C. Hotchkin, K. Ball, L.P. Pelletier, J. Partan, D. Peters, J. Kemp, and others. 2019. Persistent near real-time passive acoustic monitoring for baleen whales from a moored buoy: System description and evaluation. *Methods in Ecology and Evolution* 10(9):1,476-1,489, <https://doi.org/10.1111/2041-210X.13244>.
- Baumgartner, M.F., J. Bonnell, P.J. Corkeron, S.M. Van Parijs, C. Hotchkin, B.A. Hodges, J. Bort Thornton, B.L. Mensi, and S.M. Bruner. 2020. Slocum gliders provide accurate near real-time estimates of baleen whale presence from human-reviewed passive acoustic detection information. *Frontiers in Marine Science* 7:100, <https://doi.org/10.3389/fmars.2020.00100>.
- Bilodeau, M. 2023. Transport Canada (TC) update on North Atlantic right whale vessel management measures. Presentation given at the annual North Atlantic Right Whale Consortium, October 24-25, 2023, Halifax, NS, Canada.
- Bowling, T. 2022. Court reinstates seasonal closure of Maine lobster fishery to protect whales. *The SandBar* 21(1):4-5.
- Daoust, P.Y., E.L. Couture, T. Wimmer, and L. Bourque. 2018. *Incident Report: North Atlantic Right Whale Mortality Event in the Gulf of St. Lawrence, 2017*. Collaborative report produced by Canadian Wildlife Health Cooperative, Marine Animal Response Society, and Fisheries and Oceans Canada, 256 pp.
- Davis, G.E., M.F. Baumgartner, J.M. Bonnell, J. Bell, C. Berchok, J. Bort Thornton, S. Brault, G. Buchanan, R.A. Charif, D. Cholewiak, and others. 2017. Long-term passive acoustic recordings track the changing distribution of North Atlantic right whales (*Eubalaena glacialis*) from 2004 to 2014. *Scientific Reports* 7(1):13460, <https://doi.org/10.1038/s41598-017-13359-3>.
- Fisheries and Oceans Canada. 2023. "Protecting North Atlantic Right Whales: Canada's Fishing Measures by Year Launched," <https://www.dfo-mpo.gc.ca/about-notre-sujet/publications/infographics-infographies/narw-bnan-by-year-par-annee-eng.html>.

- Fisheries and Oceans Canada. 2024. "2024 Fishery Management Measures," <https://www.dfo-mpo.gc.ca/fisheries-peches/commercial-commerciale/atl-arc/narw-bnan/management-gestion-eng.html>.
- Gowan, T.A., and J.G. Ortega-Ortiz. 2014. Wintering habitat model for the North Atlantic right whale (*Eubalaena glacialis*) in the southeastern United States. *PLoS ONE* 9(4):e95126, <https://doi.org/10.1371/journal.pone.0095126>.
- Gowan, T.A., J.G. Ortega-Ortiz, J.A. Hostetler, P.K. Hamilton, A.R. Knowlton, K.A. Jackson, R.C. George, C.R. Taylor, and P.J. Naessig. 2019. Temporal and demographic variation in partial migration of the North Atlantic right whale. *Scientific Reports* 9(1):353, <https://doi.org/10.1038/s41598-018-36723-3>.
- Johnson, M., and T. Hurst. 2007. The DMON: An open-hardware/open-software passive acoustic detector. Paper presented at the 3rd International Workshop on the Detection and Classification of Marine Mammals Using Passive Acoustics, July 24–26, 2007, Boston, MA, USA.
- Matthews, L.P., and S.E. Parks. 2021. An overview of North Atlantic right whale acoustic behavior, hearing capabilities, and responses to sound. *Marine Pollution Bulletin* 173(Pt B):113043, <https://doi.org/10.1016/j.marpolbul.2021.113043>.
- Maxwell, S.M., E.L. Hazen, R.L. Lewison, D.C. Dunn, H. Bailey, S.J. Bograd, D.K. Briscoe, S. Fossette, A.J. Hobday, M. Bennett, and others. 2015. Dynamic ocean management: Defining and conceptualizing real-time management of the ocean. *Marine Policy* 58:42–50, <https://doi.org/10.1016/j.marpol.2015.03.014>.
- Meyer-Gutbrod, E., C. Greene, and K. Davies. 2018. Marine species range shifts necessitate advanced policy planning: The case of the North Atlantic right whale. *Oceanography* 31(2):19–23, <https://doi.org/10.5670/oceanog.2018.209>.
- Meyer-Gutbrod, E.L., C.H. Greene, K.T.A. Davies, and D.G. Johns. 2021. Ocean regime shift is driving collapse of the North Atlantic right whale population. *Oceanography* 34(3):22–31, <https://doi.org/10.5670/oceanog.2021.308>.
- Meyer-Gutbrod, E.L., K.T.A. Davies, C.L. Johnson, S. Plourde, K.A. Sorochan, R.D. Kenney, C. Ramp, J.F. Gosselin, J.W. Lawson, and C.H. Greene. 2022. Redefining North Atlantic right whale habitat-use patterns under climate change. *Limnology and Oceanography* 68:571–586, <https://doi.org/10.1002/lno.12242>.
- NASEM (National Academies of Science, Engineering, and Medicine). 2024. *Potential Hydrodynamic Impacts of Offshore Wind Energy on Nantucket Shoals Regional Ecology: An Evaluation from Wind to Whales*. The National Academies Press, Washington, DC, <https://doi.org/10.17226/27154>.
- NMFS (National Marine Fisheries Service). 2020. *North Atlantic Right Whale (Eubalaena glacialis) Vessel Speed Rule Assessment*. National Marine Fisheries Service, Office of Protected Resources, Silver Spring, MD, 53 pp.
- NMFS. 2025. "2017–2025 North Atlantic Right Whale Unusual Mortality Event," <https://www.fisheries.noaa.gov/national/marine-life-distress/2017-2025-north-atlantic-right-whale-unusual-mortality-event>.
- Parks, S.E., A. Searby, A. Célérier, M.P. Johnson, D.P. Nowacek, and P.L. Tyack. 2011. Sound production behavior of individual North Atlantic right whales: Implications for passive acoustic monitoring. *Endangered Species Research* 15(1):63–76, <https://doi.org/10.3354/esr00368>.
- Parks, S.E., D.A. Cusano, S.M. Van Parijs, and D.P. Nowacek. 2019a. Acoustic crypsis in communication by North Atlantic right whale mother-calf pairs on the calving grounds. *Biology Letters* 15(10):20190485, <https://doi.org/10.1098/rsbl.2019.0485>.
- Parks, S.E., D.A. Cusano, S.M. Van Parijs, and D.P. Nowacek. 2019b. North Atlantic right whale (*Eubalaena glacialis*) acoustic behavior on the calving grounds. *The Journal of the Acoustical Society of America* 146(1):EL15–EL21, <https://doi.org/10.1121/1.5115332>.
- Redfern, J.V. 2023. *Examining the Impacts of the National Oceanic and Atmospheric Administration's Proposed Changes to the North Atlantic Right Whale Vessel Strike Reduction Rule*. United States House of Representatives Natural Resources Subcommittee on Water, Wildlife, and Fisheries, Washington, DC, 12 pp.
- Ruckdeschel, G.S., K.T.A. Davies, and T. Ross. 2020. Biophysical drivers of zooplankton variability on the Scotian Shelf observed using profiling electric gliders. *Frontiers in Marine Science* 7:627, <https://doi.org/10.3389/fmars.2020.00627>.
- Sharp, S.M., W.A. McLellan, D.S. Rotstein, A.M. Costidis, S.G. Barco, K. Durham, T.D. Pitchford, K.A. Jackson, P.Y. Daoust, T. Wimmer, and others. 2019. Gross and histopathologic diagnoses from North Atlantic right whale *Eubalaena glacialis* mortalities between 2003 and 2018. *Diseases of Aquatic Organisms* 135(1):1–31, <https://doi.org/10.3354/dao03376>.
- Transport Canada. 2024. "Protecting North Atlantic right whales from collisions with vessels in the Gulf of St. Lawrence," <https://tc.canada.ca/en/marine-transportation/navigation-marine-conditions/protecting-north-atlantic-right-whales-collisions-vessels-gulf-st-lawrence>.
- Van Parijs, S.M., K. Baker, J. Carduner, J. Daly, G.E. Davis, C. Esch, S. Guan, A. Scholik-Schlomer, N.B. Sisson, and E. Staaterman. 2021. NOAA and BOEM minimum recommendations for use of passive acoustic listening systems in offshore wind energy development monitoring and mitigation programs. *Frontiers in Marine Science* 8, <https://doi.org/10.3389/fmars.2021.760840>.
- Webb, D.C., P.J. Simonetti, and C.P. Jones. 2001. SLOCUM: An underwater glider propelled by environmental energy. *IEEE Journal of Oceanic Engineering* 26(4):447–452, <https://doi.org/10.1109/48.972077>.
- Wilder, J., G. Davis, A. DeAngelis, S. Van Parijs, and M. Baumgartner. 2023. *Low-Frequency Detection and Classification System (LFDCS) Reference Guide*. National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Northeast Fisheries Science Center, Woods Hole, Massachusetts, 142 pp.
- Zoidis, A.M., K.S. Lomac-MacNair, D.S. Ireland, M.E. Rickard, K.A. McKown, and M.D. Schlesinger. 2021. Distribution and density of six large whale species in the New York Bight from monthly aerial surveys 2017 to 2020. *Continental Shelf Research* 230:104572, <https://doi.org/10.1016/j.csr.2021.104572>.

## ACKNOWLEDGMENTS

We would like to thank the following for funding support: Transport Canada, Fisheries and Oceans Canada, the Ocean Tracking Network, Marine Environmental Observation, Prediction and Response Network, the Murphy Family Foundation, the Broad Reach Fund, NOAA Northeast Fisheries Science Center, the Northeast Regional Association of Coastal Ocean Observing Systems, the New York State Environmental Protection Fund Ocean and Great Lakes Program via the New York State Department of Environmental Conservation, the New York State Energy Research and Development Authority, the Mid-Atlantic Regional Association Coastal Ocean Observing System, the Tides Foundation, and the Southeast Coastal Ocean Observing Regional Association.

We would also like to express our sincere appreciation to the following individuals, who were key to glider operations and mission support: A. Comeau, J. van der Meer, Z. Viva, B. Hodges, M. Meier, J. Wilder, C. Flagg, D. Aragon, N. Waite, J. Kohut, T. Miles, A. Kreuser, A. Sefah-Twerefour, K. Dreger, J. Bird, F. McQuarrie, and A. Vincent.

## AUTHORS

**Katherine L. Indeck** ([katherine.indeck@unb.ca](mailto:katherine.indeck@unb.ca)), University of New Brunswick, Saint John, NB, Canada. **Mark F. Baumgartner**, Woods Hole Oceanographic Institution, Woods Hole, MA, USA. **Laurence Lecavalier**, Transport Canada Innovation Centre, Ottawa, ON, Canada. **Frederick Whoriskey**, Ocean Tracking Network, Halifax, NS, Canada. **Delphine Durette-Morin**, Canadian Whale Institute, Welshpool, NB, Canada. **Neal R. Pettigrew**, School of Marine Sciences, The University of Maine, Orono, ME, USA. **Jacqueline M. McSweeney**, **Lesley H. Thorne**, and **Katherine L. Gallagher**, Stony Brook University, Stony Brook, NY, USA. **Catherine R. Edwards**, Skidaway Institute of Oceanography, University of Georgia, Savannah, GA, USA. **Erin Meyer-Gutbrod**, School of the Earth, Ocean, and Environment, University of South Carolina, Columbia, SC, USA. **Kimberley T.A. Davies**, University of New Brunswick, Saint John, NB, Canada.

**ARTICLE DOI.** <https://doi.org/10.5670/oceanog.2025e111>