

OCS-A  
0501



MASS  
USA

# VINEYARD WIND

## Draft Construction and Operations Plan

### Volume III Text

## Vineyard Wind Project

June 3, 2020

**Submitted by**

**Vineyard Wind LLC**  
700 Pleasant Street, Suite 510  
New Bedford, Massachusetts 02740

**Submitted to**

**Bureau of Ocean Energy Management**  
45600 Woodland Road  
Sterling, Virginia 20166

**Prepared by**

**Epsilon Associates, Inc.**  
3 Mill & Main Place, Suite 250  
Maynard, Massachusetts 01754

# Draft Construction and Operations Plan

## Volume III Text

### Vineyard Wind Project

*Submitted to:*

**BUREAU OF OCEAN ENERGY MANAGEMENT**  
45600 Woodland Rd  
Sterling, VA 20166

*Submitted by:*

**VINEYARD WIND LLC**  
700 Pleasant Street, Suite 510  
New Bedford, MA 02740

*Prepared by:*

**EPSILON ASSOCIATES, INC.**  
3 Mill & Main Place, Suite 250  
Maynard, MA 01754

*In Association with:*

Baird & Associates  
Biodiversity Research Institute  
C2Wind  
Capitol Air Space Group  
Clarendon Hill Consulting  
Ecology and Environment  
Foley Hoag  
Geo SubSea LLC  
Gray & Pape

JASCO Applied Sciences  
Morgan, Lewis & Bockius LLP  
Public Archaeology Laboratory, Inc.  
RPS  
Saratoga Associates  
Swanson Environmental Associates  
Wood Thilsted Partners Ltd  
WSP

June 3, 2020

## Section 9.0

---

---

References

## 9.0 REFERENCES

---

- Abend, A. G., & Smith, T. D. (1999). Review of distribution of the long-finned pilot whale (*Globicephala melas*) in the North Atlantic and Mediterranean. NOAA Technical Memorandum NMFS-NE-117.
- Abraham, B. J., & Dillon, P. L. (1986). Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (mid-Atlantic)—softshell clam. US Fish and Wildlife Service. 82(11.68). US Army Corps of Engineers, TR EL-82-4. 18 pp. Retrieved from [http://www.nwrc.usgs.gov/wdb/pub/species\\_profiles/82\\_11-068.pdf](http://www.nwrc.usgs.gov/wdb/pub/species_profiles/82_11-068.pdf)
- ACS (American Cetacean Society). (2016). Pilot whale. Retrieved from [http://www.acsonline.org/index.php?option=com\\_content&view=article&id=65:pilot-whale&catid=20:site-content](http://www.acsonline.org/index.php?option=com_content&view=article&id=65:pilot-whale&catid=20:site-content). Accessed November 22, 2017.
- Adams, J., Kelsey, E. C., Felis, J. J., & Pereksta. (2016). Collision and displacement vulnerability among marine birds of the California Current System associated with offshore wind energy infrastructure: US. Geological Survey Open-File Report 2016-1154, 116 p., <http://dx.doi.org/10.3133/ofr20161154>
- Adams, T.P., Miller, R.G., Aleynik, D., and Burrows, M.T. (2014). Offshore marine renewable energy devices as stepping stones across biogeographical boundaries. *J. Appl. Ecol.* 51: 330–338.
- AECOM. (2012). ENV12 CZM 01 Benthic Infaunal Analysis Report. Final Report prepared for Massachusetts Office of Coastal Zone Management. Retrieved from <http://www.mass.gov/eea/docs/czm/seafloor/benthic-infauna-report-2011.pdf>
- Aguilar de Soto, N.A., Delorme, N., Atkins, J., Howard, S., Williams, J., Johnson, M. (2013) Anthropogenic noise causes body malformations and delays development in marine larvae. *Scientific Reports* 3(2831):1–5.
- Ahlen, I., Baagoe, H. J. & Bach, L. (2009). Behavior of Scandinavian bats during migration and foraging at sea. *Journal of Mammalogy*, 90, 1318-1323
- Ainley, D. G., Nettleship, D. N., Carter, H. R., & Storey, A. E. (2002). Common Murre (*Uria aalge*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY.
- Alerstam, T. (1985). Strategies of migratory flight, illustrated by Arctic and Common Terns, *Sterna paradisaea* and *Sterna hirundo*. *Contributions in Marine Science* 27:580–603.

- Algonquin Gas Transmission Company (2000). US Army Corps of Engineers Section 404/10 Permit Application for the HubLine Pipeline Project. November 30, 2000.
- Alpine Ocean Seismic Surveying, Inc. (2017). Vineyard Wind HRG Survey – Field Verification and Vessel Signature Report. Survey Report for Alpine Ocean Seismic Survey Inc. on behalf of Vineyard Wind LLC. Gardline Report Ref 10878.
- American Wind Wildlife Institute ("AWWI"). (2016). Wind turbine interactions with wildlife and their habitats: a summary of research results and priority questions. (Updated June 2016). Washington, DC. Retrieved from [www.awwi.org](http://www.awwi.org)
- Amirault-Langlais, D. L., Imlay, T. L., & Boyne, A. W. (2014). Dispersal Patterns Suggest Two Breeding Populations of Piping Plovers in Eastern Canada. *The Wilson Journal of Ornithology* 126:352–359. doi: 10.1676/13-056.1
- Anderson, E.P. & Mackas, D.L. (1986). Lethal and sublethal effects of a molybdenum mine tailing on marine zooplankton: mortality, respiration, feeding and swimming behavior in *Calanus marshallae*, *Metridia pacifica* and *Euphausia pacifica*. *Marine Environmental Research* 19:131-155.
- Anderson, L. W., & Olsen, M. T. (2010). Distribution and population structure of North Atlantic harbour seals (*Phoca vitulina*). *NAMMCO Scientific Publications*, 8, 173-188.
- André, M., Solé, M., Lenoir, M., Durfort, M., Quero, C., Mas, A., Lombarte, A., van der Schaar, M., López-Bejar, M., Morell, M., Zaugg, S., & Houégnigan, L. (2011). Low-frequency sounds induce acoustic trauma in cephalopods. *Frontiers in Ecology and the Environment*, 10, 18-28.
- Arnett, E. B., Brown, W. K., Erickson, W. P., Fieldler, J. K., Hamilton, B. L., Henry, T. H., Jain, A., Johnson, G. D., Kerns, J., Koford, R. R., Nicholson, C. P., O'Connell, T. J., Piorkowski, M. D. & Tankersley, R. D. (2008). Patterns of Bat Fatalities at Wind Energy Facilities in North America. *Journal of Wildlife Management*, 72, 61-78
- Atlantic States Marine Fisheries Commission (ASMFC). (1998). Interstate Fishery Management Plan for Horseshoe Crab. Fishery Management Report No. 32 of Atlantic States Marine Fisheries Commission. Retrieved from <http://www.asmfc.org/uploads/file/hscFMP.pdf>.
- Atlantic States Marine Fisheries Commission (ASMFC). (2015). Interstate Fishery Management Plan for Jonah Crab. NOAA Award No. NA15NMF4740069.
- Atlantic States Marine Fisheries Commission (ASMFC). (2010). Species Profile: Horseshoe Crab- New Assessment Finds Trends in Horseshoe Crab Populations Vary by Region. *Fisheries Focus* 19. [Online.] Retrieved from <http://www.asmfc.org/uploads/file/hscProfile.pdf>

- Atlantic States Marine Fisheries Commission (ASMFC). (2018). Jonah Crab. [Online.] Retrieved from [http://www.asmfc.org/uploads/file/5a70a176Jonah\\_Crab.pdf](http://www.asmfc.org/uploads/file/5a70a176Jonah_Crab.pdf)
- Atlantic Sturgeon Status Review Team (AASRT). (2007). Status Review of Atlantic sturgeon (*Acipenser oxyrinchus oxyrinchus*). Report to National Marine Fisheries Service, Northeast Regional Office. February 23, 2007. 174 pp.
- Audubon. (2017). Guide to North American Birds. [Online.] Retrieved from <http://www.audubon.org/bird-guide>
- Bailey, H., Senior, B., Simmons, D., Rush, J., Picken, G., & Thompson, P. (2010). Assessing underwater noise levels during pile-driving at offshore windfarm and its potential effects on marine mammals. *Marine Pollution Bulletin*, 60, 888-897.
- Baird, R. W., & Stacey, P. J. (1990). Status of Risso's dolphin, *Grampus griseus*, in Canada. *Can. Field-Nat.*, 105, 233-242.
- Baker, A., Gonzalez, P., Morrison, R. I. G., & Harrington, B. A. Harrington. (2013). Red Knot (*Calidris canutus*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, New York. doi: 10.2173/bna.563
- Baldassarre, G. A., & Bolen, E. G. (2006). *Waterfowl Ecology and Management*. In. 2nd edition. Krieger, Malabar FL.
- Barber, M. R. (2017). *Effects of Hydraulic Dredging and Vessel Operation on Atlantic Sturgeon Behavior in a Large Coastal River*. Masters Thesis from Virginia Commonwealth University, Richmond, Virginia.
- Barbour, R. W. & Davis, W. H. (1969). *Bats of America*, Lexington, Univ. Press of Kentucky.
- Barco, S., Law, M., Drummond, B., Koopman, H., Trapani C., Reinheimer S., ... & Williard, A. (2016). Loggerhead turtles killed by vessel and fishery interactions in Virginia, USA, are healthy prior to death. *Marine Ecology Progress Series*, 555, 221-234.
- Barkaszi, M.J., Butler, M., Compton, R., Unietis, A., & Bennet, B. (2012). Seismic survey mitigation measures and marine mammal observer reports. OSC Study BOEM 2012-015. Prepared under BOEM Contract M08PC20051 by GeoCet Group, LLC, Houston, Texas.
- Barlas, M. E. (1999). The distribution and abundance of harbor seals (*Phoca vitulina concolor*) and gray seals (*Halichoerus grypus*) in southern New England, winter 1998-summer 1999. Graduate School of Arts and Sciences, Boston University.

- Bartol, S. M., & Ketten, D. R. (2006). Turtle and tuna hearing. In Swimmer, Y., & Brill, R. (eds.). Sea turtle and pelagic fish sensory biology: Developing techniques to reduce sea turtle bycatch in longline fisheries. pp. 98-103. NOAA Technical Memorandum NMFS-PIFSC-7. NMFS, Pacific Islands Fisheries Science Center, Honolulu, Hawaii.
- Bartol, S. M., Musick, J. A., & Lenhardt, M. (1999). Auditory evoked potentials of the loggerhead sea turtle (*Caretta caretta*). *Copeia* 1999(3), 836-840.
- Baruah, E. (2016). A Review of the Evidence of Electromagnetic Field (EMF) Effects on Marine Organisms. *Research & Reviews: Journal of Ecology and Environmental Sciences*, 4(4), 22-26.
- Bauer, G. B., Fuller, M., Perry, A., Dunn, J. R., & Zoeger, J. (1985). *Magnetoreception and biomineralization of magnetite in cetaceans*. In Kirschvink, J. L., Jones, D. S., & MacFadden, B. J. (eds.). *Magnetic biomineralization and magnetoreception in living organisms*. pp. 487-507. New York, New York: Plenum Press.
- Beamish, P. (1979). Behavior and significance of entrapped baleen whales. In Winn, H. E., & Olla, B. L. (eds.). *Behavior of marine animals, Volume 3: Cetaceans*, pp. 291-309. New York, New York: Plenum Press.
- Beamish, P., & Mitchell, E. (1973). Short pulse length audio frequency sounds recorded in the presence of a Minke whale (*Balalenopectera acutorostrata*). *Deep-Sea Research and Oceanographic Abstracts*, 20, 375- 386.
- Beauchamp, K., & Geist, M. (2011). *Agricultural Use of Cape Cod: Looking to the Future*. Association to Preserve Cape Cod. Barnstable, MA.
- Bergström, L., Sundqvist, F., & Bergström, U. (2013). Effects of an offshore wind farm on temporal and spatial patterns in the demersal fish community. *Marine Ecology Progress Series*, 485, 199-210.
- Berlin, A. M., Savoy, L., Gray, C. E., Gilbert, A. T., Olsen, G. H., Ford, S., Montevecchi, W. A., & Stenhouse, I. J. (2017). Occurrence and Migration of Surf Scoters Wintering in Offshore Waters of the Mid-Atlantic United States. In *Determining Fine-Scale Use and Movement Patterns of Diving Bird Species in Federal Waters of the Mid-Atlantic United States Using Satellite Telemetry* (C. S. Spiegel, Editor). US. Department of the Interior, Bureau of Ocean Energy Management, Division of Environmental Sciences, Sterling, VA.
- Berry, W.J., Rubinstein, N.I., Hinchey, E.H., Klein-MacPhee, G., & Clarke, D.G. (2011). Assessment of dredging-induced sedimentation effects on Winter Flounder (*Pseudopleuronectes americanus*) hatching success: results of laboratory investigations. Proceedings of the Western Dredging Association Technical Conference and Texas A&M Dredging Seminar, Nashville, Tennessee, June 5-8, 2011. Retrieved from [http://nerdt.org/wp-content/uploads/2014/12/WEDA\\_Berry.pdf](http://nerdt.org/wp-content/uploads/2014/12/WEDA_Berry.pdf)

- Bevan, E., Wibbels, T., Najera, B. M., Sarti, L., Martinez, F. I., Cueva, J. M., ... & Burchfield, P. M. (2016). Estimating the historic size and current status of the Kemp's ridley sea turtle (*Lepidochelys kempii*) population. *Ecosphere* 7(3), e01244.
- Bierregaard, R. (2017). Rob Bierregaard's Website. [Online.] Retrieved from <http://www.ospreytrax.com/>.
- Bigg, M. A. (1981). Harbor seal - *Phoca vitulina* and *P. largha*. In Ridgway, S. H., & Harrison, R. J. (eds.). Handbook of marine mammals. London: Academic Press.
- Bjorndal, K. A. (1997). Foraging ecology and nutrition of sea turtles. In: Lutz, P. L. & Musick, J. A. (eds.). The biology of sea turtles. Boca Raton, Florida: CRC Press. pp. 199-231.
- Blodget, B. C. (2002). Bird List for the Commonwealth of Massachusetts. In. Massachusetts Division of Fisheries and Wildlife.
- BOEM. (2007). Programmatic Environmental Impact Statement for Alternative Energy Development and Production and Alternate Use of Facilities on the Outer Continental Shelf. Final Environmental Impacts Statement. OCS EIS/EA MMS 2007-046. Retrieved from <https://www.boem.gov/Guide-To-EIS/>
- BOEM. (2012). Commercial wind lease issuance and site assessment activities on the Atlantic Outer Continental Shelf offshore Rhode Island and Massachusetts: Environmental Assessment. [Online.] Retrieved from [http://www.boem.gov/uploadedFiles/BOEM/Renewable\\_Energy\\_Program/Smart\\_from\\_the\\_Start/BOEM\\_RI\\_MA\\_EA\\_2012-070.pdf](http://www.boem.gov/uploadedFiles/BOEM/Renewable_Energy_Program/Smart_from_the_Start/BOEM_RI_MA_EA_2012-070.pdf)
- BOEM. (2013). Collaborative archaeological investigations and sound source verifications within the Massachusetts Wind Energy Area. U.S. Department of the Interior, Bureau of Ocean Energy, Office of Renewable Energy Programs, Herndon, VA. OCS Study 201-221.
- BOEM. (2014). *Commercial Wind Lease Issuance and Site Assessment Activities on the Atlantic Outer Continental Shelf Offshore Massachusetts Revised Environmental Assessment*. U.S. Department of the Interior, Bureau of Ocean Energy Management, Office of Renewable Energy Programs. OCS EIS/EIA BOEM 2014-603.
- BOEM. (2017). Gulf of Mexico OCS proposed geological and geophysical activities western, central, and eastern planning areas, Final Programmatic Environmental Impact Statement. OCS EIA/EA BOEM 2017-051.
- BOEMRE. (2011) Literature Synthesis for the North and Central Atlantic Ocean. U.S. Dept. of the Interior, Bureau of Ocean Energy Management, Regulation and Enforcement, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study BOEMRE 2011-012. 447 pp.



- Bonner, W. N. (1971). Grey seal *Halichoerus grypus fabricus*. In Ridgway, S. H., & Harrison, R. J. (eds.). Handbook of marine mammals. London: Academic Press.
- Bonner, W. N. (1990). The natural history of seals. Facts on File, New York. 196 pp.
- BOWind. (2008) Barrow Offshore Wind Farm Post Construction Monitoring Report. First annual report. 15 January 2008, 60pp.
- Brandt, M. J., Diederichs, A., Betke, K., & Nehls, G. (2011). Responses of harbour porpoises to pile driving at the Horns Rev II offshore wind farm in the Danish North Sea. *Marine Ecology Progress Series*, 421, 205-216.
- Broders, H. G. & Forbes, G. J. 2004. Interspecific and intersexual variation in roost-site selection of northern long-eared and little brown bats in the Greater Fundy National Park ecosystem. *Journal of Wildlife Management*, 68, 602-610.
- Brodin, Y. & Andersson, M.H. (2009). The marine splash midge *Telmatogon japonicus* (Diptera; chironomidae) – extreme and alien? *Biological Invasions*, 11(6), 1311-1317.
- Brooks, R. T. & Ford, W. M. (2005). Bat activity in a forest landscape of central Massachusetts. *Northeastern Naturalist*, 12, 447-462.
- Brown, M. W., Fenton, D., Smedbol, K., Merriman, C., Robichaud-Leblanc, K., & Conway, J. D. (2009). Recovery strategy for the North Atlantic right whale (*Eubalaena glacialis*) in Atlantic Canadian waters [Final]. Fisheries and Oceans Canada.
- Brown, M. W., & Marx, M. K. (2000). Surveillance, monitoring, and management of North Atlantic right whales, *Eubalaena glacialis*, in Cape Cod Bay, Massachusetts: January to mid-May, 2000. Final Report submitted to Division of Marine Fisheries, Department of Fisheries, Wildlife and Environmental Law Enforcement, Boston, Massachusetts. Contract No. SCFWE3000-8365027.
- Bruderer, B., & F. Lietchi. (1999). Bird migration across the Mediterranean. In Proceedings of the 22nd International Ornithological Congress (N. J. Adams and R. H. Slotow, Editors). Durban, Johannesburg, South Africa, pp. 1983–1999.
- Buehler, D. A. (2000). Bald Eagle (*Haliaeetus leucocephalus*). In The Birds of North America (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, New York. doi: 10.2173/bna.506
- Burger, J. (2015). Laughing Gull (*Leucophaeus atricilla*). In The Birds of North America (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca.

- Burger, J., Gordon, C., Lawrence, J., Newman, J., Forcey, G. , & Vlietstra, L. (2011). Risk evaluation for federally listed (roseate tern, piping plover) or candidate (red knot) bird species in offshore waters: A first step for managing the potential impacts of wind facility development on the Atlantic Outer Continental Shelf. *Renewable Energy* 36:338–351. doi: 10.1016/j.renene.2010.06.048
- Bull, L. S., S. Fuller, & D. Sim (2013). Post-construction avian mortality monitoring at Project West Wind. *New Zealand Journal of Zoology* 40:28–46. doi: 10.1080/03014223.2012.757242
- Burns, J. J. (2002). Harbor seal and spotted seal *Phoca vitulina* and *P. largha*. In Perrin, W.F., Wursig, B., & Thewissen J. G. M. (eds.). *Encyclopedia of marine mammals*. Academic Press. 552-560.
- British Wind Energy Association (BWEA). (2007). Investigation of Technical and Operational Effects on Marine Radar Close to Kentish Flats Offshore Wind Farm. MARICO Marine. Retrieved from [http://users.ece.utexas.edu/~ling/EU3%20UK%20Kentish%20Flats%20marine\\_radar%20study.pdf](http://users.ece.utexas.edu/~ling/EU3%20UK%20Kentish%20Flats%20marine_radar%20study.pdf)
- Cake, E.W., Jr. (1983). Habitat suitability index models: Gulf of Mexico American oyster. U.S. Fish and Wildlife Service FWS-OBS 82/10.57.
- Cape Cod Commission (CCC). (2012). *Cape Cod Regional Policy Plan*. Barnstable, MA.
- Cape Cod Commission (CCC). (2015). *Cape Cod Metropolitan Planning Organization, Cape Cod 2016 Regional Transportation Plan 2016-2040*. Barnstable, MA.
- Cargnelli, L.M., S.J. Griesbach, D.B. Packer, E. Weissberger. (1999a). Essential Fish Habitat Source Document: Atlantic SurfClam, *Spisula solidissima*, Life History and Habitat Characteristics. NOAA Technical Memorandum NMFS-NE-142. Retrieved from: <https://www.nefsc.noaa.gov/publications/tm/tm142/tm142.pdf>.
- Cargnelli, L.M., S.J. Griesbach, D.B. Packer, E. Weissberger. (1999b). Essential Fish Habitat Source Document: Ocean Quahog, *Artica islandica*, Life History and Habitat Characteristics. NOAA Technical Memorandum NMFS-NE-148. Retrieved from: <https://www.nefsc.noaa.gov/publications/tm/tm148/tm148.pdf>.
- Carter, T. C. & Feldhamer, G. A. (2005). Roost tree use by maternity colonies of Indiana bats and northern long-eared bats in southern Illinois. *Forest Ecology and Management*, 219, 259-268
- Casey, J. P., James, M. C., & Williard, A. S. (2014). Behavioral and metabolic contributions to thermoregulation in freely swimming leatherback turtles at high latitudes. *The Journal of Experimental Biology*, 217, 2331-2337.

- Casale, P., Affronte, M., Insacco, G., Freggi, D., Vallini, C., D'Astore, P.P., Basso, R., Paolillo, G., Abbate, G., & Argano, R. (2010). Sea turtle strandings reveal high anthropogenic mortality in Italian waters. *Aquatic Conservation: Marine and Freshwater Ecosystems* 20, 611-620.
- Casper, B.M., Popper, A.N., Matthews, F., Carlson, T.J., & Halvorsen, M.B. (2012). Recovery of Barotrauma Injuries in Chinook Salmon *Oncorhynchus tshawytscha* from Exposure to Pile Driving Sound. *PLoS ONE*, 7(6), e39593. Retrieved from doi:10.1371/journal.pone.0039593
- CBD (Convention on Biological Diversity). (2012). "Scientific Synthesis on the Impacts of Underwater Noise on Marine and Coastal Biodiversity and Habitats". Subsidiary Body on Scientific, Technical and technological Advice, Sixteenth meeting, Montreal.
- CCS. (2017). [Water quality monitoring data file]. Retrieved from <http://www.capecodbay-monitor.org/>
- Center for Coastal Studies. (2016). Cape Cod seals. Retrieved from <http://coastalstudies.org/seal-research/cape-cod-seals/>. Accessed November 22, 2017.
- CETAP (Cetacean and Turtle Assessment Program), University of Rhode Island. (1982). A characterization of marine mammals and turtles in the Mid- and North Atlantic areas of the US Outer Continental Shelf, final report. Contract AA551-CT8-48. Bureau of Land Management, Washington, DC. 586 pp.
- Chaloupka, M., Work, T.M., Balazs, G.H., Murakawa, S.K.K., & Morris, R. (2008). Cause-specific temporal and spatial trends in green sea turtle strandings in the Hawaii Archipelago (1982-2003). *Marine Biology* 154, 887-898.
- Chamberlain, D. E., Rehfisch, M. R., Fox, A. D., Desholm, , & Anthony, S. J. (2006). The effect of avoidance rates on bird mortality predictions made by wind turbine collision risk models. *Ibis* 148:198–202. doi: 10.1111/j.1474-919X.2006.00507.x
- Changsheng Chen, Beardsley R. C., Qi J. and Lin H. (2016). Use of Finite-Volume Modeling and the Northeast Coastal Ocean Forecast System in Offshore Wind Energy Resource Planning. Final Report to the U.S. Department of the Interior, Bureau of Ocean Energy Management, Office of Renewable Energy Programs. BOEM 2016-050. 131pp.
- Chardine, J. W. J., Rail, J.-F., & Wilhelm, S. I. (2013). Population dynamics of Northern gannets in North America, 1984–2009. *Journal of Field Ornithology* 84:187–192. doi: 10.1111/jofo.12017
- Cheesman, S. (2016). Measurements of operational wind turbine noise in UK waters, in *Effects of Noise on Aquatic Life II*. Advances in Experimental Medicine and Biology, edited by A. N. Popper and A. Hawkins. Vol. 875, 153–160. Springer, New York.

- Chen, C., Guan, S., & Chou, L. (2016). Noise field characterization in the habitat of the East Taiwan Strait Indo-Pacific humpback dolphin during the pile driving activity of demonstration offshore wind farm. *The Journal of the Acoustical Society of America*, 140, 3033.
- Cipriano, F. (2002). Atlantic white-sided dolphin. Perrin, W. F., Wursig, B., & Thewissen J. G. M. (eds.). San Diego, California: Academic Press. pp. 49-51.
- Clapham, P. J., & Mayo, C. A. (1987). Reproduction and recruitment of individually identified humpback whales, *Megaptera novaeangliae*, observed in Massachusetts Bay, 1979-1985. *Can. J. Zool.*, 65, 2853-2863.
- Clark, C. W., Ellison, W. T., Southall, B. L., Hatch, L., Van Parijs, S. M., Frankel, A., & Ponirakis, D. (2009). Acoustic masking in marine ecosystems: intuitions, analysis, and implication. *Marine Ecology Progress Series*, 395, 201-222.
- Clark, W. W. (1991). Recent studies of temporary threshold shift (TTS) and permanent threshold shift (PTS) in animals. *Journal of the Acoustical Society of America*, 90(1), 155-163.
- Cleasby, I. R., Wakefield, E. D., Bearhop, S., Bodey, T. W., Votier, S. C., & Hamer, K. C. (2015). Three-dimensional tracking of a wide-ranging marine predator: Flight heights and vulnerability to offshore wind farms. *Journal of Applied Ecology* 52:1474–1482. doi: 10.1111/1365-2664.12529
- CMACS. (2015). Ormonde Offshore Wind Farm. Year 2 Post-construction Benthic Monitoring Survey. Technical Report (2013 survey). Ref. J3264. Prepared for RPS Energy. May 2015.
- Cochran, W. W. (1985). Ocean migration of Peregrine Falcons: is the adult male pelagic? In Proceedings of Hawk Migration Conference IV (M. Harwood, Editor). Hawk Migration Association of North America, Rochester, NY, pp. 223–237.
- Colburn, L.L., Clay, P.M., Olson, J., Pinto da Silva, P., Smith, S.L., Westwood, A., & Ekstrom, J. (2010). *Community Profiles for Northeast U.S. Marine Fisheries*. Northeast Fisheries Science Center Social Sciences Branch, NOAA Fisheries. Woods Hole, MA *Community profiles available at: <https://www.nefsc.noaa.gov/read/socialsci/communityProfiles.html>*
- Colden, A. M. & Lipcius, R.N. 2015. Lethal and sublethal effects of sediment burial on the eastern oyster *Crassostrea virginica*. *Marine Ecology Progress Series* 527: 105-117.
- Conant, T. A., Dutton P. H., Eguchi T., Epperly S. P., Fahy C. C., Godfrey M. H., & Witherington, B. E. (2009). Loggerhead sea turtle (*Caretta caretta*) 2009 status review under the Endangered Species Act. Report of the Loggerhead Biological Review Team to the National Marine Fisheries Service, August 2009. 222 pp.

- Conn, P. C., & Silber, G. K. (2013). Vessel speed restrictions reduce risk of collision-related mortality for North Atlantic right whales. *Ecosphere*, 4(4), 43. Retrieved from <http://dx.doi.org/10.1890/ES13-00004.1>. Accessed November 29, 2017.
- Conservation of Endangered Species and Other Fish or Wildlife, 35 Fed. Reg. 8491 (June 2, 1970).
- Convention on Biological Diversity (CBD). (2014). Sea turtle hearing and sensitivity to acoustic impacts. Expert Workshop on Underwater Noise and its Impacts on Marine and Coastal Biodiversity. pp. 25-27. February 2014. London, United Kingdom of Great Britain and Northern Ireland.
- Cook, A. S. C. P., Johnston, A., Wright, L. J., & Burton, N. H. K. (2012). A Review of Flight Heights and Avoidance Rates of Birds in Relation to Offshore Wind Farms. *Report prepared on behalf of The Crown Estate*. [Online.] Retrieved from [http://www.bto.org/sites/default/files/u28/downloads/Projects/Final\\_Report\\_SOSS02\\_BTORview.pdf](http://www.bto.org/sites/default/files/u28/downloads/Projects/Final_Report_SOSS02_BTORview.pdf)
- Cook, S. L., & Forrest T. G. (2005). Sounds produced by nesting leatherback sea turtles (*Dermochelys coriacea*). *Herpetological Review*, 36, 387-390.
- Cooper, R. A. & Uzmann, J.R. (1980). Ecology of Juvenile and Adult Homarus. In *The Biology and Management of Lobsters*, edited by J. S. Cobb and B. F. Phillips. Volume 2, 97-142. Academic Press, New York.
- Copping, A., Sather, N., Hanna, L., Whiting, J., Zydlewski, G., Staines, G., Gill, A., Hutchison, I., O'Hagan, A., Simas, T., Bald, J., Sparling C., Wood, J., & Masden, E. (2016). Annex IV 2016 State of the Science Report: Environmental Effects of Marine Renewable Energy Development Around the World.
- Council on Environmental Quality (CEQ). (1997). *Environmental Justice Guidance Under the National Environmental Policy Act*. Executive Office of the President. Washington, DC.
- Cox, T. M., Read, A. J., Solow, A., & Tregenza, N. (2001). Will harbour porpoises (*Phocoena phocoena*) habituate to pingers? *Journal of Cetacean Research and Management*, 3, 81–86.
- CR Environmental, Inc. (2017). Memorandum: Underwater Video Review, Vineyard Wind Project, Nantucket Sound and Atlantic Ocean. Dated 15 November 2017, 21 pp.
- Cryan, P. M. (2008). Mating behavior as a possible cause of bat fatalities at wind turbines. *Journal of Wildlife Management*, 72, 845-849.
- Cryan, P. M. & Barclay, R. M. R. (2009). Causes of bat fatalities at wind turbines: Hypotheses and predictions. *Journal of Mammalogy*, 90, 1330-1340.

- Cryan, P. M. & Brown, A. C. (2007). Migration of bats past a remote island offers clues toward the problem of bat fatalities at wind turbines. *Biological Conservation*, 139, 1-11.
- Curry, B. E., & Smith, J. (1997). Phylogeographic structure of the bottlenose dolphin (*Tursiops truncatus*): Stock identification and implications for management. In Dizon, A. E., Chivers, S. J., & Perrin, W. F. (eds.). *Molecular genetics of marine mammals*. pp. 327-347. Society for Marine Mammology.
- Curtice, C., Cleart, J., Schumchenia, E., & Halpin, P. (2016). Marine-life Data and Analysis Team ("MDAT") technical report on the methods and development of marine-life data to support regional ocean planning and management. Prepared on behalf of the Marine-life Data and Analysis Team (MDAT). [Online.] Retrieved from [http://seamap.env.duke.edu/models/MDAT/MDAT-Technical-Report-v1\\_1.pdf](http://seamap.env.duke.edu/models/MDAT/MDAT-Technical-Report-v1_1.pdf)
- Czech-Damal, N., Dehnhardt, G., Manger, P., & Hanke, W. (2012). Passive electroreception in aquatic mammals. *Journal of Comparative Physiology*. doi: 10.1007/s00359-012-0780-8.
- Dadswell, M. J., Taubert, B. D., Squiers, T. S., Marchette, D., & Buckley, J. (1984). Synopsis of biological data on shortnose sturgeon *Acipenser brevirostrum*. NOAA Technical Report NMFS, 14. 45.
- Dahl, P., de Jong, C., & Popper, A. (2015). The underwater sound field from impact pile driving and its potential effects on marine life. *Acoustic Today*, 11.
- Dähne, M., Gilles, A., Lucke, K., Peschko, D., Adler, Krugel, S., ... & Siebert, U. (2013). Effect of pile-driving on harbour porpoise (*Phocoena phocoena*) at the first offshore wind farm in Germany. *Environmental Research Letters*, doi:10.1088/1748-9326/8/2/025002.
- Dames and Moore. (1993). Benthic animal-sediment assessment of potential beachfill borrow source for the Rehoboth/Dewey Beach, Delaware interim feasibility study. Report to U.S. Army Corps of Engineers, Philadelphia District. Contract No. DACW61-93-D-0001.
- Damon-Randall, K., Colligan, M., & Crocker, J. (2013). Composition of Atlantic sturgeon in rivers, estuaries, and in marine waters. White paper. NOAA/NMFS, Gloucester, Massachusetts: Protected Resources Division.
- Daoust, P. Y., Couture, E. L., Wimmer, T., & Bourque, L. (2017). Incident report: North Atlantic right whale mortality event in the Gulf of St. Lawrence, 2017. Collaborative report, produced by Canadian Wildlife Health Cooperative, Marine Animal Response Society, and Fisheries and Oceans Canada. 224 pp.
- David, J. A. (2006). Likely sensitivity of bottlenose dolphins to pile-driving noise. *Water and Environment Journal*, 20, 48–54.

- Davies, J. L. (1957). The geography of the gray seal. *Journal of Mammalian Evolution*, 38, 297-310.
- Davis, J.P., & Sisson, R.T. (1988). Aspects of the biology relating to the fisheries management of New England population of the whelks, *Busycotypus canaliculatus* and *Busycon carica*. *J. Shellfish Res.*, 7, 453-460.
- DeGraaf, R. & Yamasaki, M. (2001). *New England Wildlife-Habitat, Natural History, and Distribution*. Hanover, NH: University Press of New England.
- Delefosse, M., Rahbek, M. L., Roesen, L., & Clausen, K. T. (2017). Marine mammal sightings around oil and gas installations in the central North Sea. *Journal of the Marine Biological Association of the United Kingdom*. Retrieved from <https://www.cambridge.org/core/journals/journal-of-the-marine-biological-association-of-the-united-kingdom/article/marine-mammal-sightings-around-oil-and-gas-installations-in-the-central-north-sea/00DD61365842F95F95F10E2FDBC3337C>. Accessed November 29, 2017.
- DeLuca, W. V, Woodworth, B. K., Rimmer, C. C., Marra, P. P., Taylor, P. D., McFarland, K. P., Mackenzie, S. A., & Norris, D. R. (2015). *Transoceanic migration by a 12 g songbird*. *Biology Letters* 11.
- Department of Fisheries and Oceans. (2011). Stock assessment of Northwest Atlantic grey seals (*Halichoerus grypus*). Canadian Science Advisory Secretariat Science Advisory Report, 2010/091.
- Dernie, K. M., Kaiser, M. J., & Warwick, R. M. (2003). Recovery rates of benthic communities following physical disturbance. *Journal of Animal Ecology*, 72 (6),1043-1056.
- DeRuiter, S. L. & Doukara, K. L. (2012). Loggerhead turtles dive in response to airgun sound exposure. *Endangered Species Research*, 16, 55-63.
- Desholm, M. (2006). Wind farm related mortality among avian migrants – a remote sensing study and model analysis. PhD thesis, Department of Wildlife Ecology and Biodiversity, National Environmental Research Institute and Center for Macroecology, Institute of Biology, University of Copenhagen.
- Desholm, M., & Kahlert, J. (2005). Avian collision risk at an offshore wind farm. *Biology letters* 1:296–298. doi: 10.1098/rsbl.2005.0336
- Desorbo, C. R., Gray, R. B., Tash, J. Tash, Gray, C. E., Williams, K. A., & Rirodan, D. (2015). Offshore migration of Peregrine Falcons (*Falco peregrinus*) along the Atlantic Flyway. In *Wildlife Densities and Habitat Use Across Temporal and Spatial Scales on the Mid-Atlantic Outer Continental Shelf: Final Report to the Department of Energy EERE Wind*.

- Desorbo, C. R., Persico, C., Gray, R. B., & Gilpatrick, L. (2017). Studying migrant raptors using the Atlantic Flyway. Block Island Raptor Research Station, RI: 2016 season.
- Desorbo, C. R., Wright, K. G., & Greay, R. (2012). Bird migration stopover sites: ecology of nocturnal and diurnal raptors at Monhegan Island.
- DiGaudio, R., & Geupel, G. R. (2014). Assessing Bird and Bat Mortality at the McEvoy Ranch Wind Turbine in Marin County, California, 2009-2012. Point Blue Conservation Science.
- Dirksen, S., & van der Winden, J. (1998). Nocturnal collision risks of birds with wind turbines in tidal and semi-offshore areas. In Wind energy and landscape (C. F. Ratto and G. Solari, Editors). Balkema, Rotterdam, The Netherlands, pp. 99–108.
- Dodd, C. K. (1988). Synopsis of the biological data on the loggerhead sea turtle *Caretta caretta* (Linnaeus 1758). US Fish and Wildlife Service. Service, Biological Reports, 88(14), 110 pp.
- Dodge, K. D., Prescott, R., Lewis, D., Murley, D., & Merigo, C. (2007). A review of cold stun strandings on Cape Cod, Massachusetts from 1979-2003. In: Mast, R. B., Hutchinson, B. J., and Hutchinson, A. H. (comp.). Proceedings of the Twenty-Fourth Annual Symposium on Sea Turtle Biology and Conservation. NOAA Technical Memorandum NMFS-SEFSC-567. pp. 123.
- Dodge, K.L., Galaurdi, B., Miller, T.J., & Lutcavage, M.E. (2014). Leatherback turtle movements, dive behavior, and habitat characteristics in ecoregions of the Northwest Atlantic Ocean. PLOSone 9(3), e91726.
- Doksaeter, L., Olsen, E., Nottestad, L., & Ferno, A. (2008). Distribution and feeding ecology of dolphins along the mid-Atlantic Ridge between Iceland and the Azores. *Deep-Sea Res. II*, 55, 243-253.
- Donovan, G. P. (1991). A review of IWC stock boundaries. Reports of the International Whaling Commission (Special Issue), 13, 39-68.
- Dorr, B. S., Hatch, J. J., & Weseloh, D. V. (2014). Double-crested Cormorant (*Phalacrocorax auritus*). In The Birds of North America (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY. doi: 10.2173/bna.441
- Dow Piniak, W. E. (2012). Acoustic ecology of sea turtles: Implications for conservation. (Unpublished doctoral dissertation). Duke University, Durham, North Carolina.
- Dow Piniak, W. E., Eckert, S. A., Harms, C. A., & Stringer, E. M. (2012). Underwater hearing sensitivity of the leatherback sea turtle (*Dermochelys coriacea*): Assessing the potential effect of anthropogenic noise. US Department of the Interior, Bureau of Ocean Energy Management, Headquarters, Herndon, Virginia. OCS Study BOEM 2012-01156. 35 pp.



- Dowling, Z., Sievert, P. R., Baldwin, E., Johnson, L., Von Oettingen, S. & Reichard, J. (2017). Flight Activity and Offshore Movements of Nano-Tagged Bats on Martha's Vineyard, MA. US Department of the Interior, Bureau of Ocean Energy Management, Office of Renewable Energy Programs, Sterling, Virginia. OCS Study BOEM 2017-054. 39 pp.
- Drewitt, A. L., & Langston, R. H. W. (2006). Assessing the impacts of wind farms on birds. *Ibis* 148:29–42. doi: 10.1111/j.1474-919X.2006.00516.x
- Drury, W. H., & Keith, J. (1962). Radar studies of songbird migration in coastal New England. *Ibis* 104:449–489.
- Dufault, S., Whitehead, H., & Dillon, M. (1999). An examination of the current knowledge on the stock structure of sperm whales (*Physeter macrocephalus*) Worldwide. *J. Cetacean Res. Manage.*, 1(1), 1-10.
- Dunlop, R. A., Noad, M. J., McCauley, R. D., Scott-Hayward, L., Kniest, E., Slade, R., . . . Cato, D. H. (2017). Determining the behavioural dose–response relationship of marine mammals to air gun noise and source proximity. *Journal of Experimental Biology*, 220(16), 2878-2886. doi:10.1242/jeb.160192
- Dunton, K.J., Jordaan, A., McKown, K.A., Conover, D.O., & Frisk, M.G. (2010). Abundance and distribution of Atlantic Sturgeon *Acipenser oxyrinchus* within the northwest Atlantic Ocean, determined from five fishery independent surveys. US National Marine Fisheries Service Fishery Bulletin, 108, 450–464.
- Dürr, T. (2011). Bird loss of wind turbines in Germany: data from the central register of the National Fund Ornithological Station State Office for Environment Office, Health and Consumer Protection, Brandenburg, Germany.
- eBird. 2017. eBird: An online database of bird distribution and abundance [web application]. eBird, Ithaca, New York. Retrieved from: <http://www.ebird.org>.
- Eckert, S.A., Eckert, K.L., Ponganis, P. & Kooyman, G.L. (1989). Diving and foraging behavior of leatherback sea turtles (*Dermochelys coriacea*). *Canadian Journal of Zoology*, 67, 2834-2840.
- Edmonds, N.J., Firmin, C.J., Goldsmith, D., Faulkner, R.C., & Wood, D.T. (2016). A review of crustacean sensitivity to high amplitude underwater noise: data needs for effective risk assessment in relation to UK commercial species. *Marine Pollution Bulletin*, 108, 5–11.
- Edrén, S. M. C., Andersen, S. M., Teilmann, J., Carstensen, J., Haders, P., Dietz, R., & Miller, L. A. (2010). The effect of a large Danish offshore wind farm on harbor and gray seal haul-out behavior. *Marine Mammal Science*, 26, 614-634.

- Elliott-Smith, E. & Haig, S.M. (2004). Piping Plover (*Charadrius melodus*), version 2.0. *The Birds of North America*. P. G. Rodewald (Ed.). Cornell Lab of Ornithology, Ithaca, NY. Retrieved from <https://doi.org/10.2173/bna.2>
- Ellison, W. T., Southall, B. L., Clark, C. W., & Frankel, A. S. (2012). A new context-based approach to assess marine mammal behavioral responses to anthropogenic sounds. *Conservation Biology*, 26(1), 21–28.
- Endangered and Threatened Species, Determination of Nine Distinct Population Segments of Loggerhead Sea Turtles as Endangered or Threatened, 76 Fed. Reg. 58868 (September 22, 2011).
- Endangered and Threatened Species; Critical Habitat for Endangered North Atlantic Right Whale, 81 Fed. Reg. 4837 (January 27, 2016) (codified at 50 C.F.R. 26).
- Endangered and Threatened Species; Identification of 14 Distinct Population Segments of the Humpback Whale (*Megaptera novaeangliae*) and Revision of Species-Wide Listing, 81 Fed. Reg. 62259 (September 8, 2016).
- English, P.A., Mason, T.I., Backstrom, J.T., Tibbles, B.J., Mackay, A.A., Smith, M.J. and Mitchell, T. (2017). Improving Efficiencies of National Environmental Policy Act Documentation for Offshore Wind Facilities Case Studies Report. US Dept. of the Interior, Bureau of Ocean Energy Management, Office of Renewable Energy Programs, Sterling. OCS Study BOEM 2017-026. 217pp.
- Environmental Protection Agency (EPA). (2017). *Environmental Justice*. Retrieved from <https://www.epa.gov/environmentaljustice/ej-2020-glossary>
- EPA. (2015). National Coastal Condition Assessment 2010. Office of Water and Office of Research and Development. (EPA 841-R-15-006). Washington, DC. December 2015.
- Erbe, C. (2002). Underwater noise of whale-watching boats and potential effects on killer whales (*Orcinus orca*), based on an acoustic impact model. *Marine Mammal Science*, 18(2), 394-418.
- Erbe, C. (2009). Underwater noise from pile driving in Moreton Bay. Australian Acoustical Society Queensland, 37(3), 87–92.
- Erfteimeijer, P.L.A. & Shuail, D.A. (2012). Seagrass habitats in the Arabian Gulf: distribution, tolerance thresholds and threats. *Aquatic Ecosystem Health & Management*, 15(sup1), 73-83.

- Erickson, W. P., Jeffrey, J. D., & Poulton, V. K. Poulton (2008). Puget Sound Energy Wild Horse Wind Facility Post-Construction Avian and Bat Monitoring. First Annual Report. January - December 2007. *A report prepared for Puget Sound Energy, Ellensburg, Washington and the Wild Horse Wind Facility Technical Advisory Committee, Kittitas County, Washington.* [Online.] Retrieved from [http://www.efsec.wa.gov/wildhorse/SupplementalEIS/DSEIS/WILDHORSE Wind Project \\_ First Annual Monitoring Report\\_012808.pdf](http://www.efsec.wa.gov/wildhorse/SupplementalEIS/DSEIS/WILDHORSEWindProject_FirstAnnualMonitoringReport_012808.pdf)
- Erickson, W. P., Wolfe, M. M., Bay, K. J., Johnson, D. H., & Gehring, J. L. (2014). A Comprehensive Analysis of Small-Passerine Fatalities from Collision with Turbines at Wind Energy Facilities. *PLoS ONE* 9:18. doi: 10.1371/journal.pone.0107491
- Essink, K. (1999). Ecological effects of dumping of dredged sediments; options for management. *Journal of Coastal Conservation* 5, 69–80.
- ESS Group. (2006). Potential Impacts to Predator-Prey Relationships as a Result of the Proposed Project in Nantucket Sound. Appendix 3.9-B. Prepared for Cape Wind Associates, LLC. ESS Project No. E159-503.8.
- ESS Group, Inc. (2017). Benthic Macroinvertebrate Sample Analysis Report. Vineyard Wind Project Offshore Lease Area OCS-A 0501 Massachusetts. Prepared for Vineyard Wind, LLC. Project No. O207-000.
- European Union (EU) Commission. (2016). MaRVEN - Environmental impacts of noise, vibrations, and electromagnetic emissions from marine renewable energy. European Union, Final study report RTD-KI-NA-27-738-EN-N.
- Evans, N.T., Ford, K.H., & Chase, B.C., and Sheppard, J.J. (2015). Recommended Time of Year Restrictions (TOYs) for Coastal Alternation Projects to Protect Marine Fisheries Resources in Massachusetts. Retrieved from <https://www3.epa.gov/region1/superfund/sites/newbedford/548591.pdf>
- Everaert, J., & Stienen, E. W. M. (2007). Impact of wind turbines on birds in Zeebrugge (Belgium): Significant effect on breeding tern colony due to collisions. *Biodiversity and Conservation* 16:3345–3359. doi: 10.1007/s10531-006-9082-1
- Exec. Order No. 12898, 3 C.F.R. (1994).
- Executive Office of Labor and Workforce Development (EOLWD). (2017). *Labor Market Information, Largest Employers by Area*. Available at: [http://lmi2.detma.org/lmi/Largest\\_employer\\_index.asp](http://lmi2.detma.org/lmi/Largest_employer_index.asp)
- Faaborg, J., Holmes, R. T., Anders, A. D., Bildsteing, K. L., Dugger, K. M., Gauthreaux, S. A., Heglun, P., Hobson, K. A. Hobson, Jahn, A. E., Johnson, D. H., & Latta, S. C. (2010). Recent advances in understanding migration systems of New World land birds. *Ecological Monographs* 80:3–48. doi: 10.1890/09-0395.1

- Fabricius, K.E. (2005). Effects of terrestrial runoff on the ecology of corals and coral reefs: review and synthesis. *Marine Pollution Bulletin* 50:125-146.
- Ferrara, C. R., Vogt, R. C., Harfush, M. R., Sousa-Lima, R. S., Albavera, E., Tavera, A. (2014). First evidence of leatherback turtle (*Dermochelys coriacea*) embryos and hatchlings emitting sounds. *Chelonian Conservation and Biology*, 13(1), 110-114.
- FFWCC (Florida Fish and Wildlife Conservation Commission). (2017). Index nesting beach survey totals (1989-2017). Retrieved from <http://myfwc.com/research/wildlife/sea-turtles/nesting/beach-survey-totals/>. Accessed on November 22, 2017.
- Filadelfo, R., Mintz, J., Michlovich, E., D'Amico, A., & Tyack, P. L. (2009). Correlating military sonar use with beaked whale mass strandings: What do the historical data show? *Aquatic Mammals*, 34, 435–444.
- Finneran, J. J. (2016). Auditory weighting functions and TTS/PTS exposure functions for Cetaceans and marine carnivores. SPAWAR Systems Center Pacific, San Diego, California. May 2016.
- Fox, A. D., Desholm, M., Kahlert, J., Christensen, T. K., & Petersen, I. K. (2006). Information needs to support environmental impact assessment of the effects of European marine offshore wind farms on birds. *Ibis* 148:129–144. doi: 10.1111/j.1474-919X.2006.00510.x
- Fraser, D. (2017). Seabirds Washing Up and Dying on Cape Cod Beaches. Cape Cod Times.
- Frederick, P. C. (2001). Birds in the Marine Environment. In *Biology of Marine Birds* (E. A. Schreiber and J. Burger, Editors). CRC Press, Boca Raton, FL., pp. 617–655.
- Friedlander, A.M., Ballesteros, E., Fay, M., & Sala, E. (2014). Marine communities on oil platforms in Gabon, West Africa: High biodiversity oases in a low biodiversity environment. *PLoSone*, 9(8), e103709.
- Freiwald, A., Fossá, J.H., Grehan, A., Koslow, T., Roberts, J.M. (2004). *Cold-water Coral Reefs*. UNEP-WCMC, Cambridge, UK.
- Furness, R. W., Wade, H. M., & Masden, E. A. (2013). Assessing vulnerability of marine bird populations to offshore wind farms. *Journal of Environmental Management* 119:56–66. doi: 10.1016/j.jenvman.2013.01.025
- Gallaway, B.J., Szedlmayer, S.T., & Gazey, W.J. (2012). A life history review for red snapper in the Gulf of Mexico with an evaluation of the importance of offshore petroleum platforms and other artificial reefs. *Reviews in Fisheries Science*, 17, 48-67.

- Garthe, S., Guse, N., Montevecchi, W. A., Rail, J.-F., & Grégoire, F. (2014). The daily catch: Flight altitude and diving behavior of northern gannets feeding on Atlantic mackerel. *Journal of Sea Research* 85:456–462.
- Garthe, S., & Hüppop, O. (2004). Scaling possible adverse effects of marine wind farms on seabirds: developing and applying a vulnerability index. *Journal of Applied Ecology* 41:724–734. doi: 10.1111/j.0021-8901.2004.00918.x
- Garthe, S., Markones, N., & Corman, A.-M. (2017). Possible impacts of offshore wind farms on seabirds: a pilot study in Northern Gannets in the southern North Sea. *Journal of Ornithology* 158:345–349. doi: 10.1007/s10336-016-1402-y
- Gaston, A., & Jones, I.L (1998). The Auks. In. Oxford University Press, Oxford, UK.
- Gates A. R., & Jones, D. O. B. (2012). Recovery of benthic megafauna from anthropogenic disturbance at a hydrocarbon drilling well (380 m depth in the Norwegian Sea). *PLoS ONE*, 7, e44114.
- Gauthreaux, S. A., & Belser, C.G. (1999). Bird migration in the region of the Gulf of Mexico. In Proceedings of the 22nd International Ornithological Congress (N. J. Adams and R. H. Slotow, Editors). BirdLife South Africa, Durban, Johannesburg, South Africa, pp. 1931–1947.
- Gedamke, J., Harrison, J., Hatch, L., Angliss, R., Barlow, J., Berchok, C., Caldow, C., Castellote, M., Cholewiak, D., Deangelis, M. L., Dziak, R., Garland, E., Guan, S., Hastings, S., Holt, M., Laws, B., Mellinger, D., Moore, S., Moore, T. J., Oleson, E., Pearson-Meyer, J., Piniak, W., Redfern, J., Rowles, T., Scholik-Schlomer, A., Smith, A., Soldevilla, M., Stadler, J., Parijs, S., & Van Wahle, C. (2016). Ocean Noise Strategy Roadmap. National Oceanic and Atmospheric Administration. Retrieved from [https://cetsound.noaa.gov/Assets/cetsound/documents/Roadmap/ONS\\_Roadmap\\_Final\\_Complete.pdf](https://cetsound.noaa.gov/Assets/cetsound/documents/Roadmap/ONS_Roadmap_Final_Complete.pdf).
- Gill A., Gloyne-Philips, I., Kimber, J., & Sigray, P. (2014). Marine renewable energy, electromagnetic (EM) fields and EM-sensitive animals. In Shields, M., & Payne, A. (eds.). *Humanity and the sea: Marine renewable energy and the interactions with the environment*.
- Gill, A. B. & Kimber, A. A. (2005). The potential for cooperative management of elasmobranchs and offshore renewable energy development in UK waters. *Journal of the Marine Biological Association of the United Kingdom* 85:1075-1081.
- Gill, A. B., Gloyne-Phillips, I., Neal, K. J., & Kimber, J. A. (2005). The potential effects of electromagnetic fields generated by sub-sea power cables associated with offshore wind farm developments on electrically and magnetically sensitive marine organisms - a review. Collaborative Offshore Wind Research into the Environment (COWRIE), Ltd, UK. 128 pp.

- Gilmour, J. (1999). Experimental investigation into the effects of suspended sediment on fertilization, larval survival, and settlement in a scleractinian coral. *Marine Biology* 135:451-462.
- Gitschlag, G.R., Herczeg, B.A., & Barcak, T.R. (1997). Observations of sea turtles and other marine life at the explosive removal of offshore oil and gas structures in the Gulf of Mexico. *Gulf Research Reports*, 9(4), 247-262.
- Glasby, T.M., Connell, S.D., Holloway, M.G., & Hewitt, C.L. (2007). Nonindigenous biota on artificial structures: could habitat creation facilitate biological invasions? *Marine Biology*, 151: 887–895.
- Good, T. P. (1998). Great Black-backed Gull (*Larus marinus*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY. doi: 10.2173/bna.330
- Goodale, M. W., & Milman, A. (2016). Cumulative adverse effects of offshore wind energy development on wildlife. *Journal of Environmental Planning and Management* 59:1–21. doi: 10.1080/09640568.2014.973483
- Goodale, M. W., & Stenhouse, I.J. (2016). A conceptual model for determining the vulnerability of wildlife populations to offshore wind energy development. *Human-Wildlife Interactions* 10:53–61.
- Gowans, S., & Whitehead, H. (1995). Distribution and habitat partitioning by small odontocetes in the Gully, a submarine canyon on the Scotian Shelf. *Canadian Journal of Zoology*, 73, 1599-1608.
- Goyert, H. F., Manne, L.L., & Veit, R.R. (2014). Facilitative interactions among the pelagic community of temperate migratory terns, tunas and dolphins. *Oikos* 123:1400–1408. doi: 10.1111/oik.00814
- Gradient Corporation. (2017). Electric and Magnetic Field (EMF) Modeling Analysis for the Vineyard Wind Connector Project. Prepared for Epsilon Associates, Inc. and Vineyard Wind LLC.
- Grady, F. V. & Olson, S. L. (2006). Fossil bats from quaternary deposits on Bermuda (chiroptera: vespertilionidae). *Journal of Mammalogy*, 87, 148-152.
- Gray, C. E., Gilbert, A.T., Stenhouse, I., & Berlin, A.M. (2017). Occurrence Patterns and Migratory Pathways of Red-throated Loons Wintering in the Offshore Mid-Atlantic US, 2012-2016. In *Determining Fine-scale Use and Movement Patterns of Diving Bird Species in Federal Waters of the Mid-Atlantic United States Using Satellite Telemetry* (C. S. Spiegel, Editor). US. Department of the Interior, Bureau of Ocean Energy Management, Division of Environmental Sciences, Sterling, VA.

- Greater Atlantic Fisheries Office (GARFO). (2016). Recovery outline for the Atlantic Sturgeon Distinct Population Segments. Report for NOAA Fisheries Protected Resources. Received from [https://www.greateratlantic.fisheries.noaa.gov/protected/atlsturgeon/docs/ats\\_recovery\\_outline\\_gar\\_ser\\_final\\_508\\_compliant\\_v2.pdf](https://www.greateratlantic.fisheries.noaa.gov/protected/atlsturgeon/docs/ats_recovery_outline_gar_ser_final_508_compliant_v2.pdf).
- Griffin, L., Rees, E., & Hughes, B. (2011). Migration routes of whooper swans and geese in relation to wind project footprints. Final Report to the UK Department of Energy and Climate Change (DECC), London. The Wildfowl and Wetlands Trust, Slimbridge, Gloucestershire, England, United Kingdom.
- Guida, V., A. Drohan, H. Welch, J. McHenry, D. Johnson, V. Kentner, J. Brink, D. Timmons, E. Estela-Gomez. (2017). Habitat Mapping and Assessment of Northeast Wind Energy Areas. Sterling, VA: US Department of the Interior, Bureau of Ocean Energy Management. OCS Study BOEM 2017-088. p.312.
- Hain, J. H. W., Ratnaswamy, M. J., Kenney, R. D., & Winn, H. E. (1992). The fin whale, *Balaenoptera physalus*, in waters of the northeastern United States continental shelf. Reports of the International Whaling Commission, 42, 653-669.
- Halvorsen M. B., Casper, B. M., Woodley, C. M., Carlson, T. J., & Popper, A.N. (2011). Predicting and mitigating hydroacoustic impacts on fish from pile installations. NCHRP Res Results Digest 363, Project 25–28, National Cooperative Highway Research Program, Transportation Research Board, National Academy of Sciences, Washington, D.C.
- Halvorsen, M. B., Casper, B. M., Woodley, C. M., Carlson, T. J., & Popper, A. N. (2012). Threshold for onset of injury in Chinook salmon from exposure to impulsive pile driving sounds. PLoS ONE 7(6), e38968. doi: 10.1371/journal.pone.0038968.
- Halvorsen, M. B., Casper, B. M., Matthews, F., Carlson, T. J., & Popper, A. N. (2012). Effects of exposure to pile driving sounds on the lake sturgeon, Nile tilapia, and hogchoker. Proceeding of the Royal Society, 279, 4705-4714.
- Hamazaki, T. (2002). Spatiotemporal prediction models of cetacean habitats in the mid-western North Atlantic Ocean (from Cape Hatteras, North Carolina, USA to Nova Scotia, Canada). *Marine Mammal Science*, 18(4), 920-939.
- Handegard, N.O., Michalsen, K., & Tjostheim, D. (2003). Avoidance behavior in cod, *Gadus morhua*, to a bottom trawling vessel. *Aquatic Living Resources*, 16, 265–270.

- Harris, D. E., Lelli, B., & Jakush, G. (2002). Harp seal records from the southern Gulf of Maine: 1997-2001. *Northeast. Nat.*, 9(3), 331-340.
- Hart, D. (2006). Atlantic sea scallop (*Placopecten magellanicus*). Status of Fishery Resources off the Northeastern US, NEFSC- Resource Evaluation and Assessment Division. Received from <http://www.nefsc.noaa.gov/sos/spsyn/iv/scallop/>
- Hartman, J. C., Krijgsveld, K.L., Poot, M.J.M., Fijn, R.C., Leopold, M.F., & Dirksen, S. (2012). Effects on birds of Offshore Wind farm Egmond aan Zee (OWEZ).
- Hatch, J. J., Brown, K.M., Hogan, G.G., & Morris, R.D. (2000). Great cormorant(*Phalacrocorax carbo*). *The Birds of North America*: 32.
- Hatch, S. K., Connelly, E.E., Divoll, T.J., Stenhouse, I.J., & Williams, K.A. (2013). Offshore observations of eastern red bats (*Lasiurus borealis*) in the mid-Atlantic United States using multiple survey methods. *PLoS ONE* 8:1–8. doi: 10.1371/journal.pone.0083803
- Hawkes, L.A., Broderick, A.C., Coyne, M.S., Godfrey, M.H., & Godley, B.J. (2007). Only some like it hot – quantifying the environmental niche of the loggerhead sea turtle. *Diversity and Distribution*, 13, 447-457.
- Hays, G.C., Adams, C.R., Broderick, A.C., Godley, B.J., Lucas, D.J., Metcalfe, J.D., & Prior, A.A. (2000). The diving behavior of green turtles at Ascension Island. *Animal Behaviour*, 59, 577-586.
- Hayes, M. A. (2013). Bats Killed in Large Numbers at United States Wind Energy Facilities. *BioScience*, 975.
- Hayes S. A., Josephson, E., Maze-Foley, K., & Rosel, P. E. (eds.). (2017). US Atlantic and Gulf of Mexico marine mammal stock assessments – 2016. NOAA Tech Memo NMFS NE 241.
- Hazel, J., Lawler, I.R., Marsh, H., & Robson, S. (2007). Vessel speed increases collision risk for the green turtle *Chelonia mydas*. *Endangered Species Research*, 3, 105-113.
- Hein, C. D., Prichard, A., Mabee, T., & Schirmacher, M.R. (2013). Avian and Bat Post-construction Monitoring at the Pinnacle Wind Farm, Mineral County, West Virginia: 2012 Final Report. *An annual report submitted to the Bats and Wind Energy Cooperative*.
- Henderson, I. G., Langston, R. H. W., & Clark, N. A. (1996). The response of common terns *Sterna hirundo* to power lines: An assessment of risk in relation to breeding commitment, age and wind speed. *Biological Conservation* 77:185–192.



- Hendrick, V. J., Hutchison, Z. L., & Last, K. S. (2016). Sediment Burial Intolerance of Marine Macroinvertebrates. *PLoS One*, 11(2) e0149114.
- Heron Conservation (2017). Herons of the World. [Online.] Retrieved from <http://www.heronconservation.org/herons-of-the-world/>
- Hersh, S. L., & Duffield, D. A. (1990). Distinction between northwest Atlantic offshore and coastal bottlenose dolphins based on hemoglobin profile and morphometry. In Leatherwood, S., & Reeves, R. R. (eds.). *The bottlenose dolphin*. 653 pp. San Diego, California: Academic Press.
- Hildebrand, J. A. (2005). Impacts of anthropogenic sound. In *Marine mammal research: Conservation beyond crisis*. Reynolds III, J. E., Perrin, W. F., Reeves, R. R., Montgomery, S., & Ragen, T. J. (eds.). Johns Hopkins University Press. Baltimore, Maryland. pp. 101-124.
- Hill, R., Hill, K., Aumuller, R., Schulz, A., Dittmann, T., Kulemeyer, C., & Coppack, T. (2014). Of birds, blades, and barriers: Detecting and analysing mass migration events at alpha ventus. In *Ecological Research at the Offshore Windfarm alpha ventus - Challenges, Results, and Perspectives* (Federal Maritime and Hydrographic Agency (BSH), Federal Ministry of the Environment Nature Conservation and Nuclear Safety (BMU), A. Beiersdorf and K. Wollny-Goerke, Editors). Springer Spektrum, Hamburg and Berlin, Germany, pp. 111–132. doi: 10.1007/978-3-658-02462-8
- Hille Ris Lambers, R. & ter Hofstede, R. (2009). Refugium Effects of the MEP NSW Windpark on Fish: Progress Report 2007. s.l.: In *Interim report demersal fish: IMARES Institute for Marine Resources & Ecosystem Studies*.
- Hooper, T., Ashley, M., & Austen, M. (2015). Perceptions of fishers and developers on the co-location of offshore wind farms and decapod fisheries in the UK. *Marine Policy*, 61, 16-22.
- Horn, J. W., Arnett, E.B. tt, & Kunz, T.H. (2008). Behavioral responses of bats to operating wind turbines. *Journal of Wildlife Management* 72, 123-132.
- Hötker, H., Thomsen, K., & Jeromin, H. (2006). Impacts on biodiversity of exploitation of renewable energy sources: the example of birds and bats - facts, gaps in knowledge, demands for further research, and ornithological guidelines for the development of renewable energy exploitation.
- Hourigan, T. F., P. J. Etnoyer, R. P. McGuinn, C. Whitmire, D.S. Dorfman, M. Dornback, S. Cross, D. Sallis. 2015. An Introduction to NOAA's National Database for Deep-Sea Corals and Sponges. NOAA Technical Memorandum NOS NCCOS 191. 27 pp. Silver Spring, MD. Available online at: [https://data.nodc.noaa.gov/coris/library/NOAA/CRCP/other/other\\_crpc\\_publications/DeepSeaCoralRT/Intro\\_Natl\\_DB\\_for\\_DSCS.pdf](https://data.nodc.noaa.gov/coris/library/NOAA/CRCP/other/other_crpc_publications/DeepSeaCoralRT/Intro_Natl_DB_for_DSCS.pdf).

- Howes, B. L., Schlezinger, D. R., Blake, J. A., & Rhoads, D. C. (1997). Infaunal “recovery” as a control of sediment organic matter remineralization and the fate of regenerated nutrients. In: Boston Harbor: 14th Biennial Estuarine Research Federation (ERF) International Conference - The State of Our Estuaries, Providence, R.I. October 12-16, Abstracts.
- Howes, B.L., Samimy, R., Schlezinger, D., Bartlett, M, Benson, J. 2011. Environmental Effects of Sediment Transport Alteration and Impacts on Protected Species: Edgartown Tidal Project; Topic Area II: Marine Science and Hydrokinetic Site Specific Environmental Studies. School for Marine Science and Technology (SMAST), Coastal Systems Program. University of Massachusetts Dartmouth, 98 pp.
- Hüppop, O., Dierschke, J., Exo, K.-M., Fredrich, E., & Hill, R. (2006). Bird migration studies and potential collision risk with offshore wind turbines. *Ibis* 148:90–109. doi: 10.1111/j.1474-919X.2006.00536.x
- Huppopp, O., & Hilgerloh, G. (2012). Flight call rates of migrating thrushes: effects of wind conditions, humidity and time of day at an illuminated offshore platform. *Journal of Avian Biology*:85.
- Hutchison, Z. L., Sigray, P., He, H., Gill, A. B., King, J., and Gibson, C. (2018). Electromagnetic Field (EMF) Impacts on Elasmobranch (shark, rays, and skates) and American Lobster Movement and Migration from Direct Current Cables. U.S. Department of the Interior, Bureau of Ocean Energy Management, Sterling VA. OCS Study BOEM 2018-003.
- Illingworth & Rodkin, Inc. (2007). Compendium of pile driving sound data. Report for the California Department of Transportation. Petaluma, California.
- Imber, M. J. (1975). Behaviour of petrels in relation to the Moon and artificial lights. *Notornis* 22:302–306.
- Inger, R., Attrill, M. J., Bearhop, S., Broderick, A. C., Grecian, W. J., Hodgson, D. J., ... & Godley, B. J. (2009). Marine renewable energy: Potential benefits to biodiversity? An urgent call for research. *Journal of Applied Ecology*, 46, 1145-1153.
- International Finance Corporation. (2007). “Environmental, Health, and Safety Guidelines – Wind Energy.” Retrieved from [http://www.ifc.org/ifcext/sustainability.nsf/AttachmentsByTitle/gui\\_EHSGuidelines2007\\_WindEnergy/\\$FILE/Final+-+Wind+Energy.pdf](http://www.ifc.org/ifcext/sustainability.nsf/AttachmentsByTitle/gui_EHSGuidelines2007_WindEnergy/$FILE/Final+-+Wind+Energy.pdf)
- International: IMO Marine Engine Regulations. (2016). Retrieved from <https://www.dieselnets.com/standards/inter/imo.php>

- International Union for the Conservation of Nature. (2010). 2010 IUCN red list of threatened species. Retrieved from <http://www.iucnredlist.org>. Accessed November 22, 2017.
- ISO New England. (2017). 2015 ISO New England electric generator air emissions report. Retrieved from [https://www.iso-ne.com/static-assets/documents/2017/01/2015\\_emissions\\_report.pdf](https://www.iso-ne.com/static-assets/documents/2017/01/2015_emissions_report.pdf)
- Jacobs, S. R., & Terhune, J. M. (2000). Harbor seal (*Phoca vitulina*) numbers along the New Brunswick coast of the Bay of Fundy in the fall in relation to aquaculture. *Northeastern Naturalist*, 7(3), 289-296.
- Jacobson, L. D. (2005). Essential fish habitat source document: Longfin inshore squid *Loligo pealeii* life history and habitat characteristics (2nd edition). NOAA Tech Memo NMFS NE 193; 42 pp.
- James, E., Smith, K., Gallien, D. R., & Khan, A. (2017). Observing cable laying and particle settlement during the construction of the Block Island wind farm. Final report to the US Department of the Interior, Bureau of Ocean Energy Management, Office of Renewable Energy Programs. OCS Study BOEM 2017-027. 225 pp.
- Jefferson, T. A., Webber, M. A., & Pitman, R. L. (2008). Marine mammals of the world: A comprehensive guide to their identification. Amsterdam: Elsevier.
- Jefferson, T. A., Weir, C. R., Anderson, R. C., Balance, L. T., Barlow, J., Kenney, R. D., & Kiszka, J. J. (2014). Global distribution of Risso's dolphin *Grampus griseus*: A review and critical evaluation. *Mamm. Rev.*, 44, 56–68.
- Jelinski, D. E., Krueger, C. C., & Duffus, D. A. (2002). Geostatistical analyses of interactions between killer whales (*Orcinus orca*) and recreational whale-watching boats. *Applied Geography*, 22(4), 393-411.
- Jennings, S., Kaiser, M., & Reynolds, J.D. (2013). *Marine Fisheries Ecology*. John Wiley & Sons. 432 pages. Retrieved from <https://books.google.com/books?id=DhpIDAAQBAJ&pg=PT522&lpg=PT522&dq=benthic+community+within+sand+waves&source=bl&ots=uOV-U5NaYf&sig=n-VuZSKHE5OMn98aPk5Z20QVwoY&hl=en&sa=X&ved=0ahUKEwjRi4PQ6dWAhVFFMAKHSv8Ag0Q6AEIRTAH#v=onepage&q=benthic%20community%20within%20sand%20waves&f=false>
- Jensen, F., Laczny, M., Piper, W., & Coppack, T. (2014). Horns Rev 3 Offshore Wind Farm - Migratory Birds. [Online.] Retrieved from <http://www.4coffshore.com/windfarms/horns-rev-1-denmark-dk03.html>

- Johnson, G. D., Perlik, M. K., Erickson, W. P., & Strickland, M. D. (2004). Bat activity, composition, and collision mortality at a large wind plant in Minnesota. *Wildlife Society Bulletin*, 32, 1278-1288.
- Johnson, J. A., Storrer, J. Fahy, K. & Reitherman, B. (2011). Determining the potential effects of artificial lighting from Pacific Outer Continental Shelf (POCS) region oil and gas facilities on migrating birds. Prepared by Applied Marine Sciences, Inc. and Storrer Environmental Services for the US. Department of the Interior, Bureau of Ocean Energy Management, Regulations and Enforcement. Camarillo, CA. OCS Study BOEMRE 2011-047:29 pp.
- Johnson, J. B., Gates, J. E. & Zegre, N. P. (2011). Monitoring seasonal bat activity on a coastal barrier island in Maryland, USA. *Environmental Monitoring and Assessment*, 173, 1-4.
- Johnson, S., Connelly, E., Williams, K., Adams, E., Stenhouse, I., & Gilbert, A. (2015). Integrating data across survey methods to identify spatial and temporal patterns in wildlife distributions. *Wildlife Densities and Habitat Use Across Temporal and Spatial Scales on the Mid-Atlantic Continental Shelf: Final Report to the Department of Energy EERE Wind & Water Power Program*. [Online.] Retrieved from [http://www.briloon.org/uploads/BRI\\_Documents/Wildlife\\_and\\_Renewable\\_Energy/MABS Project Chapter 17 - Johnson et al 2015.pdf](http://www.briloon.org/uploads/BRI_Documents/Wildlife_and_Renewable_Energy/MABS_Project_Chapter_17_-_Johnson_et_al_2015.pdf)
- Johnston, A., Cook, A. S. C. P., Wright, L. J., Humphreys, E. M., & Burton, N. H. K. (2014). Modelling flight heights of marine birds to more accurately assess collision risk with offshore wind turbines. *Journal of Applied Ecology* 51:31–41. doi: 10.1111/1365-2664.12191
- Jongbloed, R. H. (2016). Flight height of seabirds. A literature study.
- Kahlert, I., Fox, A., Desholm, M., Clausager, I., & Petersen, J. (2004). Investigations of Birds During Construction and Operation of Nysted Offshore Wind Farm at Rødsand. Report by National Environmental Research Institute (NERI). pp 88.
- Kahnle, A. W., Hattala, K. A. & McKown, K. A. (2007). Status of Atlantic sturgeon in the Hudson River Estuary, New York, USA. In *Anadromous Sturgeons: Habitats, Threats, and Management* (Munro, J., Hatin, D., Hightower, J. E., McKown, K., Sulak, K. J., Kahnle, A. W. & Caron, F., eds), pp. 347–363. American Fisheries Society Symposium 56.
- Kastak, D. and R. Schusterman. (1999). In-air and underwater hearing sensitivity of a northern elephant seal (*Mirounga angustirostris*). *Canadian Journal of Zoology*, 77: 1751-1758.
- Kastelein, R. A., Gransier, R., & Jennings, N. (2013). Hearing thresholds of two harbour seals (*Phoca vitulina*) for playbacks of multiple pile driving strike sounds. *The Journal of the Acoustical Society of America*, 134(3), 2307–2312.

- Kastelein, R., Gransier, R., Marijt, M., & Hoek, L. (2015). Hearing frequency thresholds of harbor porpoises (*Phocoena phocoena*) temporarily affected by played back offshore pile driving sounds. *The Journal of the Acoustical Society of America*, 137, 556-564.
- Kastelein, R., Helder-Hoek, L., Covi, J., & Gransier, R. (2016). Pile driving playback sounds and temporary threshold shift in harbor porpoises (*Phocoena phocoena*): Effect of exposure duration. *Journal of the Acoustical Society of America*, 139, 2842-2851.
- Kastelein, R. P. Wensveen, L. Hoek, and J. Terhune. (2009). Underwater hearing sensitivity of harbor seals (*Phoca vitulina*) for narrow noise bands between 0.2 and 80 kHz. *Journal of the Acoustical Society of America*, 126: 476-483.
- Katona, S. K., & Beard, J. A. (1990). Population size, migrations, and feeding aggregations of the humpback whale (*Megaptera novaeangliae*) in the western North Atlantic Ocean. *Rep. Int. Whal. Comm. (Special Issue)*, 12, 295-306.
- Katona, S. K., Rough, V., & Richardson, D. T. (1993). A field guide to whales, porpoises, and seals from Cape Cod to Newfoundland. Washington, DC: Smithsonian Institution Press.
- Katzner, T., Smith, B.W., Miller, T.A., Brandes, D., Cooper, J., Lanzone, M. Brauning, D., Farmer, C., Harding, S., Kramar, D.E., Koppie, C., et al. (2012). Status, biology, and conservation priorities for North America's eastern Golden Eagle (*Aquila chrysaetos*) population. *Auk* 129:168–176. doi: 10.1525/auk.2011.11078
- Kenney, M. K. (1994). Harbor seal population trends and habitat use in Maine. University of Maine.
- Kenney, R. D. (1990). Bottlenose dolphins off the northeastern United States. pp. 369-386. In Leatherwood, S., & Reeves, R. R. (eds.). *The bottlenose dolphin*. 653 pp. San Diego, California: Academic Press.
- Kenney, R. D., Hyman, M. A. M., Owen, R. E., Scott, G. P., & Winn, H. E. (1986). Estimation of prey densities required by western North Atlantic right whales. *Marine Mammal Science*, 2, 1-13.
- Kenney, R. D., Scott, G. P., Thompson, T. J., & Winn, H. E. (1997). Estimates of prey consumption and trophic impacts of cetaceans in the USA northeast continental shelf ecosystem. *Journal of Northwest Atlantic Fishery Science*, 22, 155-171.
- Kenney, R. D. & Vigness-Raposa, K. J. (2010). RICRMC (Rhode Island Coastal Resources Management Council) Ocean Special Area Management Plan (SAMP), Volume 2. Appendix, Chapter 10. Marine Mammals and Sea Turtles of Narragansett Bay, Block Island Sound, Rhode Island Sound, and Nearby Waters: An Analysis of Existing Data for the Rhode Island Ocean Special Area Management Plan. 337 pp.

- Kenney, R. D., Winn, H. E., & Macaulay, M. C. (1995). Cetaceans in the Great South Channel, 1979-1989: Right whale (*Eubalaena glacialis*). *Continental Shelf Research*, 15, 385-414.
- Kerlinger, P. (1985). Water-crossing behavior of raptors during migration. *Wilson Bulletin* 97:109–113.
- Kirkpatrick, J.A., Benjamin, S., DePiper, G.D., Murphy, T., Steinback, S., & Demarest, C. (2017). *Socio-Economic Impact of Outer Continental Shelf Wind Energy Development on Fisheries in the U.S. Atlantic, Vol. I – Report Narrative*. U.S. Dept. of the Interior, Bureau of Ocean Energy Management, Atlantic OCS Region. Washington, D.C. OCS Study BOEM 2017-012.
- Kirschvink, J. L. (1990). Geomagnetic sensitivity in cetaceans: An update with live stranding records in the United States. In Thomas, J. A., & Kastelein, R. A. (eds.). *Sensory abilities of cetaceans: Laboratory and field evidence*. pp. 639-649. New York, New York: Plenum Press.
- Kirschvink, J. L., Dizon, A. E., & Westphal, J. A. (1986). Evidence from strandings of geomagnetic sensitive cetaceans. *Journal of Experimental Biology*, 120, 1-24.
- Kinlan, B., Poti, M., Dorfman, D., Caldow, C., Drohan, A., Packer, D., and Nizinski, M. (2016). Model output for deep-sea coral habitat suitability in the U.S. North and Mid-Atlantic from 2013 (NCEI Accession 0145923). NOAA National Centers for Environmental Information. Dataset. [Online]. Retrieved from [https://coastalscience.noaa.gov/data\\_reports/model-output-for-deep-sea-coral-habitat-suitability-in-the-u-s-north-and-mid-atlantic-from-2013-ncei-accession-0145923/](https://coastalscience.noaa.gov/data_reports/model-output-for-deep-sea-coral-habitat-suitability-in-the-u-s-north-and-mid-atlantic-from-2013-ncei-accession-0145923/).
- Kinlan, B. P., Winship, A.J., White, T.P., & Christensen, J. (2016). Modeling At-Sea Occurrence and Abundance of Marine Birds to Support Atlantic Marine Renewable Energy Planning Phase I Report. *Phase I Report*.
- Klima, E. F., Gitschlag, G. R., & Renaud, M. L. (1988). Impacts of the explosive removal of offshore petroleum on sea turtles and dolphins. *Marine Fisheries Review*, 50, 33–42.
- Knowlton, A. R., & Kraus, S. D. (2001). Mortality and serious injury of North Atlantic right whales (*Eubalaena glacialis*) in the North Atlantic Ocean. *Journal of Cetacean Resource and Management* (Special Issue), 2, 193-208.
- Kochert, M. N., Steenhof, K., McIntyre, C.L., & Craig, E.H. (2002). Golden Eagle (*Aquila chrysaetos*). In *The Birds of North America*, No. 684 (A. Poole and F. Gill, Editors). Philadelphia, PA, p. 44.
- Kranz, P.M. 1974. The anastrophic burial of bivalves and its Paleoecological significance. *The Journal of Geology* 82(2): 237-265.
- Kraus, S. D., Brown, M. D., Caswell, H., Clark, C. W., Fujiwara, M., Hamilton, P. K., ... & Rolland, R. M. (2005). North Atlantic right whales in crisis. *Science*, 309(5734), 561-562.

- Kraus, S. D., Leiter, S., Stone, K., Wikgren, B., Mayo, C., Hughes, P. .... & Tielens, J. (2016). Northeast large pelagic survey collaborative aerial and acoustic surveys for large whales and sea turtles. US Department of the Interior, Bureau of Ocean Energy Management, Sterling, Virginia. OCS Study BOEM 2016-054. 117 pp. + appendices.
- Kraus, S. D., Taylor, J. K. D., Wikgren, B., Kenney, R. D., Mayo, C., Ganley, L., ... & Rice, A. N. (2013). Field surveys of whales and sea turtles for offshore wind energy planning in Massachusetts. 2011-2012. Massachusetts Clean Energy Center.
- Kraus, S. D., Taylor, J. K. D., Wikgren, B., Kenney, R. D., Mayo, C., Ganley, L., ... & Tielens, J. (2016). Northeast large pelagic survey collaborative aerial and acoustic surveys for large whales and sea turtles. US Department of the Interior, Bureau of Ocean Energy Management, Sterling, Virginia. OCS Study BOEM 2016-054. 17 pp. + appendices.
- Krijgsveld, K. L., Fljn, R.C., Japink, M., van Horssen, P.W., Heunks, C., Collier, M.P., Poot, M.J.M., Beuker, D. & Birksen, S. (2011). Effect Studies Offshore Wind Farm Egmond aan Zee: Final report on fluxes, flight altitudes and behaviour of flying birds. *Report commissioned by NoordzeeWind*. [Online.] Retrieved from [http://www.noordzeewind.nl/wp-content/uploads/2012/02/OWEZ\\_R\\_231\\_T1\\_20080304-Flying-birds.pdf](http://www.noordzeewind.nl/wp-content/uploads/2012/02/OWEZ_R_231_T1_20080304-Flying-birds.pdf)
- Kunz, T. H., Arnett, E. B., Erickson, W. P., Hoar, A. R., Johnson, G. D., Larkin, R. P., Strickland, M. D., Thresher, R. W. & Tuttle, M. D. (2007). Ecological impacts of wind energy development on bats: Questions, research needs, and hypotheses. *Frontiers in Ecology and the Environment*, 5, 315-324.
- LaBrecque, E., Curtice, C., Harrison, J., Van Parijs, S. M., & Halpin, P. N. (2015). 2. Biologically important areas for cetaceans within US waters – East Coast region. In Van Parijs, S. M., Curtice, C., & Ferguson, M. C. (eds.). *Biologically important areas for cetaceans within US waters*. pp. 17-29. *Aquatic Mammals* (Special Issue), 41(1), 128 pp.
- Laist, D. W., Knowlton, A. R., Mead, J. G., Collet, A. S., & Podesta, M. (2001). Collisions between ships and whales. *Marine Mammal Science*, 17(1), 35-75.
- Langhamer, O. (2012). Artificial reef effect in relation to offshore renewable energy conversion: state of the art. *The Scientific World Journal* 2012: 8 pages. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3541568/pdf/TSWJ2012-386713.pdf>
- Langston, R. H. W. (2013). Birds and wind projects across the pond: A UK perspective. *Wildlife Society Bulletin* 37:5–18. doi: 10.1002/wsb.262
- Lawton, P., & Lavalli, K. L. (1995). Postlarval, Juvenile, Adolescent, and Adult Ecology, In *Biology of the Lobster, Homarus americanus*, edited by J. R. Factor. p. 51. Academic Press, San Diego, California.

- Larsen, J. K., & Guillemette, M. (2007). Effects of wind turbines on flight behaviour of wintering common eiders: implications for habitat use and collision risk. *Journal of Applied Ecology* 44:516–522. doi: 10.1111/j.1365-2664.2007.1303.x
- LaSalle, M.W., Clarke, D.G., Homziak, J., Lunz, J.D., & Fredette, T.J. (1991). A framework for assessing the need for seasonal restrictions on dredging and disposal operations. US Army Engineer Waterways Experiment Station, Technical report D-91-1, Vicksburg, Mississippi.
- Lavers, J., Hipfner, J.M., & Chapdelaine, G. (2009). Razorbill (*Alca torda*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY.
- Lavigne, D. M., & Kovacs, K. M. (1988). Harps and hoods: Ice breeding seals of the Northwest Atlantic. 174 pp. Waterloo, Ontario, Canada: University of Waterloo Press.
- Laviguer, L., & Hammill, M. O. (1993). Distribution and seasonal movements of grey seals, *Halichoerus grypus*, born in the Gulf of St. Lawrence and eastern Nova Scotia shore. *The Canadian Field Naturalist*, 107, 329-340.
- Lazell, J. D., Jr. (1980). New England waters: Critical habitat for marine turtles. *Copeia*, 2, 290-295.
- Lavender, A. L., Bartol, S. M., & Bartol, I. K. (2014). Ontogenetic investigation of underwater hearing capabilities in loggerhead sea turtles (*Caretta caretta*) using a dual testing approach. *Journal of Experimental Biology*, 217, 2580-2589.
- Lawson, J. W., & Gosselin, J. F. (2009). Distribution and preliminary abundance estimates for cetaceans seen during Canada's Marine Megafauna Survey - A component of the 2007 TNASS. Canadian Science Advisory Secretariat Research Document, 2009/031. 33 pp.
- Lawson, J. W., & Gosselin, J. F. (2011). Fully-corrected cetacean abundance estimates from the Canadian TNASS survey. Working Paper 10. National Marine Mammal Peer Review Meeting. Ottawa, Canada. 28 pp.
- Leatherwood, S., Caldwell, D. K., & Winn, H. E. (1976). Whales, dolphins, and porpoises of the western North Atlantic. A guide to their identification. NOAA Tech. Rep. NMFS Circ. 396. 176 pp.
- Leddy, K. L., Higgins, K.F., & Naugle, D.E. (1999). Effects of wind turbines on upland nesting birds in Conservation Reserve Program grasslands. *Wilson Bulletin* 111:100–104.
- Lee, D. S. (1989). Jaegers and skuas in the Western North Atlantic : some historical misconceptions. *American Birds* 43:18–20.



- Leonhard, S. B., Pedersen, J., Grøn, P.N., Skov, H., Jansen, J., Topping, C.J., & Petersen, I.K. (2013). Wind farms affect Common Scoter and Red-throated Diver behaviour. *Danish Offshore Wind: Key Environmental Issues – A Follow-up*. [Online.] Retrieved from [http://orbit.dtu.dk/ws/files/55080896/Havvindm\\_llebog.pdf](http://orbit.dtu.dk/ws/files/55080896/Havvindm_llebog.pdf)
- Lessage, V., & Hammill, M. O. (2001). The status of the grey seal, *Halichoerus grypus*, in the Northwest Atlantic. *The Canadian Field Naturalist*, 115(4), 653-662.
- Lichtman, I.D., J.H. Baas, L.O. Amoudry, P.D. Thorne, J. Malarkey, J.A. Hope, J. Peakall, D.M. Paterson, S.J. Bass, R.D. Cooke, A.J. Manning, A.G. Davies, D.R. Parsons, L. Ye. (2018). Bedform migration in a mixed sand and cohesive clay intertidal environment and implications for bed material transport predictions. *Geomorphology* 315: 17-32.
- Lindeboom, H. J., Kouwenhoven, H.J., Bergman, M.J.N., Bouma, S., Brasseur, S., Daan, R., Fijn, R.C., Dirksen, S., van Hal, R., Hille Ris Lambers, R., et al. (2011). Short-term ecological effects of an offshore wind farm in the Dutch coastal zone; a compilation. *Environmental Research Letters* 6. doi: 10.1088/1748-9326/6/3/035101
- Lindholm, J.B., Auster, P.J., Ruth, M., & Kaufman, L. (2001). Modeling the effects of fishing and implications for the design of marine protected areas: juvenile fish responses to variations in seafloor habitat. *Conservation Biology*, 15(2), pp.424-437.
- Linnane, A., Ball, B., Mercer, J.P., Van der Meeren, G., Bannister, C., Mazzoni, D., Munday, B. & Ringvold, H. (1999). Understanding the factors that influence European lobster recruitment, a trans-European study of cobble fauna. *J. Shellfish Res.* 18, (2), 719–720.
- List of Endangered Foreign Fish and Wildlife, 35 Fed. Reg. 18319 (December 2, 1970).
- Listing and Protecting Loggerhead Sea Turtles as “Threatened Species” and Populations of Green and Olive Ridley Sea Turtles as Threatened Species or “Endangered Species,” 43 Fed. Reg. 32800 (July 28, 1978).
- Lohmann, K. J. (1991). Magnetic orientation by hatchling loggerhead sea turtles (*Caretta caretta*). *Journal of Experimental Biology*, 155, 37-49.
- Lohmann, K. J., & Lohmann, C. M. F. (1994). Acquisition of magnetic directional preference in hatchling loggerhead sea turtles. *Journal of Experimental Biology*, 190, 1-8.
- Lohmann, K. J., & Lohmann, C. M. F. (1996). Detection of magnetic field intensity by sea turtles. *Nature*, 380, 59-61.

- Lohmann, K. J., Lohmann, C. M. F., and Putman, N. F. (2007). Magnetic maps in animals: Nature's GPS. *Journal of Experimental Biology*, 190, 1-8. 210: 3697-3705.
- Løkkeborg, S., Humborstad, O.B., Jørgensen, T., & Soldal, A.V. (2002). Spatio-temporal variations in gillnet catch rates in the vicinity of North Sea oil platforms. *ICES Journal of Marine Science*, 59, 294-299.
- Longhurst, A. R. (1998). *Ecological geography of the sea*. Second edition. Elsevier Academic Press. 560 pp.
- Loring, P., Goyert, H., Griffin, C., Sievert, P., & Paton, P. (2017). Tracking Movements of Common Terns, Endangered Roseate Terns, and Threatened Piping Plovers in the Northwest Atlantic: 2017 Annual Report to the Bureau of Ocean Energy Management (BOEM). In. Interagency Agreement No. M13PG00012 to US. Fish and Wildlife Service Northeast Region Division of Migratory Birds, Hadley, Massachusetts.
- Love, M. S., Nishimoto, M. M., Clark, S., McCrea, M., & Bull, A. S. (2017). The Organisms Living Around Energized Submarine Power Cables, Pipe, and Natural Sea Floor in the Inshore Waters of Southern California. *Bulletin, Southern California Academy of Sciences* 116(2), pp.61-87.
- Love, M., Robbins, C., Baldera, A., Eastman, S., Bolten, A., Hardy, R., ... & Wallace, B. (2017). Restoration with borders: An Assessment of cumulative stressors to guide large-scale, integrated restoration of sea turtles in the Gulf of Mexico. Retrieved from [https://oceanconservancy.org/wp-content/uploads/2017/10/Restoration Without Borders.pdf](https://oceanconservancy.org/wp-content/uploads/2017/10/Restoration_Without_Borders.pdf). Accessed on November 1, 2017.
- Lowe, C.G., Anthony, K.M., Jarvis, E.T., Bellquist, L.F., & Love, M.S. (2009). Site fidelity and movement patterns of groundfish associated with offshore petroleum platforms in the Santa Barbara channel. *Marine and Coastal Fisheries: Dynamics, Management and Ecosystem Science*, 1, 71-89.
- Lowther, P. E., Diamond, A.W., Kress, S.W., Robertson, G.J., & Russell, K. (2002). Atlantic Puffin (*Fratercula arctica*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY.
- MacArthur, R. (1955). Fluctuations of Animal Populations and a Measure of Community Stability. *Ecology*, 36, 533-536.
- MacKenzie, C.L. Jr., D.J. Radosh, and R.N. Reid. (1985). Densities, growth, and mortalities of juveniles of the surf clam (*Spisula solidissima*) (Dillwyn) in the New York Bight. *J. Shellfish Res.* 5: 81-84.

- Madsen, P. T., Møhl, B., Nielsen, K., & Wahlberg, M. (2002). Male sperm whale behaviour during exposures to distant seismic survey pulses. *Aquatic Mammals*, 28, 231–240.
- Madsen, P. T., Wahlberg, M., Tougaard, J., Lucke, K., & Tyack, P. (2006). Wind turbine underwater noise and marine mammals: Implications of current knowledge and data needs. *Marine Ecology Progress Series*, 309, 279-295.
- [MAFMC] Mid-Atlantic Fishery Management Council. (1997). Amendment #10 to the Fishery Management Plan for Atlantic surfclam and ocean quahog fisheries. March 1997. MAFMC. [Dover, DE.] 58 p. + appendices.
- Mansfield, A. W. (1966). The grey seal in eastern Canadian waters. *Can. Audubon Mag.*, 28, 161-166.
- Martha's Vineyard Commission (2006). *Martha's Vineyard Economic Profile*. Oak Bluffs, MA. Report available at: <http://www.mvcommission.org/sites/default/files/docs/economicprofile.pdf>
- Martha's Vineyard Commission (2008). *Martha's Vineyard Economic Profile*. Oak Bluffs, MA. Report available at: <http://www.mvcommission.org/sites/default/files/docs/economicprofileryanwithappendices.pdf>
- Martha's Vineyard Transit Authority. (2017) *About Us*. Retrieved from: [http://www.vineyardtransit.com/Pages/VTA\\_Webdocs/about](http://www.vineyardtransit.com/Pages/VTA_Webdocs/about)
- Martin, C. M., Arnett, E. B., Stevens, R. D. & Wallace, M. C. (2017). Reducing bat fatalities at wind facilities while improving the economic efficiency of operational mitigation. *Journal of Mammalogy*, 98, 378-385.
- Martin, K. J., Alessi, S. C., Gaspard, J. C., Tucker, A. D., Bauer, G. B., & Mann, D. A. (2012). Underwater hearing in the loggerhead sea turtles (*Caretta caretta*): A comparison of behavioural and auditory evoked potential audiograms. *Journal of Experimental Biology*, 215, 3001-3009.
- Masden, E. A., Haydon, D.T., Fox, A.D., Furness, R.W., Bullman, R., & Desholm, M. (2009). Barriers to movement: impacts of wind farms on migrating birds. *ICES Journal of Marine Science* 66:746–753. doi: 10.1093/icesjms/fsp031
- Massachusetts Department of Energy Resources (DOER). (2017). Request for proposals for long-term contracts for offshore wind energy projects. Retrieved from <https://macleanenergy.files.wordpress.com/2017/02/section-83c-request-for-proposals-for-long-term-contracts-for-offshore-wind-energy-projects-june-29-2017.pdf>

- Massachusetts Department of Environmental Protection (MassDEP), Bureau of Air and Waste, Division of Air and Climate Programs. (2017). Massachusetts 2016 Air Quality Report. Retrieved from <http://www.mass.gov/eea/docs/dep/air/priorities/16aqrpt.pdf>
- Massachusetts Department of Transportation (MassDOT). (2010). *2010 Massachusetts Statewide Airport Plan*. Boston, MA.
- Massachusetts Executive Office of Energy and Environmental Affairs (EEA). (2015). *Massachusetts Ocean Management Plan*. Boston, MA.
- Massachusetts Natural Heritage and Endangered Species Program, Massachusetts Division of Fisheries and Wildlife, Trustees of Reservations, Nature Conservancy, Massachusetts Audubon Society. (2013). An Action Plan for the Conservation of State-listed Obligate Grassland Birds in Massachusetts. Retrieved from <http://www.mass.gov/eea/docs/dfg/nhesp/species-and-conservation/grassland-bird-plan-final.pdf>
- Mass.Gov. (2016). Governor Baker signs comprehensive energy diversity legislation. Retrieved from <http://www.mass.gov/governor/press-office/press-releases/fy2017/governor-baker-signs-comprehensive-energy-diversity-law.html>
- Masswildlife. (2017). The Northern Long-eared Bat. Retrieved from: <https://http://www.mass.gov/service-details/the-northern-long-eared-bat>.
- Masswildlife. (2012). Northern Long-eared Bat. Retrieved from: <http://www.mass.gov/eea/docs/dfg/nhesp/species-and-conservation/nhfacts/myotis-septentrionalis.pdf>.
- Masswildlife News.(2008). Bat Mortalities in Massachusetts. Retrieved from: [https://http://www.whitenosesyndrome.org/sites/default/files/files/mass\\_bat\\_mortalities.pdf](https://http://www.whitenosesyndrome.org/sites/default/files/files/mass_bat_mortalities.pdf).
- Mayo, C. A., & Marx, M. K. (1990). Surface foraging behaviour of the North Atlantic right whale, *Eubalaena glacialis*, and associated zooplankton characteristics. *Canadian Journal of Zoology*, 68, 2214-2220.
- McAlpine, D. F., James, M. C., Lien, J., & Orchard, S. A. (2007). Status and conservation of sea turtles in Canadian waters. In: Seburn, C. N. L., & Bishop, C. A. (eds.) Ecology, Conservation, and Status of Reptiles in Canada. pp. 85-112. Herpetological Conservation Number 2. Society for the Study of Amphibians and Reptiles, Salt Lake City, Utah.
- McCaughey, R. D., Fewtrell, J., Duncan, A. J., Jenner, C., Jenner, M. N., Penrose, J. D., ... & McCabe, K. (2000). Marine seismic surveys – a study of environmental implications. *APPEA Journal*, 692-708.

- McGrady, M. J., Young, G.S., & Seegar, W.S. (2006). Migration of a Peregrine Falcon *Falco peregrinus* over water in the vicinity of a hurricane. *Ringing and Migration* 23:80–84.
- McKenna, L. N. (2016). Vocalizations of sea turtle hatchlings and embryos. MS thesis, Purdue University, Fort Wayne, Indiana. Retrieved from [http://opus.ipfw.edu/masters\\_theses/50](http://opus.ipfw.edu/masters_theses/50). Accessed November 1, 2017.
- Mead, J. G., & Potter, C. W. (1995). Recognizing two populations for the bottlenose dolphin (*Tursiops truncatus*) off the Atlantic coast of North America: Morphologic and ecologic considerations. *International Biological Research Institute Reports*, 5, 31-43.
- Meek, E. R., Ribbands, J.B., Christer, W.G., Davy, P.R. & Higginson, I. (1993). The effects of aerogenerators on moorland bird populations in the Orkney Islands, Scotland. *Bird Study* 40:140–143. doi: 10.1080/00063659309477139
- Mendelsohn, J. M., Kemp, A.C., Biggs, H.C., Biggs, R., & Brown, C.J. (1989). Wing areas, wing loadings and wing spans of 66 species of African raptors. *Ostrich* 60:35–42.
- Menzel, M. A., Owen, S. F., Ford, W. M., Edwards, J. W., Wood, P. B., Chapman, B. R. & Miller, K. V. (2002). Roost tree selection by northern long-eared bat (*Myotis septentrionalis*) maternity colonies in an industrial forest of the central Appalachian mountains. *Forest Ecology and Management*, 155, 107-114.
- Meyer, T.L., R.A. Cooper, and K.J. Pecci. (1981). The performance and environmental effects of a hydraulic clam dredge. *Mar. Fish. Rev.* 43(9): 14-22.
- Miller, P. J. O., Johnson, M. P., Madsen, P. T., Biassoni, N., Quero, M., & Tyack, P. L. (2009). Using at-sea experiments to study the effects of airguns on the foraging behavior of sperm whales in the Gulf of Mexico. *Deep Sea Research I*. doi:10.1016/j.dsr.2009.02.008.
- Mizrahi, D., Fogg, R., Peters, K.A., & Hodgetts, P.A. (2009). Assessing nocturnal bird and bat migration patterns on the Cape May peninsula using marine radar: potential effects of a suspension bridge spanning Middle Thoroughfare, Cape May County, New Jersey.
- MMS. (2007). Programmatic Environmental Impact Statement for alternative energy development and production and alternate use of facilities on the Outer Continental Shelf: Final Environmental Impact Statement.
- MMS. (2008). Cape Wind Energy Project Nantucket Sound Biological Assessment (Appendix G). In Cape Wind Energy Project Final EIS. p. 296.
- Moein, S. E., Musick, J. A., Keinath, J. A., Barnard, D. E., Lenhardt, M. L., & George, R. (1995). Evaluation of seismic sources for repelling sea turtles from hopper dredges. In: Hales, L. Z. (ed.), *Sea Turtle Research Program: Summary Report*. pp. 90-93. Technical Report CERC-95.

- Mojica, E. K., Meyers, J.M., Millsap, B.A., & Haley, K.L. (2008). Migration Of Florida Sub-Adult Bald Eagles. *The Wilson Journal of Ornithology* 120:304–310. doi: 10.1676/07-079.1
- Mojica, E. K., Watts, B.D., & Turrin, C.L. (2016). Utilization Probability Map for Migrating Bald Eagles in Northeastern North America: A Tool for Siting Wind Energy Facilities and Other Flight Hazards. *PLoS ONE* 11. doi: 10.1371/journal.pone.0157807
- Molvaer, O. and T. Gjestland. (1981). Hearing damage risk to divers operating noisy tools under water. *Scandinavian Journal of Work, Environment, and Health*, 7: 263-270
- Montevecchi, W. A. (2006). Influences of Artificial Light on Marine Birds. In *Ecological Consequences of Artificial Night Lighting* (C. Rich and T. Longcore, Editors). Island Press, Washington DC, pp. 94–113.
- Morreale, S. J., Meylan, A., Sadove, S. S., & Standora, E. A. (1992). Annual occurrence and winter mortality of marine turtles in New York waters. *Journal of Herpetology*, 26, 301-308.
- Morris, S. R., Richmond, M.E., & Holmes, D.W. (1994). Patterns of stopover by warblers during spring and fall migration on Appledore Island, Maine. *Wilson Bulletin* 106:703–718.
- Mostello, C. S., & Longsdorf, J. (2017). Inventory of Terns, Laughing Gulls, and Black Skimmers Nesting in Massachusetts in 2015.
- Mostello, C. S., Nisbet, I.C.T, Oswald, S.A., & Fox, J.W. (2014). Non-breeding season movements of six North American Roseate Terns *Sterna dougallii* tracked with geolocators. *Seabird* 27:1–21.
- Mowbray, T. B. (2002). Northern Gannet (*Morus bassanus*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY. doi: 10.2173/bna.693
- Mrosovsky, N. (1972). Spectrographs of the sounds of leatherback turtles. *Herpetologica*, 29(3), 256-258.
- Murie, O. J. (1940). Food Habits of the Northern Bald Eagle in the Aleutian Islands, Alaska. *The Condor* 42:198–202.
- Murphy, R.C. (1985). Factors affecting the distribution of the introduced bivalve, *Mercenaria*, in a California lagoon- the importance of bioturbation. *Journal of Marine Research* 43:673-692.
- Nachtigall, P. E., & Supin, A. Y. (2008). A false killer whale adjusts its hearing when it echolocates. *Journal of Experimental Biology*, 211, 1714-1718.
- Nantucket Planning Board. (2009). *Nantucket Master Plan, Prepared in Accordance with M.G.L. Ch 41, Section 81D*. Nantucket, MA

- National Aeronautics and Space Administration (NASA). (2015). Science briefs: Sea level rise, after the ice melted and today. [http://www.giss.nasa.gov/research/briefs/gornitz\\_09/](http://www.giss.nasa.gov/research/briefs/gornitz_09/). Accessed October 29, 2015.
- National Oceanic and Atmospheric Administration (NOAA). (1991). Recovery plan for the humpback whale (*Megaptera novaeangliae*). Silver Spring, Maryland.
- National Oceanic and Atmospheric Administration (NOAA). (1993). Stellwagen Bank Management Plan and Final Environmental Impact Statement.
- National Oceanic and Atmospheric Administration (NOAA). (2010). Recovery plan for the sperm whale (*Physeter macrocephalus*). National Marine Fisheries Service, Silver Spring, Maryland. 165 pp.
- National Oceanic and Atmospheric Administration (NOAA). (2012). Annual Average Ambient Noise Modeling Methodology: Global Shipping. Retrieved from [https://cetsound.noaa.gov/sound\\_data](https://cetsound.noaa.gov/sound_data). Accessed on December 8, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2013). Commercial wind lease issuance and site assessment activities on the Atlantic Outer Continental Shelf in Massachusetts, Rhode Island, New York, and New Jersey wind energy area Endangered Species Act Section 7 consultation biological opinion.
- National Oceanic and Atmospheric Administration (NOAA). (2014a). Harbor porpoise (*Phocoena phocoena*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/porpoises/harbor-porpoise.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2014b). Minke whale (*Balaenoptera acutorostrata*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/whales/minke-whale.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2015a). Harp seal (*Pagophilus groenlandicus*). Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/seals/harp-seal.html>. Accessed on November 20, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2015b). Risso's dolphin (*Grampus griseus*). Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/dolphins/rissos-dolphin.html>. Accessed on November 20, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2015c). Sei Whale (*Balaenoptera borealis*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/whales/sei-whale.html>. Accessed October 17, 2017.

- National Oceanic and Atmospheric Administration (NOAA). (2015d). Shortnose Sturgeon (*Acipenser brevirostrum*). Retrieved from <http://www.nmfs.noaa.gov/pr/species/fish/shortnose-sturgeon.html>
- National Oceanic and Atmospheric Administration (NOAA). (2016a). Amendment 10 to the 2006 Consolidated Atlantic Highly Migratory Species Fishery Management Plan: Essential Fish Habitat. Retrieved from [http://www.nmfs.noaa.gov/sfa/hms/documents/fmp/am10/draft\\_maps/hms\\_efh\\_amendment\\_10\\_draft\\_ea\\_0648\\_xd990\\_final\\_090716.pdf](http://www.nmfs.noaa.gov/sfa/hms/documents/fmp/am10/draft_maps/hms_efh_amendment_10_draft_ea_0648_xd990_final_090716.pdf)
- National Oceanic and Atmospheric Administration (NOAA). (2016b). Atlantic salmon (*Salmo salar*). Retrieved from <http://www.nmfs.noaa.gov/pr/species/fish/atlantic-salmon.html>
- National Oceanic and Atmospheric Administration (NOAA). (2016c). Atlantic white-sided dolphin (*Lagenorhynchus acutus*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/dolphins/atlantic-white-sided-dolphin.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2016d). Bottlenose dolphin (*Tursiops truncatus*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/dolphins/bottlenose-dolphin.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2016e). Fin whale (*Balaenoptera physalus*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/whales/fin-whale.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2016f). Harbor seal (*Phoca vitulina*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/seals/harbor-seal.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2016g). Humpback whale (*Megaptera novaeangliae*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/whales/humpback-whale.html#description>. Accessed November 19, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2016h). Gray seal (*Halichoerus grypus*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/seals/gray-seal.html>. Accessed November 22, 2017.



- National Oceanic and Atmospheric Administration (NOAA). (2016i). Long-finned pilot whale (*Globicephala melas*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/whales/long-finned-pilot-whale.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2016j). Integrative Statistical Modeling and Predictive Mapping of Marine Bird Distributions and Abundance on the Atlantic Outer Continental Shelf: Maine – Florida. [Online.] Retrieved from <https://coastalscience.noaa.gov/projects/detail?key=279>
- National Oceanic and Atmospheric Administration (NOAA). (2016k). North Atlantic right whales (*Eubalaena glacialis*). Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/whales/north-atlantic-right-whale.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2016l). Short-beaked common dolphin (*Delphinus delphis*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/dolphins/short-beaked-common-dolphin.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2016m). Short-finned pilot whale (*Globicephala macrorhynchus*) species description. Retrieved from <http://www.nmfs.noaa.gov/pr/species/mammals/whales/short-finned-pilot-whale.html>. Accessed November 22, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2017a). 2016-2017 Humpback Whale Unusual Mortality Event along the Atlantic Coast. Retrieved from <http://www.nmfs.noaa.gov/pr/health/mmume/2017humpbackatlanticume.html>. Accessed October 19, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2017b). Large Marine Ecosystems of the World. [Online.] Retrieved from [http://www.lme.noaa.gov/index.php?option=com\\_content&view=featured&Itemid=101](http://www.lme.noaa.gov/index.php?option=com_content&view=featured&Itemid=101)
- National Oceanic and Atmospheric Administration (NOAA). (2017c). NOAA Deep-Sea Coral Data Portal. Funded by NOAA's Deep Sea Coral Research and Technology Program (DSCRTP). Retrieved from <https://deepseacoraldata.noaa.gov/website/AGSViewers/DeepSeaCorals/mapSites.htm>
- National Oceanic and Atmospheric Administration (NOAA). (2017d). 2017 North Atlantic right whale unusual mortality event. Retrieved from <http://www.nmfs.noaa.gov/pr/health/mmume/2017northatlanticrightwhaleume.html>. Accessed November 18, 2017.

- National Oceanic and Atmospheric Administration (NOAA). (2017e). Sperm whale (*Physeter macrocephalus*) species description. Retrieved from <http://www.fisheries.noaa.gov/pr/species/mammals/whales/sperm-whale.html>. Accessed November 17, 2017.
- National Oceanic and Atmospheric Administration (NOAA). (2017f). *United States Coast Pilot 2 Atlantic Coast: Cape Cod, MA to Sandy Hook, NJ (46th Edition)*. US Department of Commerce, National Oceanic and Atmospheric Administration (NOAA), National Ocean Service. Retrieved October 4, 2017, from [https://www.nauticalcharts.noaa.gov/nsd/coastpilot/files/cp2/CPB2\\_E46\\_20170804\\_1803\\_WEB.pdf](https://www.nauticalcharts.noaa.gov/nsd/coastpilot/files/cp2/CPB2_E46_20170804_1803_WEB.pdf)
- National Science Foundation and US Geological Survey (NSF & USGS). (2011). Programmatic environmental impact statement/overseas environmental impact statement for marine seismic research funded by the National Science Foundation or the US Geological Survey.
- NatureServe. (2010). NatureServe Explorer: An outline encyclopedia of life [web application]. Version 7.1, NatureServe. Arlington, Virginia.
- Navy (Department of the Navy). (2007). Navy OPAREA density estimates (NODE) for the northeast OPAREAS: Boston, Narragansett Bay, and Atlantic City. Naval Facilities Engineering Command, Atlantic; Norfolk, Virginia. Contract N62470-02-D-9997, Task Order 0045. Prepared by Geo-Marine, Inc., Plano, Texas.
- NEAMAP. (2018). Virginia Institute of Marine Science Northeast Area Monitoring and Assessment Program. Retrieved from [http://www.vims.edu/research/departments/fisheries/programs/multispecies\\_fisheries\\_research/neamap/index.php](http://www.vims.edu/research/departments/fisheries/programs/multispecies_fisheries_research/neamap/index.php)
- Nedelec, S. L., Radford, A. N., Simpson, S. D., Nedelec, B., Lecchini, D., & Mills, S. C. (2014). Anthropogenic noise playback impairs embryonic development and increases mortality in a marine invertebrate. *Scientific reports*, 4.
- Nedwell, J. & Howell, D. (2004). A review of offshore windfarm related underwater noise sources. Report commissioned by COWRIE. Retrieved from <https://tethys.pnnl.gov/publications/review-offshore-windfarm-related-underwater-noise-sources>. Accessed on September 21, 2017.
- Nedwell, J., Parvin, S., Edwards, B., Workman, R., Brooker, A., & Jynoch, J. (2007). Measurement and interpretation of underwater noise during construction and operation of offshore windfarms in UK waters.
- Neill, W. H., Stevens, E. D. (1974). Thermal inertia versus thermoregulation in "warm" turtles and tunas. *Science*, 184, 1008-1010.

- NEODP (Northeast Ocean Data Portal). (2017). Northeast Ocean Data: Maps and Data for Ocean Planning in the Northeastern United States. Data Explorer. Retrieved from <http://www.northeastoceandata.org/data-explorer/>. Accessed September 2017
- NEFSC. (2014). Annual Report of a Comprehensive Assessment of Marine Mammal, Marine Turtle, and Seabird Abundance and Spatial Distribution in US Waters of the Western North Atlantic Ocean. Appendix C: Shipboard habitat survey during March – April 2014: Northeast Fisheries Science Center. Retrieved from [http://www.nefsc.noaa.gov/psb/AMAPPS/docs/NMFS AMAPPS 2014 annual report Final.pdf](http://www.nefsc.noaa.gov/psb/AMAPPS/docs/NMFS_AMAPPS_2014_annual_report_Final.pdf)
- NEFSC. (2017a). Multispecies bottom trawl survey [Water quality monitoring data file]. Retrieved from [http://comet.nefsc.noaa.gov/cgi-bin/ioos/index\\_hydro1.pl](http://comet.nefsc.noaa.gov/cgi-bin/ioos/index_hydro1.pl)
- NEFSC. (2017b). Spatial Analyses of Invertebrate Species. Retrieved from <https://www.nefsc.noaa.gov/ecosys/spatial-analyses/>.
- NEFSC. (2018). Surfclam and Ocean Quahog Survey Resource Survey Reports (RSR). NEFSC Ecosystems Surveys Branch. Retrieved from [https://www.nefsc.noaa.gov/femad/ecosurvey/mainpage/resource\\_survey\\_reports.html](https://www.nefsc.noaa.gov/femad/ecosurvey/mainpage/resource_survey_reports.html).
- NEFSC & SEFSC. (2010). 2010 Annual report to the inter-agency agreement M10PG00075/0001: A comprehensive assessment of marine mammal, marine turtle, and seabird abundance and spatial distribution in US waters of the western North Atlantic Ocean.
- NEFSC & SEFSC. (2011). 2011 Annual report to the inter-agency agreement M10PG00075/0001: A comprehensive assessment of marine mammal, marine turtle, and seabird abundance and spatial distribution in US waters of the western North Atlantic Ocean.
- NEFSC & SEFSC. (2012). 2012 Annual report to the inter-agency agreement M10PG00075/0001: A comprehensive assessment of marine mammal, marine turtle, and seabird abundance and spatial distribution in US waters of the western North Atlantic Ocean.
- NEFSC & SEFSC. (2013). 2013 Annual report to the inter-agency agreement M10PG00075/0001: A comprehensive assessment of marine mammal, marine turtle, and seabird abundance and spatial distribution in US waters of the western North Atlantic Ocean.
- NEFSC & SEFSC. (2014). 2014 Annual report to the inter-agency agreement M10PG00075/0001: A comprehensive assessment of marine mammal, marine turtle, and seabird abundance and spatial distribution in US waters of the western North Atlantic Ocean.
- NEFSC & SEFSC. (2015). 2015 Annual report to the inter-agency agreement M10PG00075/0001: A comprehensive assessment of marine mammal, marine turtle, and seabird abundance and spatial distribution in US waters of the western North Atlantic Ocean. – AMAPPS II.

- NEFSC & SEFSC. (2016). Annual report to the inter-agency agreement M10PG00075/0001: A comprehensive assessment of marine mammal, marine turtle, and seabird abundance and spatial distribution in US waters of the western North Atlantic Ocean. – AMAPPS II.
- New Bedford Harbor Development Commission (NBHDC). (2016). *Economic Impact Study of New Bedford/Fairhaven Harbor: Examining the Economic Impact of the Port of New Bedford and the Phase V & Navigational Dredging in New Bedford/Fairhaven Harbor*. New Bedford, MA.
- Newcombe, C.P., & MacDonald, D.D. (1991). Effects of suspended sediments on aquatic ecosystems. *North American Journal of Fisheries Management*, 11, 72-82.
- Newcombe, C.P., & Jensen, J.O.T. (1996). Channel suspended sediment and fisheries: a synthesis for quantitative assessment of risk and impact. *North American Journal of Fisheries Management*, 16, 693-727.
- Newell, R.I.E. (1989). Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (North and Mid-Atlantic)–blue mussel. US Fish. Wildl. Serv. Biol. Rep. 82(11. 102). US Army Corps of Engineers, T R EI-82-4. 25 pp. Retrieved from [http://www.nwrc.usgs.gov/wdb/pub/species\\_profiles/82\\_11-102.pdf](http://www.nwrc.usgs.gov/wdb/pub/species_profiles/82_11-102.pdf)
- New Jersey Department of Agriculture (n.d.). *New Jersey Fishing and Aquaculture: Harvesting the Garden State's Waters*. Trenton, NJ. Report available at: <http://www.jerseyseafood.nj.gov/seafoodreport.pdf>
- Nisbet, I. C. T. (1963). Measurements with radar of the height of nocturnal migration over Cape Cod, Massachusetts. *Bird-Banding* 34:57–67.
- Nisbet, I. C. T. (1984). Migration and winter quarters of North American Roseate Terns as shown by banding recoveries. *Journal of Field Ornithology* 55:1–17.
- Nisbet, I. C. T., Gochfeld, M. & Burger, J. (2014). Roseate Tern (*Sterna dougallii*). *The Birds of North America Online*. [Online.] Retrieved from <http://bna.birds.cornell.edu/bna/species/370>. doi: 10.2173/bna.370
- Nisbet, I. C. T., & Veit, R.R. (2015). An explanation for the population crash of Red-necked Phalaropes *Phalaropus lobatus* staging in the Bay of Fundy in the 1980s. *Marine Ornithology* 43:119–121.
- Nisbet, I. C. T., Veit, R.R., Auer, S.A. & White, T.P. (2013). Marine Birds of the Eastern United States and the Bay of Fundy: Distribution, Numbers, Trends, Threats, and Management. In. No. 29. Nuttall Ornithological Club, Cambridge, MA.

- Nisbet, I. C. T., Weseloh, D.V., Hebert, C.E., Mallory, M.L., Poole, A.F., Ellis, J.C., Pyle, P. & Patten, M.A. (2017). Herring Gull (*Larus argentatus*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY. doi: 10.2173/bna.hergul.03
- NJSDEP. (2010). Ocean/wind power ecological baseline studies. Volume I: Overview, summary, and application.
- NMFS. (2002). Endangered Species Act Section 7 consultation on proposed Gulf of Mexico outer continental shelf multi-lease sales (185, 187, 190, 192, 194, 196, 198, 200, 201). Biological Opinion. November 29, 2002.
- NMFS & USFWS. (1991). *Recovery plan for U.S. population of Atlantic Green Turtle*. National Marine Fisheries Service, Silver Spring, MD.
- NMFS & USFWS. (1992a). Recovery plan for leatherback turtles (*Dermochelys coriacea*) in the US Caribbean, Atlantic and Gulf of Mexico. NMFS, Silver Spring, Maryland. 69 pp.
- NMFS & USFWS. (1992b). Recovery plan for the Kemp's Ridley Sea Turtle (*Lepidochelys kempi*). NMFS, Silver Spring, Maryland. 47 pp.
- NMFS & USFWS. (1993). *Recovery plan for Hawksbill Turtles in the U.S. Caribbean Sea, Atlantic Ocean, and Gulf of Mexico*. National Marine Fisheries Service, Silver Spring, MD.
- NMFS & USFWS. (2008). Recovery plan for the Northwest Atlantic Population of the Loggerhead Sea Turtle (*Caretta caretta*). NMFS, Silver Spring, Maryland. 325 pp.
- NMFS, USFWS, & SEMARNAT (National Marine Fisheries Service, US Fish and Wildlife Service, and Secretariat of Environment and Natural Resources). (2011). Bi-national recovery plan for the Kemp's ridley sea turtle (*Lepidochelys kempi*), second revision. NMFS, Silver Spring, Maryland. 156 pp. + appendices.
- NOAA NBDC (National Oceanic Atmospheric Administration National Data Buoy Center). (2017). [Water quality monitoring data file]. Retrieved from [http://www.ndbc.noaa.gov/historical\\_data.shtml](http://www.ndbc.noaa.gov/historical_data.shtml)
- NODP (Northeast Ocean Data Portal). (2017). Fish: Individual Species. Retrieved from <http://www.northeastoceandata.org/data-explorer/?fish>
- Norberg, U. M. & Rayner, J. M. V. (1987). Ecological morphology and flight in bats (*Mammalia; Chiroptera*): wing adaptations, flight performance, foraging strategy and echolocation. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 316, 335-427.

- Norling P., Kautsky, N. (2007). Structural and functional effects of *Mytilus edulis* on diversity of associated species and ecosystem functioning. *Marine Ecology Progress Series* 351:163–175, DOI 10.3354/meps07033.
- Normandeau Associates Inc. (2011). New Insights and New Tools Regarding Risk to Roseate Terns, Piping Plovers, and Red Knots from Wind Facility Operations on the Atlantic Outer Continental Shelf. *Report No. BOEMRE 048-2001. Contract No. M08PC20060*. [Online.] Retrieved from [www.data.boem.gov/PI/PDFImages/ESPIS/4/5119.pdf](http://www.data.boem.gov/PI/PDFImages/ESPIS/4/5119.pdf)
- Normandeau Associates, Inc. (2012). Effects of Noise on Fish, Fisheries, and Invertebrates in the US Atlantic and Arctic from Energy Industry Sound-Generating Activities. A Workshop Report for the US Dept. of the Interior, Bureau of Ocean Energy Management, 72 pp. plus Appendices.
- Normandeau Associates, Inc. (2017). Benthic Sample Processing Results, Vineyard Wind Cable Route Survey, September 2017, 29 pp.
- Normandeau Associates Inc., Exponent Inc., Tricas, T., & Gill, A. (2011). Effects of EMFs from undersea power cables on elasmobranchs and other marine species, final report. Prepared under BOEMRE Contact M09P C00014. OCS Study BOEMRE 2011-09. Published by the US Department of the Interior Pacific OCS Region.
- Normandeau, E., Tricas, T., & Gill, A. (2011). Effects of EMFs from Undersea Power Cables on Elasmobranchs and Other Marine Species. US Dept. of the Interior, Bureau of Ocean Energy Management, Regulation, and Enforcement, Pacific OCS Region, Camarillo, CA. OCS Study BOEMRE 2011-09.
- Northeast Ocean Data Portal (“NEODP”). 2017. Northeast Ocean Data: Maps and Data for Ocean Planning in the Northeastern United States. Data Explorer. Retrieved from <http://www.northeastoceandata.org/data-explorer/>
- Northeast Regional Planning Body (2016). *Northeast Ocean Plan*. Retrieved from: [http://neoplan.org/wp-content/uploads/2016/10/Northeast-Ocean-Plan\\_Full.pdf](http://neoplan.org/wp-content/uploads/2016/10/Northeast-Ocean-Plan_Full.pdf)
- Northridge S. P., Tasker M. L., Webb A., Camphuysen, K., & Leopold, M. (1997). White-beaked Lagenorhynchus albirostris and Atlantic white-sided dolphin L. acutus distributions in Northwest European and US North Atlantic waters. Forty-seventh report of the International Whaling Commission, 47, 797-805.
- Nowacek, D. P., Johnson, M. P., & Tyack, P. L. (2004). North Atlantic right whales (*Eubalaena glacialis*) ignore ships but respond to alerting stimuli. In Proceedings of the Royal Society of London. *Series B: Biological Sciences*, 271, 227-231.

- NSF & USGS (National Science Foundation & US Geological Survey). (2011). Programmatic environmental impact statement/overseas environmental impact statement for marine seismic research funded by the National Science Foundation or the US Geological Survey.
- Nunny, R., Graham, E., & Bass, S. (2008). Do sea turtles use acoustic cues when nesting? NOAA Technical Memorandum NMFS SEFSC No. 582:83. Retrieved from <http://www.nmfs.noaa.gov/pr/pdfs/species/turtlesymposium2005.pdf>. Accessed December 7, 2017.
- NWP Offshore Ltd. (2007). North Hoyle Offshore Wind Farm Annual FEPA Monitoring Report (2005-6). March (2007). [Online]. Retrieved from: <https://www.innogy.com/web/cms/mediablob/en/3170614/data/3170712/1/rwe-innogy/rwe-innogy-uk/sites/wind-offshore/in-operation/north-hoyle/environment/1.-Executive-summary-and-contents-.pdf>. Accessed: June 2018. 9 pp.
- NYSERDA (2010). Pre-development assessment of avian species for the proposed Long Island New York City offshore wind project area.
- NYSERDA (2015). Advancing the Environmentally Responsible Development of Offshore Wind Energy in New York State: A Regulatory Review and Stakeholder Perceptions NYSERDA Report 15-16.
- Oakley, B.A. and Boothroyd, J.C., 2012. Reconstructed topography of southern New England prior to isostatic rebound with implications of total isostatic depression and relative sea level. *Quaternary Research*, doi:10.1016/j.yqres.2012.03.002.
- O'Hara, J., & Wilcox, J. R. (1990). Avoidance responses of loggerhead turtles, *Caretta caretta*, to low frequency sound. *Copeia*, 1990(2), 564-567.
- Oldale, R.N. 1992. Cape Cod and the Islands: The Geologic Story. Parnassus Imprints, Orleans, Massachusetts, 208 pp.
- Olsen, J., & Olsen, P. (1980). Alleviating the impact of human disturbance on the breeding peregrine falcon II: Public and Recreational Lands. *Corella* 4:54-57.
- Orr, T. L., Herz, S.M. & Oakley, D.L. (2013). Evaluation of Lighting Schemes for Offshore Wind Facilities and Impacts to Local Environments. *Report prepared under BOEM Contract M12-PD-00007*. [Online.] Retrieved from [https://www.fbo.gov/index?s=opportunity&mode=form&id=51b9df02523d759b04b160168567b62f&tab=core&\\_cview=0](https://www.fbo.gov/index?s=opportunity&mode=form&id=51b9df02523d759b04b160168567b62f&tab=core&_cview=0)
- Overholtz, W. J., & Tyler, A. V. (1985). Long-term responses of the demersal fish assemblages of Georges Bank. *Fisheries Bulletin*, 83(4), 507-520.

- Owen, M., & Black, J.M. (1990). Waterfowl Ecology. In. Chapman & Hall, New York, NY.
- Pace, R. M. III, Corkeron, P. J., & Kraus, S. D. (2017). State-space mark-recapture estimates reveal a recent decline in abundance of North Atlantic right whales. *Ecology and Evolution*. doi: 10.1002/ece3.3406.
- Palka, D. (2012). Cetacean abundance estimates in US Northwestern Atlantic Ocean waters from summer 2011 line transect survey. US Department of Commerce, Northeast Fisheries Science Center Reference Document, 12-29.
- Palsbøll, P. J., Allen, J., Berube, M., Clapham, P., Feddersen, T., Hammond, P., ... & Oien, N. (1997). Genetic tagging of humpback whales. *Nature*, 388, 767-769
- Pangerc, T., S. Robinson, P. Theobald, and L. Galley. (2016). Underwater sound measurement data during diamond wire cutting: First description of radiated noise. *Proceedings of Meetings on Acoustics*, 27: 040012.
- Parks, S. E., Johnson, M., Nowacek, D., & Tyack, P. L. (2011). Individual right whales call louder in increased environmental noise. *Biology Letters*, 7, 33-35.
- Parks, S. E., Ketten, D. R., O'Malley, J. T., & Arruda, J. (2007). Anatomical predictions of hearing in the North Atlantic right whale. *Anatomical Record*, 290, 734-744.
- Paton, P. (2016). Assessing nearshore and offshore movements of Piping Plovers in southern New England. In Presentation at the 2016 North American Ornithological Conference (NAOC). Washington, D. C.
- Payne, K., & Payne, R. S. (1985). Large-scale changes over 17 years in songs of humpback whales in Bermuda. *Zeitschrift für Tierpsychologie*, 68, 89-114.
- Payne, P. M., & Heinemann, D. W. (1990). A distributional assessment of cetaceans in the shelf and shelf edge waters of the northeastern United States based on aerial and shipboard surveys, 1978-1988. Woods Hole, Massachusetts.
- Payne, P. M., & Heinemann, D. W. (1993). The distribution of pilot whales (*Globicephala* sp.) in shelf/shelf edge and slope waters of the northeastern United States, 1978-1988. *Reports of the International Whaling Commission (Special Issue)*, 14, 51-68.
- Payne, P. M., & Selzer, L. A. (1989). The distribution, abundance, and selected prey of the harbor seal, *Phoca vitulina concolor*, in southern New England. *Marine Mammal Science*, 5, 173-192.



- Payne, P. M., Selzer, L. A., & Knowlton, A. R. (1984). Distribution and density of cetaceans, marine turtles and seabirds in the shelf waters of the northeast US, June 1980 - Dec. 1983, based on shipboard observations. National Marine Fisheries Service, Woods Hole. NA81FAC00023: 245.
- Pelletier, S., Peterson, T., Boyden, S., Watrous, K. & Perkins, J. (2013). Ongoing offshore Atlantic & Great Lakes acoustic bat research. US Department of Energy Offshore Wind Webinar. Retrieved from: [http://mhk.pnnl.gov/wiki/images/0/00/Pelletier OSW Webinar 4 24 13.pdf](http://mhk.pnnl.gov/wiki/images/0/00/Pelletier_OSW_Webinar_4_24_13.pdf).
- Peng, C., Zhao, X., & Liu, G. (2015). Noise in the sea and its impacts on marine organisms. International Journal of Environmental Research and Public Health, 12(10), 12304-12323.
- Percival, S. M. (2010). Kentish Flats Offshore Wind Farm: Diver Surveys 2009-10. [Online.] Retrieved from [http://www.vattenfall.co.uk/en/file/Diver\\_follow-up\\_report\\_2010\\_\\_PDF\\_1970\\_kB\\_\\_16639636.pdf](http://www.vattenfall.co.uk/en/file/Diver_follow-up_report_2010__PDF_1970_kB__16639636.pdf)
- Perkins, S., Allison, T., Jones, A., & Sadoti, G. (2004). A Survey of Tern Activity Within Nantucket Sound, Massachusetts During the 2003 Fall Staging Period. Final Report to the Massachusetts Technology Collaborative, 10 September 2004.
- Perrin, W. F., Wursig, B., Thewissen, J. G. M. (eds.). (2002). Encyclopedia of marine mammals. Academic Press.
- Perrow, M. R., Gilroy, J.J., Skeate, E.R. & Tomlinson, M.L. (2011). Effects of the construction of Scroby Sands offshore wind farm on the prey base of Little tern *Sternula albifrons* at its most important UK colony. Marine Pollution Bulletin 62:1661–70. doi: 10.1016/j.marpolbul.2011.06.010
- Peters, E.C., Pilson, M.E.Q. (1985). A comparative study of the effects of sedimentation on symbiotic and asymbiotic colonies of the coral *Astrangia danae* Milne Edwards and Hime 1849. Journal of Experimental Marine Biology and Ecology, 92, 215–230.
- Petersen, I., & Fox, A. (2007). Changes in bird habitat utilisation around the Horns Rev 1 offshore wind project, with particular emphasis on Common Scoter. *Report commissioned by Vattenfall A/S*. [Online.] Retrieved from [http://www.vattenfall.se/sv/file/Horns-Rev-Habitat-Changes\\_11336653.pdf](http://www.vattenfall.se/sv/file/Horns-Rev-Habitat-Changes_11336653.pdf)
- Petersen, I. K., Christensen, T.K., Kahlert, J., Desholm, M. & Fox, A.D. (2006). Final results of bird studies at the offshore wind farms at Nysted and Horns Rev, Denmark. *Report commissioned by DONG Energy and Vattenfall A/S*. [Online.] Retrieved from [http://www.ens.dk/sites/ens.dk/files/undergrund-forsyning/vedvarende-energi/vindkraft-vindmoeller/havvindmoeller/idriftsatte-parker-nye/birds\\_final\\_2005.pdf](http://www.ens.dk/sites/ens.dk/files/undergrund-forsyning/vedvarende-energi/vindkraft-vindmoeller/havvindmoeller/idriftsatte-parker-nye/birds_final_2005.pdf).

- Petersen, I. K., Monique, M., Rexstad, E.A., Wisz, M.S., & Fox, A.D. (2011). Comparing pre- and post-construction distributions of long-tailed ducks *Clangula hyemalis* in and around the Nysted offshore wind farm, Denmark: a quasi-designed experiment accounting for imperfect detection, local surface features and autocorrelation. [Online.] Retrieved from <http://research-repository.st-andrews.ac.uk/handle/10023/2008>.
- Petersen, J.K., & Malm, T. (2006). Offshore windmill farms: Threats to or possibilities for the marine environment. *Royal Swedish Academy of Sciences*, 35, 75–80.
- Peterson, T. S., Pelletier, S. K., Boyden, S. A. & Watrous, K. S. (2014). Offshore acoustic monitoring of bats in the Gulf of Maine. *Northeastern Naturalist*, 21, 86-107.
- Pettersson, J. (2005). The impact of offshore wind farms on bird life in Southern Kalmar Sound Sweden final report based on studies 1999-2003. [Online.] Retrieved from [http://www.wind-energie.de/fileadmin/dokumente/Themen\\_A-Z/Vogelschutz/Voegel\\_Offshore\\_Sweden.pdf](http://www.wind-energie.de/fileadmin/dokumente/Themen_A-Z/Vogelschutz/Voegel_Offshore_Sweden.pdf)
- Pettersson, J., & Fågelvind, J. (2011). Night Migration of Songbirds and Waterfowl at the Utgrunden Off-Shore Wind Farm: A Radar-Assisted Study in Southern Kalmar Sound.
- Pettis, H. M., Pace III, R. M., Schick, R. S., & Hamilton, P. K. (2017). North Atlantic right whale consortium 2015 annual report card. Retrieved from <http://www.narwc.org/pdf/2017%20Report%20CardFinal.pdf>. Accessed on October 19, 2017.
- Pettit, J. L. & O'Keefe, J. M. (2017). Day of year, temperature, wind, and precipitation predict timing of bat migration. *Journal of Mammalogy*, gyx054.
- Piniak, W. E. D., Mann, D. A., Harms, C. A., Jones, T. T., & Eckert, S. A. (2016). Hearing in the juvenile green sea turtle (*Chelonia mydas*): A comparison of underwater and aerial hearing using auditory evoked potentials. *PLoS ONE* 11(10), e0159711.
- Plonczkier, P., & Simms, I.C. (2012). Radar monitoring of migrating pink-footed geese: behavioural responses to offshore wind farm development. *Journal of Applied Ecology* 49:1187–1194. doi: 10.1111/j.1365-2664.2012.02181.x
- Pollet, I. L., Shutler, D., Chardine, J.W., & Ryder, J.P. (2012). Ring-billed Gull (*Larus delawarensis*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY.
- Poot, H., Ens, B. J., de Vries, H., Donners, M.A.H., Wernand, M.R., & Marquenie, J.M. (2008). Green Light for Nocturnally Migrating Birds. *Ecology & Society* 13.

- Popper, A. N. & Fay, R. R. (1993). Sound detection and processing by fish: Critical review and major research questions. *Brain, Behavior and Evolution*, 41, 14-38.
- Popper, A. N. & Hastings, M. C. (2009). The effects of human-generated sound on fish. *Integrative Zoology*, 4(1), pp.43-52.
- Popper, A. N., Hawkins, A. D., Fay, R. R., Mann, D. A., Bartol, S., Carlson, ... & Tavolga, W. N. (2014). Sound exposure guidelines for fishes and sea turtles: A technical report prepared by ANSI - Accredited Standards Committee S3/SC1 and registered with ANSI. ASA Press/Springer, Cham. 88 pp.
- Popper, A. N., Smith, M. E., Cott, P. A., Hanna, B. W., MacGillivray, A. O., Austin, M. E., & Mann, D. A. (2014). Effects of exposure to seismic airgun use on hearing of three fish species. *The Journal of the Acoustical Society of America*, 117, pp. 3958-3971.
- Pratt, S. D. (1973). Benthic fauna. In: Coastal and offshore environmental inventory. Cape Hatteras to Nantucket Shoals. Marine Publication Series No. 2. University of Rhode Island, Kingston, R.I.
- Pritchard, P. C. H. (1997). Evolution, phylogeny, and current status. In: Lutz, P.L., & Musick, J. A. (eds.). *The biology of sea turtles*. Boca Raton, Florida: CRC Press. pp. 1-28.
- Provincetown Center for Coastal Studies. (2005). *Toward an Ocean Vision for the Nantucket Shelf Region*. Provincetown Center for Coastal Studies, Coastal Solutions Initiative. Retrieved from [http://www.coastalstudies.org/pdf/ocean\\_vis\\_rep.pdf](http://www.coastalstudies.org/pdf/ocean_vis_rep.pdf)
- Raoux, A., Tecchio, S., Pezy, J. P., Lassalle, G., Degraer, S., Wilhelmsson, D., Cachera, M., Ernande, B., Le Guen, C., Haraldsson, M., & Grangeré, K. (2017). Benthic and fish aggregation inside an offshore wind farm: Which effects on the trophic web functioning? *Ecological Indicators*, 72, pp.33-46.
- Rayment, W.J., (2002). Semi-permanent tube-building amphipods and polychaetes in sublittoral mud or muddy sand. Marine Life Information Network: Biology and Sensitivity Key Information Sub-programme [on-line]. Plymouth: Marine Biological Association of the United Kingdom. [cited 13/06/03]. Available from: <http://www.marlin.ac.uk>
- Read, P.A., Anderson, K.J., Matthews, J.E., Watson, P.G., Halliday, M.C., & Shiells, G.M. (1982). Water quality in the Firth of Forth. *Marine Pollution Bulletin* 13:421-425.
- Read, P.A., Anderson, K.J., Matthews, J.E., Watson, P.G., Halliday, M.C., & Shiells, G.M. (1983). Effects of pollution on the benthos of Firth of Forth. *Marine Pollution Bulletin* 14:12-16.

- Redfern, J. V., Hatch, L. T., Caldow, C., DeAngelis, M. L., Gedamke, J., Hastings, S., ... & Porter, M. B. (2017). Assessing the risk of chronic shipping noise to baleen whales off southern California. *Endangered Species Research*, 32, 153–167.
- Reeves, R. R., & Reed, A. J. (2003). Bottlenose dolphin, harbor porpoise, sperm whale, and other toothed cetaceans *Tursiops truncatus*, *Phocoena phocoena*, and *Physeter macrocephalus*. In Feldhamer, G. A., Thomson, B. C., & Chapman, J. A. (eds.). *Wild mammals of North American biology, management, and conservation*. pp. 397-424. Baltimore, Maryland: Johns Hopkins University Press.
- Reeves, R. R., Stewart, B. S., & Leatherwood, S. (1992). *The Sierra Club handbook of seals and sirenians*. San Francisco, California: Sierra Club Books.
- Reeves, R. R., & Whitehead, H. (1997). Status of sperm whale, *Physeter macrocephalus*, in Canada. *Can. Field Nat.*, 111, 293-307.
- Reichmuth, C., M.M. Holt, J. Mulsow, J.M. Sills, and B.L. Southall. (2013). Comparative assessment of amphibious hearing in pinnipeds. *Journal of Comparative Physiology A* 199: 491-507.
- Reine, K., D. Clarke, and C. Dickerson. (2012). *Characterization of Underwater Sound Produced by a backhoe dredge excavating rock and gravel*. (ERDC TN-DOER-E36). Vicksburg, MS: U.S. Army Engineer Research and Development Center. [www.wes.army.mil/el/dots/doer](http://www.wes.army.mil/el/dots/doer)
- Renaud, M.L. (1995). Movement patterns and submergence patterns of Kemp's ridley turtles (*Lepidochelys kempi*). *Journal of Herpetology*, 29(3), 370-374.
- Renski, H., Strate, S., Hodge, D., Proulx, W., Paik, K., Herter, S. (2015). *Long-term Population Projections for Massachusetts Regions and Municipalities*. University of Massachusetts. Amherst, MA.
- Rhoads, D. C., McCall, P. L., & Yingst, J. Y. (1978). Disturbance and production on the estuarine seafloor. *Am. Sci.*, 66, 577-86.
- Rhode Island Coastal Management Council. (2010). *Rhode Island Special Area Management Plan*. Wakefield, RI
- Rice, D. W. (1998). *Marine mammals of the world: Systematics and distribution*. Society for Marine Mammalogy, Special Publication No. 4. Lawrence, Kansas.
- Richardson, D. T. (1976). Assessment of harbor and gray seal populations in Maine 1974-1975. In Final Report to Marine Mammal Commission.

- Richardson, W. J. (1979). Southeastward shorebird migration over Nova Scotia and New Brunswick in autumn: a radar study. *Canadian Journal of Zoology* 57:107–124. doi: 10.1139/z79-009
- Richardson, W. J., Greene, C. R., Malme, C. I., & Thomson, D. H. (1995). *Marine mammals and noise*. San Diego, California: Academic Press.
- Ridgway, S. H., Wever, E. G., McCormick, J. G., Palin J., & Anderson, J. H. (1969). Hearing in the giant sea turtle, *Chelonia mydas*. *Proceedings of the National Academy of Sciences*, 64, 884-890.
- Riefolo, L., Lanfredi, C., Azzellino, A., Tomasicchio, G. R., Felice, D. A., Penchev, V., & Vicinanza, D. (2016). Offshore wind turbines: An overview of the effects on the marine environment. In The 26th International Ocean and Polar Engineering Conference. International Society of Offshore and Polar Engineers.
- Risch, D., Clark, C. W., Dugan, P. J., Popescu, M., Siebert, U., & VanParijs, S. M. (2013). Minke whale acoustic behavior and multi-year seasonal and diel vocalization patterns in Massachusetts Bay, USA. *Marine Ecology Progress Series*. 489, 279-295.
- Rogers, A. (2004). The biology, ecology and vulnerability of deep-water coral reefs. International Union for Conservation of Nature and Natural Resources. 10 pp.
- Rogers, B.A. (1969). Tolerance levels of four species of estuarine fishes to suspended mineral solids. Master's thesis. University of Rhode Island, Kingston.
- Rolland R. M., Parks, S. E., Hunt, K. H., Castellotte, M., Corkeron, P. J., Nowacek, D. P., ... & Kraus, S. D. (2012). Evidence that ship noise increases stress in right whales. *Proceedings of the Royal Society of London, Series B*, 279, 2363–2368
- Ronald, K., & Healey, P. J. (1981). Harp seal. In Ridgway, S. H., & Harrison, R. J. (eds.). *Handbook of marine mammals, Vol. 2: Seals*. New York, New York: Academic Press.
- Rone, B. K., & Pace, III, R. M. (2012). A simple photograph-based approach for discriminating between free-ranging long-finned (*Globicephala melas*) and short-finned (*G. macrorhynchus*) pilot whales off the east coast of the United States. *Mar. Mamm. Sci.*, 28(2), 254-275.
- Roos, P.C., and S.J.M.H. Hulscher. (2003). Large-scale seabed dynamics in offshore morphology: modeling human intervention. *Reviews of Geophysics* 41(2): 5-1 – 5-16.
- Rosel, P. E., Hansen, L., & Hohn, A. A. (2009). Restricted dispersal in a continuously distributed marine species: Common bottlenose dolphins (*Tursiops truncatus*) in coastal waters of the western North Atlantic. *Molecular Ecology*, 18, 5030-5045.

- Rosenberg, D. M., & Resh, V. H., eds. (1993). *Freshwater Biomonitoring and Benthic Macroinvertebrates*. New York: Chapman & Hall.
- Rosenfeld, M., George, M., & Terhune, J. M. (1988). Evidence of fall harbour seal, *Phoca vitulina*, movement from Canada to the United States. *The Canadian Field Naturalist*, 102(3), 527-529.
- RPS. (2012). *LID Year 3 Post-Construction Monitoring Summary Report*. Report to Centrica Energy Renewables Ref. LD-0-CE-013-0000-000000-324-D. December 2012.
- Rubega, M.A., Schamel, D. & Tracy, D. M. (2000). Red-necked Phalarope (*Phalaropus lobatus*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY. doi: 10.2173/bna.538
- Russel, D. F., Hastie, D., Thompson, V., Janik, P., Hammond, L., Scott-Hayward, J., ... & McConnell, B. (2016). Avoidance of wind farms by harbor seals is limited to pile driving activities. *Journal of Applied Ecology*, 53, 1642-1652.
- Rydell, J. & Wickman, A. (2015). Bat Activity at a Small Wind Turbine in the Baltic Sea. *Acta Chiropterologica*, 17, 359-364.
- Sammarco, P.W., Liretta, A., Tung, Y.F., Boland, G.S., Genazzio, M., & Sinclair, J. (2014). Coral communities on artificial reefs in the Gulf of Mexico: standing vs. toppled oil platforms. *ICES Journal of Marine Science*, 71(2), 417-426.
- Samuel, Y., Morreale, S. J., Clark, C. W., Greene, C. H., & Richmond, C. E. (2005). Underwater, low-frequency noise in a coastal sea turtle habitat. *Journal of the Acoustical Society of America*, 117(3), 1465-1472.
- Sanders, H. L. (1956). Oceanography of Long Island Sound. Chapter X. The biology of marine bottom communities. *Bulletin Bingham Oceanography Collection*, 15, 345-414. Retrieved from [http://peabody.yale.edu/sites/default/files/documents/scientific-publications/BOC15\\_06.pdf](http://peabody.yale.edu/sites/default/files/documents/scientific-publications/BOC15_06.pdf).
- Sanford, T. and Flick, R. (1975). On the relationship between transport and motional electric potentials in broad shallow currents. *Journal of Marine Research*, 33 (1), 123-139.
- Scheidat, M., Tougaard, J., Brasseur, S., Carstensen, J., van Polanen Petel, T., Teilmann J., & Reijnders P. (2011). Harbour porpoises (*Phocoena phocoena*) and wind farms: A case study in the Dutch North Sea. *Environmental Research Letters*. 6:025102. doi:10.1088/1748-9326/6/2/025102.

- Schneider, D., & Payne, P. M. (1983). Factors affecting haul-out of harbor seals at a site in southeastern Massachusetts. *Journal of Mammology*, 64(3), 518-520.
- Schreer, J., & Kovacs, K. (1997). Allometry of diving capacity in air-breathing vertebrates. *Canadian Journal of Zoology*, 75, 339-358.
- Schroeder, C. L. (2000). Population status and distribution of the harbor seal in Rhode Island waters. University of Rhode Island.
- Scott, T. M., & Sadove, S. S. (1997). Sperm whale, *Physeter macrocephalus*, sightings in the shallow shelf waters off Long Island, New York. *Mar. Mamm. Sci.*, 13, 317-321.
- SEFSC. (2017). Sea Turtle Stranding and Salvage Network (STSSN). Retrieved from <https://www.sefsc.noaa.gov/species/turtles/strandings.htm>. Accessed on November 22, 2017.
- Selzer, L. A., & Payne, P. M. (1988). The distribution of white-sided (*Lagenorhynchus acutus*) and common dolphins (*Delphinus delphis*) vs. environmental features of the continental shelf of the northeastern United States. *Marine Mammal Science*, 4(2), 141-153.
- Seminoff, J. A., Allen, C. D., Balazs, G. H., Dutton, P. H., Eguchi, T., Haas, H. L., ... & Waples, R. S. (2015). Status review of the green turtle (*Chelonia mydas*) under the US Endangered Species Act. NOAA Technical Memorandum, NOAA NMFS-SWFSC-539. 571pp.
- Sergeant, D. E. (1965). Migrations of harp seal *Pagophilus groenlandicus* (Erxleben) in the Northwest Atlantic. *J. Fish. Res. Board Can.*, 22, 433-464.
- Sergeant, D. E. (1977). Stocks of fin whales (*Balaenoptera physalus* L.) in the North Atlantic Ocean. Reports of the International Whaling Commission 27.
- Sergeant, D. E., Mansfield, A. W., & Beck, B. (1970). Inshore records of cetacea for eastern Canada, 1949-68. *Journal of the Fisheries Research Board of Canada*, 27, 1903-1915.
- Sherk, J.A., & Cronin, L.E. (1970). The effects of suspended and deposited sediments on estuarine organisms: an annotated bibliography of selected references. University of Maryland Natural Resources Institute, Reference 70-19, Solomons.
- Sherk, J.A., O'Connor, J.M., Neumann, D.A., Prince, R.D. & Wood, K.V. (1974). Effects of suspended and deposited sediments on estuarine organisms. Phase II. University of Maryland Natural Resources Institute, Reference 74-20, Solomons.
- Shimada, T., Limpus, C., Jones, R., & Hamann, M. (2017). Aligning habitat use with management zoning to reduce vessel strike of sea turtles. *Ocean and Coastal Management*, 142, 163-172.

- Shmookler, R. (2015). *Vessel Monitoring Systems (VMS) Commercial Fishing Density Northeast and Mid-Atlantic Regions*. Northeast Regional Ocean Council. Report available at: <http://www.northeastoceandata.org/files/metadata/Themes/CommercialFishing/VMSCommercialFishingDensity.pdf>
- Shoop, C. R. & Kenney, R. D. (1992). Distributions and abundances of loggerhead and leatherback sea turtles in northeastern US waters. *Herpetological Monographs*, 6, 43-67.
- Shortnose Sturgeon Status Review Team. (2010). A Biological Assessment of Shortnose Sturgeon (*Acipenser brevirostrum*). Report to National Marine Fisheries Service, Northeast Regional Office. November 1, 2010. 417 pp.
- Siemann, L. & Smolowitz, R. (2017). Southern New England Juvenile Fish Habitat Research Paper. OCS Study BOEM 2017-028. Retrieved from <https://www.boem.gov/ESPIS/5/5592.pdf>
- Sills, J. B. Southall, and C. Reichmuth. (2014). Amphibious hearing in spotted seals (*Phoca largha*): underwater audiograms, aerial audiograms and critical ratio measurements. *The Journal of Experimental Biology*, 217: 726-734.
- Sills, J. B. Southall, and C. Reichmuth. (2015). Amphibious hearing in ringed seals (*Pusa hispida*): underwater audiograms, aerial audiograms and critical ratio measurements. *The Journal of Experimental Biology*, 218: 2250-2259.
- Silverman, E. D., Leirness, J.B., Saalfeld, T., Koneff, M.D., & Richkus, K.D. (2012). Atlantic Coast Wintering Sea Duck Survey, 2008-2011.
- Silverman, E. D., Saalfeld, D.T., Leirness, J.B., & Koneff, M.D. (2013). Wintering Sea Duck Distribution Along the Atlantic Coast of the United States. *Journal of Fish and Wildlife Management* 4:178–198. doi: 10.3996/122012-JFWM-107
- Sjollemas, A. L., Gates, J. E., Hilderbrand, R. H. & Sherwell, J. (2014). Offshore Activity of Bats Along the Mid-Atlantic Coast. *Northeastern Naturalist*, 21, 154-163.
- Slater, M., Schultz, A., Jones, R., & Fischer, C. (2011). Electromagnetic field study. Oregon Wave Energy Trust. Retrieved from [file:///C:/Users/CourbisS/Downloads/OWET-EMF-on-Marine-Species\\_FINAL\\_Full\\_web1.pdf](file:///C:/Users/CourbisS/Downloads/OWET-EMF-on-Marine-Species_FINAL_Full_web1.pdf). Accessed November 19, 2017.
- Slavik, K., Lemmen, C., Zhang, W., Kerimoglu, O., and Wirtz, K. W. (2017). The large scale impact of offshore windfarm structures on pelagic primary production in the southern North Sea, *Hydrobiologia*. doi: 10.1007/s10750-018-3653-5.
- Smallwood, K. S. (2013). Comparing bird and bat fatality-rate estimates among North American wind-energy projects. *Wildlife Society Bulletin*, n/a.



- Smallwood, K. S. (2013). Long-term trends in fatality rates of birds and bats in the Altamont Pass Wind Resource Area, California.
- SMAST. (2016). "SMAST Benthic Survey Pyramid Study." Retrieved from <http://www.northeastoceandata.org/files/metadata/Themes/Habitat/AveragePresenceAbundanceSMAST.pdf>
- Smit, M. G., Holthaus, K. I., Trannum, H. C., Neff, J. M., Kjeilen-Eilertsen, G., Jak, R. G., Singaas, I., Huijbregts, M. A., & Hendriks, A. J. (2008). Species sensitivity distributions for suspended clays, sediment burial, and grain size change in the marine environment. *Environmental Toxicology and Chemistry*, 27(4),1006-12.
- Smolowitz, R. J., Patel, S. H., Haas, H. L., & Miller, S. A. (2015). Using a remotely operated vehicle (ROV) to observe loggerhead sea turtle (*Caretta caretta*) behavior on foraging grounds off the mid-Atlantic United States. *Journal of Experimental Marine Biology and Ecology*, 471, 84–91.
- Solan, M., Hauton, C., Godbold, J. A., Wood, C. L., Leighton, T. G., and White, P. (2016). Anthropogenic sources of underwater sound can modify how sediment-dwelling invertebrates mediate ecosystem properties. *Scientific Reports* 6, 20540.
- Solé M., Lenoir, M., Durfort, M., López-Bejar, M., & Lombarte, A. (2013), Ultrastructural Damage of *Loligo vulgaris* and *Illex coindetii* statocysts after Low Frequency Sound Exposure. *PLoS ONE*, 8(10): e78825.
- Southall, B. L. (2005). Shipping noise and marine mammals: A forum for science, management, and technology. Final report of the National Oceanic and Atmospheric Administration (NOAA) International Symposium. NOAA Fisheries, Arlington, Virginia. May 18-19, 2004. 40 pp.
- Southall, B., Bowles, A., Ellison, W., Finnerman, J., Gentry, R., Greene Jr., C., ... & Tyack, P. (2007). Marine mammal noise exposure criteria: Initial scientific recommendations. *Aquatic Mammals*, 33(4), 411-509.
- Southall, B. L., Dowacek, D. P., Miller, P. J. O., & Tyack, P. L. (2016). Experimental field studies to measure behavioral responses of cetaceans to sonar. *Endangered Species Research*, 31, 293–315.
- Spalding, M. D., Fox, H. E., Allen, G. R., Davidson, N., Ferdaña, Z. A., Finlayson, M., ... & Robertson, J. (2007). Marine ecoregions of the world: A bioregionalization of coastal and shelf areas. *BioScience*, 57(7), 573–583.
- Spiga, I., Caldwell, G.S., and Bruintjes, R. (2016). Influence of pile driving on the clearance rate of the blue mussel, *Mytilus edulis*. *Proceedings of Meetings on Acoustics*, 27: 040005. doi: 10.1121/2.

- Spotila, J. R., O'Connor, M. P., & Paladino, F. V. (1996). Thermal biology. In: Lutz, P. L. & Musick, J. A. (eds.). *The biology of sea turtles*. Boca Raton, Florida: CRC Press. pp. 297-313.
- Staine, K. J., & Burger, J. (1994). Nocturnal foraging behavior of breeding Piping Plovers (*Charadrius melodus*) in New Jersey. *The Auk* 111:579–587.
- Stanley, D.R. & Wilson, C.A. (1989). Utilization of offshore platforms by recreational fisherman and SCUBA divers off the Louisiana coast. *Bulletin of Marine Science*, 44(2), 767-775.
- Stantial, M. L. (2014). Flight behavior of breeding piping plovers: Implications for risk of collision with wind turbines.
- Stein, A. B., Friedland, K. B. & Sutherland, M. (2004). Atlantic sturgeon marine bycatch mortality on the continental shelf of the Northeastern United States. *North American Journal of Fisheries Management* 24, 171–183.
- Steinback, S. & Brinson, A. (2013). *The Economics of the Recreational For-hire Fishing Industry in the Northeast United States, 2<sup>nd</sup> ed.* Northeast Fisheries Science Center Social Sciences Branch, NOAA Fisheries. Woods Hole, MA
- Stenberg, C., Støttrup, J., Deurs, M. V., Berg, C. W., Dinesen, G. E., Mosegaard, H., Grome, T., & Leonhard, S. B. (2015). Long-term effects of an offshore wind farm in the North Sea on fish communities. *Marine Ecology Progress Series*, 528, pp.257-265.
- Stenhouse, I. J., Montevocchi, W.A., Gray, C.E., Gilbert, A.T., Burke, C.M., & Berlin, A.M. (2017). Occurrence and Migration of Northern Gannets Wintering in Offshore Waters of the Mid-Atlantic United States. In *Determining Fine-scale Use and Movement Patterns of Diving Bird Species in Federal Waters of the Mid-Atlantic United States Using Satellite Telemetry* (C. S. Spiegel, Editor). US. Department of the Interior, Bureau of Ocean Energy Management, Division of Environmental Sciences, Sterling, VA.
- Stokesbury, K. D. E. 2013. MA Windfarm Survey, Final Report. SMAST video survey of Western portion of the offshore Windfarm area, School for Marine Science and Technology (SMAST), University of Massachusetts Dartmouth, 13 pp.
- Stokesbury, K. D. E. 2014. MA Windfarm Survey, Final Report. SMAST video survey of Western portion of the offshore Windfarm area, School for Marine Science and Technology (SMAST), University of Massachusetts Dartmouth, 18 pp.
- Stoneburner, D.L. (1982). Satellite telemetry of loggerhead sea turtle movement in the Georgia bight. *Copeia*, 1982(2), 400-408.

- Strobel, S. M., & Mooney, T.A. (2012). Detection of low-frequency tones and whale predator sounds by the American sand lance *Ammodytes americanus*. *Journal of Fish Biology* 81:1646–1664. doi: 10.1111/j.1095-8649.2012.03423.x
- Sutcliffe, M. H., & Brodie, P. F. (1977). Whale distributions in Nova Scotia waters. Fisheries and Marine Service Technical Report, 722, 1-89.
- Swain, P.C., & Kearsley, J.B. (2001). Classification of the natural communities of Massachusetts. Retrieved from <http://www.mass.gov/eea/docs/dfg/nhosp/natural-communities-facts/nhcom-toc-ref.pdf>
- Swingle, W. M., Barco, S. G., Pitchford, T. D., McLellan, W. A., & Pabst, D. A. (1993). Appearance of juvenile humpback whales feeding in nearshore waters of Virginia. *Marine Mammal Science*, 9, 309-315.
- Taylor, D. A. R. (2006). *Forest management and bats*. Bat Conservation International, 13.
- Teilmann, J., & Carstensen, J. (2012). Negative long term effects on harbour porpoises from a large scale offshore wind farm in the Baltic—evidence of slow recovery. *Environmental Research Letters*, 7(4), 045101.
- Tempte, J. L., Bigg, M. A., & Wiig, O. (1991). Clines revisited: The timing of pupping in the harbour seal (*Phoca vitulina*). *Journal of Zoology*, 224, 617-632.
- TetraTech. (2012). Block Island wind farm and Block Island transmission system environmental report/construction and operations plan. Prepared for Deepwater Wind.
- TEWG. (2007). An assessment of leatherback turtles in the Atlantic Ocean. NOAA technical memorandum NMFS-SEFSC-555. 116 pp.
- TEWG. (2009). An assessment of the loggerhead turtle population in the western North Atlantic Ocean. NOAA technical memorandum. NMFS-SEFSC-575. 131 pp.
- Thaxter, C. B., Ross-Smith, V.H., Bouten, W., Clark, N.A., Conway, G.J., Rehfisch, M.M., & Burton, N.H.K. (2015). Seabird-wind farm interactions during the breeding season vary within and between years: A case study of lesser black-backed gull *Larus fuscus* in the UK. *Biological Conservation* 186:347–358.
- Theroux, R. B., & Wigley, R. L. (1998). Quantitative composition and distribution of the microbenthic invertebrate fauna of the continental shelf ecosystems of the northeastern United States. Technical Report NMFS 140 Washington, D.C.: US Dept. of Commerce, National Oceanic and Atmospheric Administration.

- Thompson, N.B. (1988). The status of loggerhead, *Caretta caretta*; Kemp's ridley, *Lepidochelys kempii*; and green, *Chelonia mydas*, sea turtle in U.S. waters. *Marine Fisheries Review*, 50(3), 16-23.
- Thompson, P. O., Cummings, W. C., & Ha, S. J. (1986). Sounds, source levels, and associated behavior of humpback whales, Southeast Alaska. *Journal of the Acoustical Society of America*, 80(3), 735-740.
- Thomsen, F. A. Gill, M. Kosecka, M. Andersson, M. Andre, St. Degraer, T. Folegot, J. Gabriel, A. Judd, T. Neumann, A. Norro, D. Risch, P. Sigray, D. Wood, B. Wilson. (2015). MaRVEN – Environmental Impacts of Noise, Vibrations and Electromagnetic Emissions from Marine Renewable Energy Final study report. Prepared for European Commission, Directorate General for Research and Innovation.
- Thomsen, F., Lüdemann, K., Kafemann, R., & Piper, W. (2006). Effects of offshore wind farm noise on marine mammals and fish. Biola, Hamburg, Germany on behalf of COWRIE Ltd, 62.
- Threatened Fish and Wildlife; Status Review of the Gulf of Maine/Bay of Fundy Population of Harbor Porpoise under the Endangered Species Act (ESA), 66 Fed. Reg. 40176 (August 2, 2001).
- Todd, C. S., Young, L. S., Owen, Jr., R.B., & Gramlich, F.J. (1982). Food habits of bald eagles in Maine. *Journal of Wildlife Management* 46:363–645.
- Tollit, D. J., Greenstreet, S. P. R., & Thompson, P. M. (1997). Prey selection by harbor seals, *Phoca vitulina*, in relation to variation in prey abundance. *Canadian Journal of Zoology*, 75, 1508-1518.
- Tougaard, J., Henriksen, O. D., & Miller, L. A. (2009). Underwater noise from three types of offshore wind turbines: Estimation of impact zones for harbor porpoises and harbor seals. *The Journal of the Acoustical Society of America*, 125, 3766-3773. doi: 10.1121/1.3117444.
- Tracy, D.M., Schamel, D. & Dale, J. (2002). Red Phalarope (*Phalaropus fulicarius*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY. doi: 10.2173/bna.698
- Turner, E.J. & Miller, D.C. (1991). Behavior and growth of *Mercenaria* during simulated storm events. *Marine Biology* 111:55-64.
- Tuttle, M. D. & Taylor, D. A. R.(1998). *Bats and mines*, Bat Conservation International.

- Tyack, P. (1986). Population biology, social behavior, and communication in whales and dolphins.
- Tyson, R. B., Piniak, W. E. D., Domit, C., Mann, D., Hall, M., Nowacek, D. P., & Fuentes, M. M. P. B. (2017). Novel bio-logging tool for studying fine-scale behaviors of marine turtles in response to sound. *Frontiers in Marine Science* 4: doi: 10.3389/fmars.2017.00219.
- United States Department of the Interior, Office of Renewable Energy Programs, Bureau of Ocean Energy Management (BOEM). (2016). Guidelines for information requirements for a renewable energy Construction and Operations Plan (COP), version 3.0. Retrieved from <https://www.boem.gov/COP-Guidelines/>
- United States Environmental Protection Agency (EPA). (2010). General conformity training module. Retrieved from [https://www.epa.gov/sites/production/files/2016-03/documents/general\\_conformity\\_training\\_manual.pdf](https://www.epa.gov/sites/production/files/2016-03/documents/general_conformity_training_manual.pdf)
- United States Environmental Protection Agency (EPA). (2015). Memorandum: Implementing the 2015 ozone National Ambient Air Quality Standards. Retrieved from [https://www.epa.gov/sites/production/files/2015-10/documents/implementation\\_memo.pdf](https://www.epa.gov/sites/production/files/2015-10/documents/implementation_memo.pdf)
- United States Environmental Protection Agency (EPA). (2017a). Greater Connecticut nonattainment area intended area designations for the 2015 ozone National Ambient Air Quality Standards technical support document. Retrieved from [https://www.epa.gov/sites/production/files/2017-12/documents/greater\\_ct.designations\\_120\\_day.tsd\\_final\\_.pdf](https://www.epa.gov/sites/production/files/2017-12/documents/greater_ct.designations_120_day.tsd_final_.pdf)
- United States Environmental Protection Agency (EPA). (2017b). New York-northern New Jersey-Long Island, NY-NJ-CT nonattainment area intended area designations for the 2015 ozone National Ambient Air Quality Standards technical support document (TSD). Retrieved from [https://www.epa.gov/sites/production/files/2017-12/documents/ny\\_nj\\_ct\\_new\\_york-northern\\_new\\_jersey-long\\_island\\_120d\\_tsd\\_final.pdf](https://www.epa.gov/sites/production/files/2017-12/documents/ny_nj_ct_new_york-northern_new_jersey-long_island_120d_tsd_final.pdf)
- United States Environmental Protection Agency (EPA). (2017c). Nonattainment areas for criteria pollutants (Green Book). Retrieved from <https://www.epa.gov/green-book>
- USACOE (US Army Corps of Engineers, New England District). (2001). Providence River and Harbor Maintenance Dredging Project. Final Environmental Impact Statement. Appendix L: Dredging Windows, August 2001.
- U.S. Coast Guard (USCG). 2012. Pollution Incidents In and Around U.S. Waters, A Spill/Release Compendium: 1969-2011: U.S. Coast Guard Marine Information for Safety and Law Enforcement (MISLE) System.

- USCG. (2009). Assessment of Potential Impacts to Marine Radar as it relates to marine navigation safety from the Nantucket Sound Wind Farm as proposed by Cape Wind, LLC. Retrieved from <https://www.boem.gov/Renewable-Energy-Program/Studies/USCGRADARfindingsandrecommendationsFINAL.aspx>
- USCG. (2006). Final Environmental Impact Statement for the Compass Port LLC Deepwater Port License Application. Volume 1 of 2. USCG 2004-17659.
- U.S. Department of the Navy. (2017). U.S. Navy Marine Species Density Database Phase III for the Atlantic Fleet Training and Testing Study Area. NAVFAC Atlantic Final Technical Report. Naval Facilities Engineering Command Atlantic, Norfolk, VA. 281 pp.
- USDOE MMS (US Department of Energy, Minerals Management Service. (2009). Final Environmental Impact Statement for the Proposed Cape Wind Energy Project, Nantucket Sound, Massachusetts (Adopted), DOE/EIS-0470. Retrieved from <https://www.boem.gov/Cape-Wind-FEIS/>
- US Fish and Wildlife Service. (n.d.), IPaC. Retrieved from <https://ecos.fws.gov/ipac/location/5ZAYYDWLYRH6PC4XGYJNSLVMOA/resources>
- US Fish and Wildlife Service (2016). 4(d) Rule for the Northern Long-Eared Bat. 50 CFR Part 17, Docket No. FWS–R5–ES–2011–0024; 4500030113. RIN 1018–AY98. Federal Register 81(9): 1900-1922.
- USFWS. (2008). Final Biological Opinion, Cape Wind Associates. [Online.] Retrieved from [http://www.fws.gov/newengland/pdfs/CapeWind-BO-21November2008\\_withCovLtrr.pdf](http://www.fws.gov/newengland/pdfs/CapeWind-BO-21November2008_withCovLtrr.pdf)
- USFWS. (2009). Piping Plover 5-Year Review: Summary and Evaluation. In. Hadley, Massachusetts and East Lansing, Michigan.
- USFWS. (2010). Caribbean Roseate Tern and North Atlantic Roseate Tern (*Sterna dougallii* dougallii) 5-Year Review: Summary and Evaluation.
- USFWS. (2014). West Indian manatee (*Trichechus manatus*) Florida stock (Florida subspecies, *Trichechus manatus latirostris*). Retrieved from [https://www.fws.gov/northflorida/manatee/SARS/20140123\\_FR00001606\\_Final\\_SAR\\_WIM\\_FL\\_Stock.pdf](https://www.fws.gov/northflorida/manatee/SARS/20140123_FR00001606_Final_SAR_WIM_FL_Stock.pdf). November 22, 2017.
- USFWS. (2015). Status of species: Red Knot (*Calidris canutus rufa*). Retrieved from [https://www.fws.gov/verobeach/StatusoftheSpecies/20151104\\_SOS\\_RedKnot.pdf](https://www.fws.gov/verobeach/StatusoftheSpecies/20151104_SOS_RedKnot.pdf)
- USFWS. (2017a). The Service’s independent “reasonable and prudent” determination regarding the originally proposed draft turbine curtailment measure in the Service’s October 2008 draft Biological Opinion (BO) for the Cape Wind project. Memorandum in response to decision.

- USFWS. (2017b). Piping Plover. Retrieved from <https://www.fws.gov/northeast/pipingplover/>
- USFWS. (2018). Massasoit National Wildlife Refuge, Draft Comprehensive Conservation Plan, unpublished.
- Vabø, R., Olsen, K., & Huse, I. (2002). The effect of vessel avoidance of wintering Norwegian spring-spawning herring. *Fisheries Research*, 58, 59–77.
- Valliere, A., Pierce, S. (2007). Southern New England industry-based yellowtail flounder survey: pilot study report. Retrieved from <http://www.dem.ri.gov/programs/bnatres/fishwild/pdf/ytibsrpt.pdf>
- Van Dalftsen, J. A., & Essink, K. (2001). Benthic community response to sand dredging and shoreface nourishment in Dutch coastal waters. *Senckenbergiana marit*, 31(2),329-32.
- Vanermen, N., Onkelinx, T., Courtens, W., Van de Walle, M., Verstraete, H., & Stienen, E. W. M. (2015). Seabird avoidance and attraction at an offshore wind farm in the Belgian part of the North Sea. *Hydrobiologia* 756:51–61. doi: 10.1007/s10750-014-2088-x
- Van Houtan, K. S., Halley, J. M., & Marke, W. (2015). Terrestrial basking sea turtles are responding to spatio-temporal sea surface temperature patterns. *Biology Letters*, 11, 20140744.
- Van Renterghem, T., Botteldooren, D., & Dekoninck, L. (2014). Airborne sound propagation over sea during offshore wind farm piling. *Journal of the Acoustical Society of America*, 135, 599-609.
- Vasconcelos, R. O., Amorim, M. C. P., Ladich, F. (2007). Effects of ship noise on the detectability of communication signals in the Lusitanian toadfish. *Journal of Experimental Biology*, 210, 2104-2112.
- Vavrinec, J., Pearson, W.H., Kohn, N.P., Skalski, J.R., Lee, C., Hall, K.D., Romano, B.A., Miller, M.C., and Khangaonkar, T.P. (2007). Laboratory Assessment of Potential Impacts to Dungeness Crabs from Disposal of Dredged Material from the Columbia River. Prepared for the U.S. Army Corps of Engineers, Portland District, Portland, OR under agreement with U.S. Department of Energy.

- Veit, R. (2015). Spatial association between seabirds and prey on the mid-Atlantic Outer Continental Shelf. *Wildlife Densities and Habitat Use Across Temporal and Spatial Scales on the Mid-Atlantic Outer Continental Shelf: Final Report to the Department of Energy EERE Wind & Water Power Technologies Office*. [Online.] Retrieved from [http://www.briloon.org/uploads/BRI\\_Documents/Wildlife\\_and\\_Renewable\\_Energy/MABS Project Chapter 10 - Veit 2015.pdf](http://www.briloon.org/uploads/BRI_Documents/Wildlife_and_Renewable_Energy/MABS_Project_Chapter_10_-_Veit_2015.pdf)
- Veit, R. R., Goyert, H. F., White, T. P., Martin, M.-C., Manne, L. L. Manne, & Gilbert, A. (2015). Pelagic Seabirds off the East Coast of the United States 2008-2013. US Dept. of the Interior, Bureau of Ocean Energy Management, Office of Renewable Energy Programs, Sterling, VA. OCS Study BOEM 2015-024. 186 pp.
- Veit, R. R., & Manne, L. L. (2015). Climate and changing winter distribution of alcids in the Northwest Atlantic. *Frontiers in Ecology and Evolution* 3:1–9. doi: 10.3389/fevo.2015.00038
- Veit, R. R., White, T. P., Perkins, S. A., & Curley, S. (2016). Abundance and Distribution of Seabirds off Southeastern Massachusetts, 2011-2015: Final Report. In. OCS Study. US. Department of the Interior, Bureau of Ocean Energy Management, Sterling, Virginia.
- Vlietstra, L. S. (2007). Potential Impact of the Massachusetts Maritime Academy Wind Turbine on Common (*Sterna hirundo*) and Roseate (*S. dougallii*) Terns.
- Vineyard Wind. (2016). Vineyard Wind protected species observer report 2016 geophysical and geotechnical surveys.
- Vineyard Wind. (2017). Vineyard Wind protected species observer report 2017 geophysical and geotechnical surveys.
- Vogt, R.J., Crum, T.D., Greenwood, W., Ciardi, E.J., & Guenther, R.G. (n.d.). *New Criteria for Evaluating Wind Turbine Impacts on NEXRAD Weather Radars*. NOAA, National Weather Service, Radar Operations Center. Norman, OK.
- Wahlberg, M. & Westerberg, H. (2005). Hearing in fish and their reactions to sounds from offshore wind farms. *Marine Ecology Progress Series*, 288, pp.295-309.
- Walker, M. (2001). Basslink project review of impacts of high voltage direct current sea cables and electrodes on chondrichthyan fauna and other marine life. Basslink Supporting Study No. 29, Marine and Freshwater Resources Institute No. 20.
- Walker, M. M., Diebel, C. E., & Kirschvink, J. L. (2003). Detection and use of the Earth's magnetic field by aquatic vertebrates. In Collins, S. P., & Marshall, N. J., (eds.). *Sensory processing in aquatic environments*. pp. 53-74. Springer, New York.



- Walker, M., Kirschvink, J., Ahmed, G., & Dizon, A. (1992). Evidence that fin whales respond to the geomagnetic field during migration. *Journal of Experimental Biology*, 171, 67-78.
- Walls, E.A., Berkson, J., Smith, S.A. (2002). The horseshoe crab, *Limulus polyphemus*, 100 years of study. *Reviews in Fisheries Science*, 10:39–73.
- Warchol, M. E. (2011). Sensory regeneration in the vertebrate inner ear: Differences at the levels of cells and species. *Hearing Research*, 273, 72-79.
- Waring, G. T., Josephson, E., Maze-Foley, K., & Rosel, P. E. (eds.). (2013). US Atlantic and Gulf of Mexico marine mammal stock assessments - 2012. NOAA Technical Memorandum NMFS-NE-223.
- Waring, G. T., Josephson, E., Maze-Foley, K., & Rosel, P. E. (eds.). (2014). US Atlantic and Gulf of Mexico marine mammal stock assessments - 2013. NOAA Technical Memorandum NMFS-NE-228.
- Waring, G. T., Josephson, E., Maze-Foley, K., & Rosel, P. E. (eds.). (2015). US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments - 2014. NOAA Technical Memorandum NMFS-NE-231.
- Warwick, R.M., & Uncles, R.J. (1980). Distribution of benthic macrofauna associations in the Bristol Channel in relation to tidal stress. *Marine Ecology Progress Series*, 3, 97-103.  
Retrieved from  
[http://researchrepository.murdoch.edu.au/id/eprint/22973/1/benthic\\_macrofauna\\_associations.pdf](http://researchrepository.murdoch.edu.au/id/eprint/22973/1/benthic_macrofauna_associations.pdf)
- Weilgart, L. S. (2007). The impacts of anthropogenic ocean noise on cetaceans and implications for management. *Canadian Journal of Zoology*, 85, 1091-1116.
- Westerberg, H. (2000). Impact Studies of Sea-Based Windpower in Sweden. - In: Merck, T. and H. von Nordheim, (eds.), *Technische Eingriffe in marine Lebensräume*. BFN-Skripten 29, Bundesamt für Naturschutz, 163-168.
- White, C. M., Clum, N. J., Cade, T. J., & Hunt, W. G. (2002). Peregrine Falcon (*Falco peregrinus*). In *The Birds of North America Online* No. 660. (A. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, p. 58. doi: 10.2173/bna.660
- White, T. P., Veit, R. R., & Perry, M. C. (2009). Feeding ecology of Long-tailed Ducks *Clangula hyemalis* wintering on the Nantucket Shoals. *Waterbirds* 32:293–299.
- Whitehead, H. (2002). Estimates of the current global population size and historical trajectory for sperm whales. *Mar. Ecol. Prog. Ser.*, 242, 295-304.

- Whitehead, H. (2003). Sperm whales: Social evolution in the ocean. The University of Chicago, Illinois: Chicago Press. 431 pp.
- Whitman, A. A., & Payne, P.A. (1990). Age of harbour seals, *Phoca vitulina concolor*, wintering in southern New England. *The Canadian Field Naturalist*, 104(4), 579-582.
- Whitt, A. D., Dudzinski, K., & Laliberté, J. R. (2013). North Atlantic right whale distribution and seasonal occurrence in nearshore waters off New Jersey, USA, and implications for management. *Endangered Species Research*, 20(1), 59-69.
- Wiese, F. K., Montevecchi, W. A., Davoren, G. K., Huettmann, F., Diamond, A. W., Linke, J., & Anonymous. (2001). Seabirds at risk around offshore oil platforms in the North-west Atlantic. *Marine Pollution Bulletin* 42:1285–1290.
- Wilber, D.H., & Clarke, D.G. (2001). Biological effects of suspended sediments: a review of suspended sediment impacts on fish and shellfish with relation to dredging activities in estuaries. *North American Journal of Fisheries Management*, 21, 855-875.
- Wilber, D.H., & Clarke, D.G. (2010). Dredging activities and the potential impacts of sediment resuspension and sedimentation on oyster reefs. Proceedings of the Western Dredging Association Technical Conference, June 6-9, 2010. San Juan, Puerto Rico, USA. pp. 61-69.
- Wilber, D. H., Brostoff, W., Clarke, D. G., & Ray, G. L. (2005). Sedimentation: potential biological effects of dredging operations in estuarine and marine environments. US Army Engineer Research and Development Center DOER-E20pg. 15 pp
- Wiley, D. N., Asmutis, R., Pitchford, T., & Gannon, D. (1995). Stranding and mortality of humpback whales, *Megaptera novaeanglia*, in the mid-Atlantic and southeast regions of the United States. *Fishery Bulletin*, 93, 196–205.
- Wiley, H. R., & Lee, D. S. (2000). Pomarine Jaeger (*Stercorarius pomarinus*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY. doi: 10.2173/bna.483
- Wiley, R. H., & Lee, D. S. (1998). Long-tailed Jaeger (*Stercorarius longicaudus*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY. doi: 10.2173/bna.365
- Wiley, R. H., & Lee, D.S. (1999). Parasitic Jaeger (*Stercorarius parasiticus*). In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY. doi: 10.2173/bna.445

- Williams, K., Connelly, E., Johnson, S., & Stenhouse, I. (Editors) (2015). Baseline Wildlife Studies in Atlantic Waters Offshore of Maryland: Final Report to the Maryland Department of Natural Resources and the Maryland Energy Administration. [Online.] Retrieved from [http://www.briloon.org/uploads/BRI\\_Documents/Wildlife\\_and\\_Renewable\\_Energy/MD Project 2015.pdf](http://www.briloon.org/uploads/BRI_Documents/Wildlife_and_Renewable_Energy/MD_Project_2015.pdf)
- Williams, R., Wright, A. J., Ashe, E., Blight, L. K., Bruintjes, R., Canessa, R., ... & Wale, M. A. (2015). Impacts of anthropogenic noise on marine life: Publication patterns, new discoveries, and future directions in research and management. *Ocean and Coastal Management*, 115, 17-24.
- Wilson, S. C. (1978). Social organization and behavior of harbor seals, *Phoca vitulina concolor*, in Maine.
- Wigley, R. (1968). Benthic invertebrates of the New England fishing banks. *Underwater Naturalist American Littoral Society* 5(1).
- Wood, J., B. L. Southall, and D. J. Tollit. (2012). PG&E offshore 3-D seismic survey project EIR—Marine mammal technical draft report. SMRU Ltd.
- Yasui, W. Y., & Gaskin, D. E. (2012). Energy budget of a small cetacean, the harbor porpoise, *Phocoena phocoena* (L.). *Ophelia*, 25, 183-197.